arXiv:math/0612375v1 [math.DG] 13 Dec 2006

THE METHOD OF ARCHIMEDES IN THE GEOMETRY OF QUADRICS

ION I. DINCĂ

Abstract. Confocal quadrics capture (encode) and geometrize spectral properties of symmetric operators. Certain metric-projective properties of confocal quadrics (most of them established in the first half of the XIX$^{\text{th}}$ century) *carry out* (stick and transfer) by rolling to and influence surfaces *applicable* (isometric) to quadrics and surfaces geometrically linked to these, thus providing a wealth of integrable systems and projective transformations of their solutions. We shall mainly follow Bianchi's discussion of deformations (through bending) of quadrics. Interestingly enough, *The Method* of Archimedes (lost for 7 centuries and rediscovered in the same year as Bianchi's discovery (1906), so unknown to Bianchi) applies *word by word in both spirit and the letter* and may provide the key to generalizations in other settings. Basically we have

Wisdom of the wise ancients+flat connection form=integrable systems.

Wisdom of the wise ancients+algebraic computations of the Bianchi Permutability Theorem=optimal discretization of integrable systems.

A four page appendix with the gist of the theory of deformations of quadrics is included as § 7 for the convenience of the rushed reader.

'It's only after we've lost everything that we're free to do anything.'
Tyler Durden

Contents

1. Introduction 1
1.1. Acknowledgements and full disclosure of the method of finding *The Method* 1
1.2. General remarks 10
1.3. Short historical perspective on quadrics 20
1.4. Statement of results 44
2. Algebraic preparatives for quadrics in $\mathbb{C}^n$ 52
2.1. Confocal quadrics in canonical form 52
2.2. Proofs of Lamé, Dupin, Ivory, Bianchi I, Jacobi and Chasles theorems; a ray of light 58
2.3. Proof of Bianchi II 59
2.4. Parametrization of confocal isotropic quadrics without center 63
2.5. Totally real quadrics in $\mathbb{C}^n$ 64
3. Algebraic preparatives for quadrics in $\mathbb{C}^3$ 65
3.1. Canonical confocal quadrics and homographies of Bianchi II 65
3.2. Totally real confocal quadrics in $\mathbb{C}^3$ and the corresponding homographies of Bianchi II 70
3.3. Two algebraic consequences of the tangency configuration 78
3.4. The Ivory affinity provides a rigid motion 79
3.5. Second iteration of the tangency configuration 81
3.6. Möbius configurations and an application of the Menelaus theorem 87
3.7. Doubly ruled Möbius configurations 91
4. The rolling problem 91
4.1. Submanifolds in $\mathbb{C}^n$ 91
4.2. Rolling surfaces 93

Key words and phrases: Bäcklund transformation, Bianchi Permutability Theorem, (confocal) quadrics, (discrete) integrable systems, *The Method* of Archimedes.
Supported by University of Notre Dame du Lac.

4.3. Curves corresponding under the applicability 98
4.4. Ruled surfaces 98
4.5. Infinitesimal deformations, Weingarten congruences, Darboux's 12 surfaces and transformations of surfaces 99
4.6. Rolling congruences 101
4.7. Rolling distributions 102
5. Deformations in $\mathbb{C}^3$ of quadrics in $\mathbb{C}^3$ 103
5.1. The Ricatti equation and the applicability correspondence provided by the Ivory affinity 103
5.2. Inversion of the Bäcklund transformation and the Weingarten congruence 105
5.3. Bianchi Permutability Theorem and moving Möbius configurations 109
5.4. Quadrics conjugate in deformation and the Hazzidakis transformation 111
5.5. (Singular) Bäcklund transformations of (singular) quadrics 112
5.6. (Totally real) deformations of totally real quadrics 115
5.7. Discrete deformations of quadrics 117
5.8. Delaunay's, Guichard's and Weingarten's results 118
5.9. Conic systems 120
6. Particular deformations in $\mathbb{C}^{n+p}$ of quadrics in $\mathbb{C}^{n+1}$ 121
6.1. Overview and particular deformations in $\mathbb{C}^{2+p}$ of quadrics in $\mathbb{C}^3$ 121
6.2. Tenenblat-Terng's Bäcklund transformation of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ 122
6.3. The Bianchi-Lie approach 126
6.4. Particular deformations in $\mathbb{C}^{2n-1}$ of quadrics in $\mathbb{C}^{n+1}$ 127
7. Appendix 1: A gist of deformations of quadrics 128
7.1. Introduction 128
7.2. Confocal quadrics 128
7.3. The Bäcklund transformation 129
7.4. Two algebraic consequences of the tangency configuration 130
7.5. Rolling surfaces and distributions 131
7.6. The method of Archimedes-Bianchi-Lie 131
References 134

1. Introduction

1.1. **Acknowledgements and full disclosure of the method of finding** *The Method*.
I am much indebted to the University of Notre Dame du Lac for academic support during the graduate program (in particular I am grateful that although my teaching duties began with the second year and mainly required four tutorials per week, most semesters I was required to hold only two; one semester even only one) and to my Advisor Professor Brian Smyth for useful discussions from Fall 2002 until Fall 2005 (he clocked two hours a week during the academic year and mostly four a week during the Summers; thus I had no choice but to present at the blackboard mostly the opinions of the classical geometers). Also I would like to thank my fellow graduate students (in particular to my officemates Daniel Cibotaru, Florin Dumitrescu and Daniel Jackson) for patiently listening and useful discussions.

I would also like to thank the Faculty of the Mathematics Department from Bucharest University for the free high quality undergraduate mathematical education (I was also granted a modest financial support).

Special thanks to Daniel Cibotaru for an enlightening discussion that took place in Fall 2005, when I was preparing a quiz for Calculus B and I noticed a behavior which suggested a smooth ride in a car with elliptic wheels and on a sinusoidal road (the textbook was stating that the arc-length of $(\cos(s), \sqrt{2}\sin(s))$ and that of $(s, \sin(s))$ are the same; this provides rolling). I knew

that the elliptic wheel has to be attached to the car at one of the foci before even trying to complete the computations, because replacing the circular wheel with the elliptic wheel was the 1-dimensional metric-projective analogy to replacing the pseudo-sphere with an arbitrary quadric for the 2-dimensional theory of isometric deformations of quadrics by means of rolling (we shall henceforth use the classical denomination *deformation* instead of *isometric deformation*). Thus I have stated my views to Daniel (clarifying them as I was talking to him) and I felt compelled to do the analytic confirmation and a Maple moving picture; the picture suggested that the wheel rotates faster when the focus is closer to the point of tangency, which led me to a book of history of mathematics and the first encounter with the wisdom of the wise ancients, who have mastered *'the science of the sections occurring in cones'*. It is this how I was able to give a possible answer to the question which was bugging me for some time, namely *Why one can do computations for quadrics?*; this was done only after I read and understood the early history of conics (up to including Newton), in the process simplifying the 2-dimensional discussion (I had all the facts, including the metric-projective denomination, but I did not put them together up to that point).

I thus kept in high esteem and added the early history of conics (including Archimedes' quote from *The Method*) to the notes in late December 2005-early January 2006; also I considered their quotation in these notes to be in the correct place because I was using a theorem of Menelaus in a place where Bianchi did not use it. At that time I saw in those quotes a confirmation of the fact that Bianchi was able to do computations for quadrics, so nothing more than a weak analogy; I was looking at them as everybody does: statements of historical importance. Unfortunately in what it seems to be my biggest blunder so far I reduced the early history of quadrics to less than a page by the end of January 2006 (thus removing Archimedes' quote in the process) and I did not read the early part of the notes up to August 2006 with the same frequency and intensity I did before January 2006 (I used to do that on a bad day, get my spirits up and do some work in the meat of the notes).

In mid-August 2006 I drank a beer with Daniel and I stated my hopes that with Levi-Civita's intuitive notion of parallel transport taken as an obvious axiom and the methods developed by Apollonius in *Conics* one can provide a synthetic proof of the theory of deformations of quadrics, thus reducing it from a proof Leibnitz would understand to a proof the wise ancients would understand; again these things became more clear to me as I was stating them to Daniel. After this discussion I noticed that at some point I was stating in that version of these notes *'All necessary identities of the moving part boil down to valid relevant algebraic identities of the static part; in turn these algebraic identities naturally appear within the static part and admit simple geometric interpretations.'* The *'naturally'* appeared from an analytic point of view (changing the ruling family), but at some point my memory failed in my favor, as I did not remember the change of ruling family, but just the reflection property. But if the reflection property naturally appeared, why would I continue to say *'and admit simple geometric interpretations'*, as the natural reflection property was itself the simple geometric interpretation (and in trying to see why it is natural I found the rolling of a surface on both sides of another applicable surface also applies in the singular case of the Bianchi-Lie ansatz: if one rolls the seed on both sides of the considered quadric, then one gets immediately the reflection property). Thus I tried to understand what I wrote; in doing so I realized that the reflection property *as a natural geometric approach* also works at the level of the Bianchi-Lie ansatz *by means of an intuitive non-rigorous argument*.

So this was the setting in late August 2006 when I had a discussion with the Chair (according to The Oracle in the *'Matrix Trilogy': 'We cannot see beyond the choices we do not understand'*); I believe I even mentioned to the Chair that again I have ideas (between January 2006 and August 2006 I just run on fumes, trying to complete a program clearly and previously established); on that day the idea was only a fancier point of view on the 2-dimensional discussion, so it would not bring new results in that setting. On the very same afternoon I went home and I decided to bring the notes to the earlier January 2006 organization (I believe I also met Florin on campus on the way home; he is interested in super parallel transport and I believe I informed him on that occasion of

Levi Civita's simple point of view). I remembered Archimedes' quote and the fact that I liked it, so first thing I did was to put it back in the notes. By reasons of internal organization of these notes Archimedes' quote stood for the first time alone, separate of its intended use in the earlier version of the notes. I wanted to see if it is correctly organized, so I read the statement above it, then Archimedes' quote, but I never got to the statement below: as soon as I read it something stuck to my mind and I had to investigate. This is how I made the connection between Archimedes and the classical geometers of the XIX$^{\text{th}}$-XX$^{\text{th}}$ century.

Now even if I saw this simplification (which may have helped me look closer at Archimedes' quote) before making the connection with Archimedes, Archimedes' point of view is still the one which must have priority, as *the first one*; on top of that Archimedes took it up a notch. I did not expect to get what I was looking for before the first attempt to seriously look at the ideas of the wise ancients as statements of current usefulness instead of statements of historical usefulness. So I got the first interpretation of Archimedes' quote by accident (it was separated from its initial place in the earlier version of the notes because I added the comment about Archimedes being generous *in disclosing the full method of investigation*; had I've put it back to its original place, then probably I would have not made the connection, as I read and re-read this quote for one month in late December 2005- early January 2006 thinking nothing else but the situation at hand, namely Archimedes' specific applications of his method). Further and maybe more irrationally than rationally I almost saw the *actual* interpretation because *I just obeyed the command shown in front of my eyes*; thus I further investigated (as Jules would say not all signs are of a divine nature, but by use of common sense, Fonzie cool and simple logic the false ones can be easily removed as such) and it was clearly revealed to me that the big fuss about what I saw and Bianchi did not see, about what I did and Bianchi did not do would be decided by giving the credit (as usually) to *a third neutral party*. What hit me was the fact that Archimedes told me that there is no spot of trouble in using the singular case *full Bianchi-Lie ansatz* just as the non-singular case *Theorem I of Bianchi's theory of deformations of quadrics*: thus basically Archimedes solved the most difficult step of the theory of deformations of quadrics 2200 years before anybody got any inkling about such things. What also hit me was the power of generality of *The Method* (the only way Archimedes could have accomplished this deed), so I tried it immediately on the higher dimensional problem and instantaneously I got *the first confirmation that the higher dimensional problem is amenable to an attack strategy*: this is my first important application of Archimedes' ideas.

Again *The Arnold Principle* worked out just fine (luckily just on time for me; I felt uneasy about criticizing Bianchi and praising myself instead; I believe even Bianchi would have approved of Archimedes' opinion on the matter), just as *'the inconceivable effectiveness of mathematics in natural sciences'*.

Of course I was glad that it worked out this way (confounded bitter-sweet-dark-comedy-*'Twilight-Zone'-'dazed-and-confused'* to be more precise), but several days later I remembered stories with mathematicians seeing nonexistent hidden messages or talking to the aliens, so I realized that there may be another rational explanation: thus I emailed an alien (Daniel) to confirm at least my intention of seriously reading the wise ancients and Daniel gracefully confirmed (although a little bit confounded; when he inquired me later for more explanations and in order to cool him down I made a bet with a beer towards implementing the ideas of the wise ancients in what I am doing; this bet is to be discounted because I already knew the outcome and Daniel had already proven me right: the key to the problem at hand was sitting on the counter the whole time; one must only have to look straight at it while having fresh in the memory the Bianchi-Lie ansatz in order to see it).

Archimedes is not only the best mathematician of the antiquity and his opinions are not only of historical importance, but useful in the study of current problems. Thus I would like to also thank Archimedes for his choice of well crafted words (I had an early interpretation, but it still did not fully comply with Archimedes' written point of view, so I had to further refine it until, I believe, Archimedes was completely satisfied).

The miracle of the *Archimedes-Bianchi-Lie* (ABL) method is that Archimedes put *The Method* in the Bianchi-Lie ansatz, previously used in a hurry as a strange defining axiom and rapidly discarded as being a nonsense with no other usefulness. Bianchi and Lie could not use the full method (although probably they would have approved its use in the exact wording of Archimedes), because it was a nonsense probably even according to their own beliefs. Thus Bianchi in ([4],§ 374) points out a singular behavior and takes part of it as a defining axiom. Moser in [24] uses cautious wording and provides a number of examples; it may be the case that Moser points out Archimedes' method, so the rabbit hole goes even deeper (but becomes slimmer) and in a full circle the study of the geometry of quadrics returns to its true beginnings and not just to the beginning of the $\mathrm{XIX}^{\mathrm{th}}$ century: integrable system means nothing more and nothing less than cases when one can apply Archimedes' method of integration. It is only the French and their abstract tendencies (see Arnold [2]) that led Darboux to dare state such nonsense as a surface of revolution being deformed to a line, but this was still not enough: one needs a line being *an actual* deformation of a surface of revolution.

Basically *The Method* of Archimedes states that $P \Rightarrow Q$ still give elegant, relevant, useful and valid consequences Q, even if P is a statement not accepted as being true by the standards of proof of the time. Such an example of Q is the statement *'E pur si move!'* or the correct exponent obtained by the scaling trick to find the needed exponent for various inequalities involving Sobolev spaces before proving those inequalities. Or in general all tricks used to see if a statement is true (*reductio ad absurdum*), without attacking directly the full proof of the validity of that statement (usually very difficult); when the tricks do not point out an error it means that one must muster courage to attack the full proof. Moreover the whole initial effort of trying to disprove P must always be kept in mind, instead of being thrown away once its apparent usefulness has been exhausted. Most of the time P will turn out to be true, even if not according to the standards of proof of the time. Note that simple logic says $\neg Q \Rightarrow \neg P$; when it is frustrating that Q is headstrong enough to remain true, that is actually a good sign that P is true, just as happened with Euclid's fifth postulate.

Luckily Archimedes had faith that P is true, even though he could not prove it and he tried *The Method* on numerous examples to confirm his beliefs and intuition, before sending a letter to Erathostenes of the University of Alexandria. To this day Archimedes' sentence P is not proven *'such as the ancients required'*, but instead and independently of Archimedes allowed as a foundational axiom by the founding fathers of Calculus Leibnitz and Newton (although the buzz around Archimedes' palimpsest is that Archimedes may have already performed the trick).

Archimedes does not mind stating that he uses an unproven sentence P, as he had already secured his back and reputation with enough examples pointing to its validity; thus we have a great mind (he coined the wording with the Master's perfection) being honest and that is all that a novice needed 2300 years later. I have tried initially to pay only partial attention to the Master's opinion only to see later that the Master's opinion was perfect and I was wrong in interpreting it.

The beauty of *The Method* of Archimedes is that it works in some cases with same efficiency even if P is not accepted as true by the standards of proof of the time and has a very small chance of being accepted as true by future generations: this is the message I got from Archimedes. So Archimedes told me to assume that a ruling on a confocal quadric is a deformation of the considered quadric and not to pay attention to such a *'ridiculous'* statement, but only to its consequences. Before Archimedes I justified the reflection property of the considered distributions in the tangent bundle of the considered quadric (a simple geometric procedure, but it does not appear naturally from a geometric point of view, as Bianchi did not think about it) with changing the ruling family of the confocal quadric (and thus naturally from an analytic point of view); also the rigid motion provided by the Ivory affinity naturally appeared from an analytic point of view, but not from a geometric point of view, as even I could not think about using ridiculous statements (at some point and while trying to find vacuum solitons I thought about using this rigid motion to see the applicability correspondence between a ruling on a confocal quadric and a region of the considered quadric: the corresponding region is a ruling on the considered quadric, but the planes of the facets are just

tangent planes to the considered quadric along the ruling, each counted with an ∞ multiplicity, so I dropped this nonsense immediately). At some point the stronger version of Darboux's statement was present in these notes as versions equivalent to it and obtained by rolling, but that just because I had a vague idea about Darboux's statement; when challenged to remove the stronger statement I knew that I had to ask Monsieur Darboux about his opinion on the matter.

Thus I saw the geometric properties of the moving picture as consequences of the geometric properties of the static picture, but the later naturally appeared only from an analytic point of view (changing the ruling family and the use of a certain rigid motion as to whose appearance I did not even dare question). After Archimedes the geometric properties of the static picture are consequences of the geometric properties of the moving picture and thus naturally appear from a geometric point of view (see again Arnold [2]).

Principles of a general nature transcend singularities and remain valid even for singularities; moreover such principles recorded for singularities are the only important information needed to confirm the validity of the initial principles of a general nature; such ideas can be found in the current literature of integrable systems (like Hilbert-Riemann or scattering data) or maybe in other domains and Archimedes' *The Method* is the first example (so Archimedes basically proved that the balance with fixed fulcrum survives integration and the only important thing to do is to find the correct balance configuration at the infinitesimal level by use of simple Euclidean properties of the studied object: *he transferred and stuck slices (the equivalent of infinitesimal leaves or facets) at their center with ∞ multiplicity at the fixed left end of the balance*; this is *precisely* the above mentioned nonsense of Bianchi-Lie's and thus Archimedes' principle of the balance surviving integration also applies to the theory of deformations of quadrics).

Certain properties of certain structures remain valid even when the *subjacent* (supporting) structure collapses to nothingness, to ridiculous, to irrational (even the method of finding the link to *The Method* of Archimedes is an application of the same method, since an irrational hypothesis (signs are shown to me) has a rational and perfectly founded conclusion), thus providing not a paradox or a false ridiculous statement, but the simplest explanation of those structures.

The area of a segment of length $f(x)$ is of course 0, but if that segment is a limit of thinning rectangles of same height $f(x)$, something sticks to it and it has a memory containing more information: the area is $f(x)dx$, an infinitesimal which by itself and for all practical purposes is of course 0, but it is definitely something different from 0. So this is the centuries old story, told and retold to high-school kids in some countries (and to college level kids in others).

Archimedes not only said essentially the same story more than two thousand years ago, *but he also gave its geometric description* and it is the power of geometry that makes Archimedes' ideas amenable to generalizations in another settings, just as *'the inconceivable effectiveness of mathematics in natural sciences'* , usually realized by a model supported on a simple geometric picture. Bianchi's contemporaries acclaimed *'Qui la geometria vive!'* and yet they did not know the whole story.

The beauty of the ABL method is that it gives almost for free more information from the beginning, so I believe it to be the essential ingredient to supporting generalizations in other settings (especially in higher dimensions); in dimension 2 is just a fancier reformulation of the old point of view. Of course the information already given in dimension 2 by the old point of view may be generalized with a heroic effort to higher dimensions without Archimedes' point of view, but Archimedes' point of view is the natural one, it appearing from a *geometric* point of view instead of an *analytic* point of view. And it is over the course of history that *not* the geometric picture subjacent to physical facts (like gravity and light), *but* its analytic interpretation changed. On top of that it bypasses complicated computations since it gives for free *all* algebraic identities of the static picture necessary for the completion of the differential identities required by the moving picture, instead of trying to complete first the later and in doing so see what are the former. To do this trick we need to find the correct discretization of the said differential identities.

Consider the theory of deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ from Tenenblat-Terng [32].

From the application of the ABL method one gets immediately the $\mathbf{O}_{n-1}(\mathbb{R})\times\mathbf{O}_{n-1}(\mathbb{R})$ symmetry due to the symmetries (in the normal bundle) of the rolling and to the symmetry of the tangency configuration. But these symmetries never use the fact that we work with $\mathbf{H}^n(\mathbb{R})$, so are valid for the deformations of a general n-dimensional real quadric in $\mathbb{R}^{2n-1}$ (Berger, Bryant, Griffiths [3]), should such a theory of deformation be developed in this case. For my birthday in Spring 2006 the true gift was not the congratulatory phone call from Romania, but finding the article of Berger, Bryant and Griffiths (I believed it to be true, began with Cartan [12] and followed the trail, only to find that the most difficult part (namely the existence of the seeds of the theory) had already been proven). Although it is not clear to me if Berger, Bryant and Griffiths proved the flat normal bundle property, it should follow as a simple intuitive consequence of the *initial value data* of the differential system subjacent to the Bäcklund transformation (counting the leaves; their numbers are the same as for Tenenblat-Terng since we just generalize their point of view from a metric-projective point of view), since no functional information is allowed in the normal bundle except a finite $\mathbf{O}_{n-1}(\mathbb{R})$ dimensionality.

Note that the Bianchi Permutability Theorem from Terng [33] is the correct discretization of the differential equations required by existence of deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$, that is the proof *'such as the ancients required'*. The metric-projective properties of this picture can be assumed to be true at the level of the static picture for deformations of n-dimensional real quadrics in $\mathbb{R}^{2n-1}$ by means of the ABL method and get the needed algebraic information (it seems that the necessary algebraic identities required by the existence of the B transformation are encoded by the second iterated tangency configuration and those required by the existence of the Bianchi Permutability Theorem are encoded by the third iteration of the tangency configuration; no further algebraic identities are needed).

Most continuous groups of symmetries and symmetric spaces are given by quadratic equations; some of their metric properties may not be due to the big group of symmetries, but to the quadratic definition: we call such properties *metric-projective*. Of course to deform such things one needs a surrounding space with a big group of symmetries and certain simplifying assumptions.

According to Terng-Thorbergsson [34]: *'...while extending the theory of sub-manifolds to ambient spaces more general than space forms proves quite difficult if one tries to use the same approach as for the space forms, at least for symmetric spaces it has proved possible to develop an elegant theory based on focal structure that reduces to the classical theory in the case of space forms.'*

Thus should a theory of deformations of higher dimensional general quadrics be developed in the Euclidean space, the only important thing remaining to do is the construction of the confocal family of a quadric in a symmetric space; probably this can be easily done so as to satisfy the usual classical metric-projective properties.

As to the undergraduate education at Bucharest University in what concerns the prerequisites from my higher mathematical education needed to read the classical geometers I wish to thank Professor Kostake Teleman for the first year Geometry course (where it was discussed almost all necessary prerequisites for the static picture for quadrics including vectorial, affine and projective spaces and their transformations; bilinear and multilinear forms; eigenvalues and eigenvectors; diagonalization of symmetric real and hermitian matrices; Sylvester's theorem; metric, affine and projective classification of real quadrics; projective classification of complex quadrics; real confocal quadrics; Bézout's theorem for algebraic curves; etc), Assistant Victor Vuletescu for being both the Instructor and the Assistant for the first semester of the second year Geometry course (where the classical geometry of curves and hyper-surfaces was summarily covered), Professor Martin Jurchescu for the second year Complex Analysis course (there was also the Assistant I do not remember the name, but I met him at Notre Dame first in Summer 2004; I distinctly remember he did a problem with a Schwartz reflection symmetry and a problem on the exam was almost instantaneously solved with a rotational $\mathbb{Z}_3$ symmetry plus the Schwartz reflection symmetry and a Cauchy continuity at the origin), for the third year Calculus on Manifolds course (when it was clearly revealed to me by Guillemin-Pollack that any abstract manifold can appear as a real Euclidean manifold in Poincaré's

interpretation) and for the fourth year Multivariable Complex Analysis course (the Romanian Dimitrie Pompeiu's generalization of Cauchy's point of view is just a fancier reformulation for one variable, but essential for more than one variable); unfortunately he got sick that year; Assistant George Marinescu took over the second semester of the course and completed it with the proof of Nirenberg-Newlander. Now the teaching method in Romania is on the French model: students go to lectures (which are not discussion style, but rather the lecturer putting a whole Rudin, Ahlfors or Schwartz on the blackboard out of memory, of course in one's own interpretation, just as a conductor sometimes allows oneself freedom in interpreting the message of the composer (meantime the audience keeps quiet) or just like QT allowed himself changing the time-line of *'Pulp Fiction'*; the favor is asked back from the student at the end of the semester) and tutorials for three quarters of the semester and have only tough *written* final examinations in the exams session of a quarter of a semester (roughly one exam per week) and somehow easier *written* examinations in a special exams session when students may choose at the pleasure of their own heart to challenge at most two of the first grades; as a consequence I still remember the page of the Complex Analysis course I was reading when, being frustrated with what I was reading, I decided together with my roommate to throw water bags from the dorm window hoping that the frustration will *stick to the water bags and transfer* to the poor passers who happened to walk below our window and conversely.

At that time my undergraduate Advisor Professor Stere Ianuş was doing harmonic maps, so he gave me a book of Eells-Rato on harmonic maps with symmetries and positive curvature targets to write the BS thesis (so all research should begin by writing some notes to oneself where one explains in detail already known results in an area of consistent research); in doing so I also almost understood the article of Eells-Sampson (it is considered the founding article of the branch of harmonic maps; using the heat flow it provided existence of harmonic maps in negative curvature targets). For my MS thesis at Bucharest University I remember reading Lawson's minimal surfaces, sub-manifolds and currents as examples of harmonic maps (I also met with Monsieur Lawson in a course of Spin Geometry); as I was leaving Romania Professor Stere Ianuş intended for me to read something about loop groups (after trying a stint with isoparametric sub-manifolds; I believe I saw something but by the time I investigated it *'I lost it, Lou!'*); I found out later that these have something to do with deformations of quadrics. Now harmonic maps with positive curvature targets (for example spheres or ellipsoids) are very difficult to study; consequently one must assume initially lots of symmetries to simplify the problem. The Ph. D. thesis of Smith in early 1980's did just that (note also that that of Wood's showed that the Schwartz reflection symmetry is valid for harmonic maps in dimension 2): CMC surfaces have harmonic Gauß map into the unit sphere $\mathbb{S}^2$; if one assumes this Gauß map to have rotational $\mathbb{S}^1$ symmetry, then one gets Delaunay's CMC surfaces of revolution (this observation on Smith's results for 2-dimensional target is due to Calabi). Note that if one assumes the rotational $\mathbb{S}^1$ symmetry only at the level of the linear element, then one gets more CMC surfaces than those of Delaunay's: such surfaces were found by Smyth in 1987, are complete, always have an umbilic and outer dihedral symmetry (so the Gauß map does not have rotational symmetry). Recently Dorfmeister, Pedit, Wu and collaborators have analyzed CMC surfaces with the loop groups tools and in particular have generalized these Smyth surfaces to various other settings.

Finally on a very short list of people which had the most benefic influence on my mathematical training by going beyond what was required of them I would like to thank my mother Floarea Dincă, *Învăţătorul Român* incarnated for me as my primary school teacher Didiţa Staicu, my 7^{th} grade mathematics teacher Lucia Velicu, my formerly high-school mathematics teacher (currently Professor Dumitru Popa of Ovidius University Constanţa), the former mathematics Inspector Gheorghe Popescu of the Ialomiţa County, the *Romanian Education Foundation* of Stanford, CA and the weekly *Academia Caţavencu*. Further this last part of the acknowledgements is definitely biased and incomplete, it being designed to suit a restricted *educational* point of view; besides that it has a personal touch (according to Tyler Durden *'You want to make an omelet, you got to break some eggs'* and it is the mouth of the sinner (the undersigned) which spells out the truth: one can

recognize the power of generality of mathematics when the same model describes a-priori unrelated situations).

It is because my mother has paid a subscription to the monthly *Gazeta Matematică* ever since I was in the 5^{th} grade that I had later things to munch on. She was my best Advisor although she had no education beyond primary school and I did not always agree with her findings (note however that I being a minor and she being my legal tutor she was my only Advisor ever who could *legally* choose and did chose notably in Summer 1990 to impose on me one's opinion first and foremost of mine). It is from her that I learned the useful lesson of *standing and defending my ground 'lest The Heavens will fall!'* (just as should God meet with the Satori sword from *'Kill Bill'* (designed according to the Romanian Gheorghe Zamfir's pan-flute instructions mind you), God Himself will be cut, should I smell that God is trying to pull a fast one on me I will hit God Himself with a confirmation delivery letter (*So Help Me God!*), because the receipt of Uncle Sam's Postal Services is the alien's only friend on Uncle Sam's territory): one of our neighbors' son rose to the rank of party leader of the Buzău county; consequently our neighbor decided to encroach on our territory and my mother took the problem to a county court of law, where of course she lost. Her lack of education (her father denied his girls education beyond primary school, as he considered that *Bărăgan* (the plain with a mind of its own and where the harsh Siberian winter wind roams free; on a good rainy year it provides the bulk of the Romanian grain which in turn used to decide the price of grain in Europe until WWII; on a sequence of bad dry years it almost starved to death its poorest inhabitants; note also that it gave to the Romanians their first unifier *Mihai Viteazu*, Flower's child of Floarea in the lost City of Flocks situated 10 miles east of Ţăndărei and from where Ţăndărei probably *split*) would provide the remaining needed education; it is only after the change to communist regime in Romania that two of her younger sisters completed at least high school education) may be the possible explanation of the fact that all her four children (boys) got a college education (beginning with August 1975 she was the head of the family and consequently worked low paying jobs as unqualified worker at a local brick factory until she retired in Summer 1992 on a low pension on account of health reasons and worked in our privately-owned agriculture during the Springs). In Summer 1995 I was almost defeated by a string of bad grades; just like Ştefan cel Mare's mother put some faith into him when he was almost defeated by the Turks, she put some faith into me, I went back and I aced the PDE exam (it was not a perfect score, but luckily for me the Instructor decided to round the score upwards before seeing my student ID and my other grades); looking back I can honestly say that no formal education was needed after that exam, as I have not seen since then interesting tricks, methods and tools after being hit that year with infinite dimensional spaces (Banach, Hilbert and locally convex) and the essence of symmetries (Lie Algebras); the *Fourier transform* words were not uttered until its natural existence Shwartz's space of tempered distribution was properly introduced. Now the custom and only publicly recognized form of corruption in Romania is to bribe *The Godfather* (Comptroller) with 30% of the train ticket price; the briber (usually student or other small income frequent train user social categories, because one needs both motivation and experience; the new student user is helped by The experienced Godfather's nonchalance in receiving the bribe and The beginner Godfather by the experienced student's nonchalance in offering the bribe) risks a civil fine up to $5-10$ times the price of the ticket if caught ticket-less by The Uber-Comptroller, that is the inspector of The Godfather's work; it seems that even The Csar of Russia initially refused the help of The Romanian Army in the 1877-1878 Russo-Turkish war on account that a nation which does not have the decency of buying a ticket before boarding a train cannot be trusted as an ally in waging war; I took full advantage of this custom as a student (in the Summers and early Falls I was doing agriculture at Ţăndărei and during the academic year I was carrying food to Bucharest every fortnight) and never paid a fine; the only time when I came close to paying one (The Uber-Comptroller confiscated our student ID's) The Wolf of Ţăndărei happened to pass in the neighborhood and solved the problem by producing a railway worker ID and shaming The Uber-Comptroller into total submission; usually when The Uber-Comptroller meets a group of students

he also meets resistance and consequently after difficult negotiations he can bestow at most a fine fine to the whole group; thus *the power lies within the group, at least five'o'em* (in which case one gets almost even).

It is because of Didiţa Staicu that I discovered the thrill of *reading and writing mathematics*: just like Jules she had an *'ultimate piercing'* look and would not let yours escape once caught and would pierce and shake the very foundations of your soul; consequently I still have trouble looking her in the eyes when we meet on the street (that may be also due to my not so proud to be of deeds).

It is because of Lucia Velicu that I let aside the life of mischief and tormenting my teachers and our neighbors from the Strachina neighborhood (I was first of my class, but my deeds were still best described by Ion Creangă's *'Memories of my childhood'*) and began doing serious mathematics: she organized a weekly off-curriculum mathematics club. I remember my first idea to generalize $\sqrt{2^4} = 2^2$ was $\sqrt{2^n} = 2^{n-2}$ (it is customarily to remember mostly unpleasant situations although the name itself explains: square root of a is the root of x squared equals a; even the Babilonians knew Newton's method on how to approximate square roots thousands of years before Newton got any linking about this method). Lucia asked us in Fall 1986 to buy a book of problems; luckily my colleague and good friend Cătălin lent me some money (first and last time I touched money not belonging to me was at age of 8 when I was playing hide-and-seek with Cătălin and his sister in their home; I hid near the porcelain pig and consequently some of their money *stuck to my sticky hands and transferred to my pockets*; luckily his father chose just to pull my ears the next morning when I met Cătălin as usually on our way to school and without bringing my deed to the knowledge of my mother) and I bought the last book from the bookstore; I read many answers to the geometry problems from the answer section of that book on the evening before the county competition in Spring 1987, *as they were explained at large* (the problem with the current mathematician is that *one's mathematics is difficult*; note however that that difficulty may also arise from assuming the student to be familiar with unfamiliar background, letting the student fill in *'the details'* by oneself as an unnecessary homework assignment (thus virtually *rebuilding in a costly masochistic fashion* the already built theory instead of *investing a fraction of the same time* in understanding the same, namely the full method of investigation) or from failing to motivate and reinforce by means of simple examples the purpose of the investigation (according to Number One of *'Austin Powers Goldmember': 'Throw me a ... bone!'*) and asking at each step of the investigation the motivation of that step: taking definitions and other things for granted produces *the sleep of ration*); I distinctly remember Euler's 9 points. Thus in Spring 1987 I barely missed the national mathematics competition, but as a consequence I acceded to county mathematics programs beginning with Fall 1987. In Spring 1988 I got almost the perfect score for the geometry examination of the national mathematics competition; in the algebraic examination I remember guessing a solution of an equation which after some change of variable (skilful handling at that time) became a quadratic equation in x^2, so it had four solutions.

It is because of Professor Dumitru Popa that I continued doing serious mathematics: he took me under his wing at a bad time in my life (I had just miserably failed the Romanian Literature and Grammar high-school entry examination; on top of that The Wolf caught me cheating first time in my life on the first English test in high-school (in a full circle last time I cheated was on the last test in high-school; it was French and I was caught for the second time only); these may have also motivated me to pursue mathematics with renewed vigor in order to redeem myself); also he imposed his will on other teachers and always arranged that I should have at least one week off school before mathematics competitions (or, as a recognition of my efforts, a whole month in 1989 after returning from a national program in mathematics), a thing I sorely missed during my last two years in high-school (in Fall 1991 *I declined participation to an intercounty mathematics competition on account of not being prepared, as no due one week notice was given to me and instead I was fighting for passing grades at subjects of study such as geography technology* (drawing maps and memorizing names of Romania's mountains)). It does not matter that the group of three-four

students preparing for the mathematics competition were in fact frustrated with the mathematics problems and instead were playing ping-pong or various games with chalk or coins during class time only to show diligence before and during recesses, when we were inspected to show progress on our work, because that work was enough (an impromptu inspection even caught us in the act of throwing darts at the door, as the door opened to let the inspector of our work do one's duty). And even playing with coins can have benefic consequences: probably Ţiţeica was playing with the change while drinking beer with friends and thus noticed the problem with four equal coins (a law of a general nature, probably amenable to interesting generalizations for conics or even in higher dimensions; it actually has something to do with *Barbilian geometries*).

It is because of former mathematics Inspector Gheorghe Popescu of the Ialomiţa County that I acceded to special national programs in mathematics in Romania: due to a certain sequence of events he chose to settle a dispute by choosing *a third neutral resolution*: he bet on my geometrical abilities in Spring 1989, in the process bending some rules.

Interestingly enough in Spring 1989 I misunderstood a difficult algebraic problem which took most of the competitors most of their time to solve; I gave in few minutes a faulty solution and as a consequence I had most of the remaining time to spend on the other three problems. The remaining problem with an algebraic character was similar to a problem which had been previously discussed by Silviu Ciupercă (formerly mathematics teacher at the Ţăndărei high-school and currently turned politician) at a county mathematics program in early 1989 (unfortunately the last such county mathematics program ended *by necessity* on the very same day with the communist regime in Romania). For one of the geometry problems I was asked to provide a synthetic proof to a metric-projective result of Apollonius (the plane locus of points with constant ratio of distances to two fixed points of the plane is a circle; when the ratio is 1 the circle passes through the line at ∞ and becomes a line in the Euclidean plane). I knew Ţiţeica's book of Euclidean geometry almost by heart, since for 4-6 hours of classes a day I was spending 9 hours on the train and waiting in the Slobozia train station hallway (I could have travelled around the equator with 40 miles a day for almost four years). I had to turn back 16 years later to Apollonius and his contemporaries, in the process discovering Arnold [2] in early 2006 and realizing that I was right in my convictions for the last three years.

It is because the *Romanian Education Foundation* paid my TOEFL and GRE general test fees that I found later a lender to borrow the money for the GRE mathematics test fee and thus I acceded to a Ph. D. program in mathematics in USA (I got this REF sponsorship tip from a fellow student one year senior to me; unfortunately their first letter was *'O scrisoare pierdută'* (A lost letter, as many other letters coming to Romania from abroad; some even reach Hungary on account of Bucharest being its capital city and thus Budapest being the capital of Romania) and did not reach my dorm address; after contacting them again the second one reached my permanent address, but due to the delay and not being able to find a lender by early Summer 1996 I had to postpone the application process with one year).

It is because the weekly *Academia Caţavencu* paid the airline ticket that I was able to show myself at such a Ph. D. program (I got this sponsorship tip from a fellow student, who with the same sponsorship was able to show himself at a summer school in Italy in 1997; on top of that their motto is *'Our readers are brighter than their readers'*, so I pressured them to deliver on their ideas). That would not have probably been necessary, as I had other options (including offers to borrow money from some of my fellow students and Faculty of the Mathematics Department of the Bucharest University), but being in debt with a smaller amount of money gave me peace of mind.

1.2. **General remarks.**

As much as possible and always keeping in mind an optimal mixture of clarity, convenience and simplicity we shall choose our working definitions, notions, style of presentation, etc in decreasing order of importance beginning from the classical ones, followed by the current ones, followed by our choices, but the later always not far removed from the context. Most likely the later are not the best choice: for example *adjugate* was *adjoint* and before that *dual*.

Take for example the use of the classical denomination *applicable* instead of the current one *isometric*; the former acknowledges that the surfaces naturally exist in a surrounding space, while the later pays attention only to the induced linear elements. For the same reason we use *deformation* instead of *isometric deformation*, as there is no such later thing (the induced linear element stays the same, so the abstract surface itself (obtained by deletion of the surrounding space) does not undergo any deformation). Another choice would be *deformation through bending*, but this implies continuity of the process and *deformation through warping* is already too complicated (see Sabitov [27] or Spivak ([29], Vol **5**)). For the same reason we use *Gauß -Weingarten equations* instead of *fundamental equations of immersion*, since they are consequences of the existence of a sub-manifold in a surrounding space and not necessary equations required for the existence of an isometric immersion of an abstract sub-manifold in another one. The main point of the theory of deformations of quadrics is that we always assume the existence of an initial (*seed*) deformation in a surrounding space.

In the spirit of the classical geometers we shall preserve the thread of exposition of facts, without (numbered) theorems, lemmas, etc.

Due to the frequent use of certain key words we shall define and use abbreviated notations.

Due to the frequent use of the reference Bianchi ([5],Vol **4**,(122,...)), it will be replaced with Bianchi (122,...). Note also that Bianchi ([4],Vol 2, Part 1) contains three chapters about deformations of quadrics; it may be the last revised version of Bianchi's.

References are restricted to the ones I am aware of being of direct interest here. There is a huge literature on surfaces and integrable systems and for example Rogers-Schieff [26] has a consistent list of references, from where one can probably get in at most one more step to much of the current literature in this area. As usual, unwarranted absence of credit does not imply claim of credit, but only that the particular fact is considered common knowledge or that I am not aware of such correlations of names and facts. A further hinder in this matter is the fact that the classical geometers and their results are not considered mainstream mathematics today; seldom there are monographs in various branches of geometry which give short descriptions of the results of the classical geometers in their respective historical perspective sections. Consequently the current geometer trusts only part of the results of the classical geometers and safely assumes that the only logical conclusion that can be drawn from the fact that notions such as rolling, the use of imaginaries, etc are not present in the current literature is that there is something fundamentally wrong with these; I had my share of doubts on the use of imaginaries and to this day I still believe such an explanation may be possible, but until *I find such a valid argument* I will side with the classical geometers. Note that a big part of the classical references is freely available online: for example Bianchi ([5],Vol 2) and an earlier one volume version from 1894, Darboux [13] and Eisenhart [20] can be found inThe University of Michigan Historical Mathematics Collection. Unfortunately from Archimedes' collected works (written in the $\mathrm{XIX}^{\mathrm{th}}$ century) is missing exactly the needed piece, but again *The Method* is freely available on the web.

There is no replacement of Bianchi's works and the present notes do not intend this; they may be used in a parallel reading, as there are many other ideas and points of view in Bianchi's works that do not appear here. While a simpler proof is sometimes useful, ideas from the original proof can be used in other settings; this is another point of these notes.

The sine-Gordon equation $\omega_{uv} = \sin\omega$ appears as the Gauß equation of the *Gauß-Codazzi-Mainardi* (GCM) equations (the integrability conditions of the *Gauß-Weingarten* (GW) equations) for *constant Gauß curvature* (CGC) -1 surfaces, where (u, v) are *Tchebyshev coordinates* (coordinates for which u is the arc-length parameter on the v =const curves and conversely; that is the taylor's problem with a piece of fabric) which further are asymptotic and ω is the angle between the asymptotic directions; the Codazzi-Mainardi equations are identically satisfied in this setting. Note that the complete CGC -1 surface can be realized as a connected component of a particular quadric (the *space-like pseudo-sphere*) in the Lorentz space of signature $(2, 1)$. The *space-like*

refers to the induced linear element being positive definite (equivalently the normal is time-like; the pseudo-sphere itself is seen as time-like from the origin, since all normals pass through the origin).

While the sine-Gordon equation received much attention in recent years, I am not aware of any similar treatment of deformations of general quadrics: Moser discusses in [24] only the geodesic flow on quadrics and points out analytic similarities between it and other integrable systems (like the *Korteweg-de Vries* (KdV) equation), while Burstall in the monograph [9] on *isothermic* surfaces (surfaces with isothermic lines of curvature) does not discuss the link to deformations of quadrics. In [36] Terng and Uhlenbeck discuss, via a highly technical Lie algebraic formalism, a natural generalization of the classical *Bäcklund* (B) transformation of CGC -1 surfaces: the B transformation of the *Zakharov, Šabat, Ablowitz, Kaup, Newell, Segur* (ZS-AKNS) $\mathbf{sl}_n(\mathbb{C})$ hierarchy. Although they do not specifically mention Bianchi's *Bäcklund* (B) transformation of deformations of general quadrics, there are similarities; thus it may be the case that their approach contains Bianchi's B transformation as a particular case, but only from an analytical point of view and without the simpler geometric picture. Note however that hierarchies must be constructed at the analytic level, as at this point any geometric meaning is above me; here again Terng-Uhlenbeck [36] may already have done that.

The *Bäcklund* (B) transformation of local solutions of an integrable system is a procedure of generating *leaves* (new local solutions) from a *seed* (given local solution). This procedure requires the integration of a simpler auxiliary differential system associated to the given local solution. Sometimes this is called an *auto*-B transformation, following that a *general* B transformation may be a procedure of generating local solutions of an integrable system from local solutions of another integrable system, again requiring the integration of a simpler auxiliary differential system (there are also algebraic procedures with the same purpose as the general B transformation, classically known as *Hazzidakis* (H) transformations, but currently as *Miura* transformations). Since we use only the auto-B transformation, we shall call it B transformation.

By necessity (all quadrics are equivalent from a complex projective point of view) we shall consider the complexification $(\mathbb{C}^n, <, >)$ of the Euclidean space $(\mathbb{R}^n, <, >)$: $< x, y >:= x^T y,\ |x|^2 := x^T x,\ x, y \in \mathbb{C}^n$. *Isotropic* (null) vectors are those vectors of length 0 : $|v|^2 = 0$; since most vectors are not isotropic we call *vector* a vector presumed non-isotropic and we shall only emphasize *isotropic* when the vector is assumed to be isotropic. This terminology will also apply in other settings: for example we call *quadric* a non-degenerate quadric, following that we shall only emphasize *degenerate* when the quadric is assumed to be degenerate. Standard geometric formulae for sub-manifolds in $\mathbb{R}^n$ remain valid (with their usual denomination) almost everywhere. If the Euclidean product on $\mathbb{C}^n$ induces a non-degenerate product on the complexification of a tangent space (this complexification is assumed to be a proper subspace of $\mathbb{C}^n$) of a general sub-manifold in $\mathbb{C}^n$, then one can find an orthogonal complement spanned by an orthonormal frame. Since the GW and the *Gauß-Codazzi-Mainardi-Ricci* (GCMR) equations for a sub-manifold $x \subset \mathbb{R}^n$ assume only the non-degeneracy of the linear element of x (and thus the existence of a moving orthonormal frame in the normal bundle), they are still valid and sufficient to describe the geometry of general sub-manifolds in $\mathbb{C}^n$ almost everywhere. For example the Lorentz space of signature $(2, 1)$ can be realized as the *totally real subspace* $\mathbb{R}^2 \times i\mathbb{R} \subset \mathbb{C}^3$ (the induced scalar product is real (valued) and non-degenerate). Note that there are many definitions of totally real subspaces, some involving a hermitian product, but all definitions coincide. All 3-dimensional totally real affine subspaces of $\mathbb{C}^3$ are obtained by applying a rigid motion $(R, t) \in \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3,\ (R, t)x := Rx + t,\ x \in \mathbb{C}^3$ to one of the Lorentz spaces $\mathbb{R}^3,\ \mathbb{R}^2 \times i\mathbb{R},\ \mathbb{R} \times (i\mathbb{R})^2,\ (i\mathbb{R})^3$.

We call *surface* any sub-manifold of $\mathbb{C}^3 \simeq \mathbb{R}^6$ of real dimension $2, 3$ or 4 such that all its *complexified tangent spaces* (called henceforth *tangent spaces*) have complex dimension 2. In this case the distribution $T \cap iT$ formed by intersecting real tangent spaces T with iT has constant real dimension (which must respectively be $0,\ 2$ or 4). This distribution is integrable (by its own definition); by an application of Frobenius one obtains leaves; on leaves the Nijenhuis condition holds and by Nirenberg-Newlander this is equivalent (locally) to prescribing $x : D \to \mathbb{C}^3$, where D is a domain of

$\mathbb{R}^2$ or $\mathbb{C}\times\mathbb{R}$ or $\mathbb{C}^2$ such that $dx\times\wedge dx\neq 0$ (the $\times$ applies to the vector structure and the $\wedge$ to the exterior form structure, so the order does not matter: $\times\wedge=\wedge\times$; note also that $dx\times\wedge dx=2dx\times dx$, but we also need at some point $dx\times\wedge dy$). The GW and GCM equations suffice to describe the geometry of most surfaces almost everywhere and surfaces such defined are the natural completion of the usual real surfaces, although the usual coordinates used in geometry (asymptotic, conjugate systems, orthogonal, principal) may not be the ones which clearly split into purely real and purely complex (for example asymptotic coordinates on a real surface of positive Gauß curvature). The change of coordinates still holds, even outside bi-holomorphic change of the complex coordinates and diffeomorphism of the real ones, but if the real and complex coordinates are mixed, they lose the character of being purely real or complex and the new parametrization may not be holomorphic in some variables. For example if we have a surface $x=x(z,t)\subset\mathbb{C}^3,\ (z,t)\in D\subseteq\mathbb{C}\times\mathbb{R}$ and consider two complex functions $u=u(z,t),\ v=v(z,t)$ of z and t with $J:=\frac{du\wedge dv}{dz\wedge dt}=\begin{vmatrix}\frac{\partial u}{\partial z} & \frac{\partial u}{\partial t}\\ \frac{\partial v}{\partial z} & \frac{\partial v}{\partial t}\end{vmatrix}\neq 0$, then one can invert $z=z(u,v),\ t=t(u,v)$ at least formally and at the infinitesimal level by the usual calculus rule of taking the inverse of the Jacobian: $dz=\frac{\partial z}{\partial u}du+\frac{\partial z}{\partial v}dv,\ dt=\frac{\partial t}{\partial u}du+\frac{\partial t}{\partial v}dv$. Although the holomorphic dependence on a variable is lost, the 2-dimensionality character remains clear. After using formal coordinates of geometric meaning to simplify certain computations, one can return to the purely real and complex coordinates, just as the use of $z,\bar{z}$ parametrization on real surfaces. Note that although on such surfaces the vector fields $\partial_z,\ \partial_{\bar{z}}$ do not admit integral curves (lines of coordinates), statements about infinitesimal behavior of such lines remain valid, so one can assume such lines to exist and derive corresponding results; thus for all practical purposes we can assume that coordinates descend upon lines of coordinates on the surface. From a practical point of view usually one needs only surfaces of real dimension 2 or 4; real 3-dimensional surfaces usually appear in the intermediary step of obtaining real 2-dimensional totally real surfaces from real 4-dimensional surfaces; they are true real 3-dimensional when appear as real 1-dimensional families of holomorphic curves and this family cannot be extended to a holomorphic 1-dimensional family; surfaces of revolution and ruled surfaces should provide the simplest examples.

This goes a little bit against the intuition, as for a real surface in the real Euclidean space $\mathbb{R}^6$ the second fundamental form is vector valued and we also need the Ricci equations. But keep in mind the fact that the Euclidean scalar product on $\mathbb{C}^3$ is completely different: it has a real and an imaginary part, both being indefinite scalar products of signature $(3,3)$ on $\mathbb{R}^6$. If we begin with a real analytic surface, then one can easily complexify not only the tangent bundle, as one does currently, but also the surface and the surrounding space; thus the asymptotic coordinates (complex conjugate $z,\bar{z}$) on a real analytic surface of positive Gauß curvature will admit integral curves on the corresponding surface parameterized by $z,\ w$ and the initial real surface is individuated by the non-real condition $w=\bar{z}$ (which implies $z=\bar{w}$; these two functionally independent conditions can be considered for computations instead of two real functionally independent conditions imposed on the four real parameters of the surface in order to get the real surface). Thus one can easily derive lines of length 0 (isotropic curves) on real analytic surfaces; this was my first encounter with imaginaries when I read Eisenhart [19] in Fall 2002, but I ignored the use of imaginaries for one more semester. The *surfaces of translation* (sum of two curves) with generating curves being isotropic conjugate are precisely the minimal surfaces and this fact is due, according to Lie, to Monge. When the curves are holomorphic and one also restricts the parameter on one to be the conjugate of the parameter on the other one gets *all real minimal surfaces*; this is one step before the *Weierstrass representation formula* (so Weierstrass essentially used the standard parametrization of the isotropic cone), recently generalized to other settings.

It is unclear to me at this point if the classical geometers considered only analytic surfaces; anyway except for the results where they specifically use the power series expansion the remaining ones remain valid in a more general setting. By the end of the XIX$^{\text{th}}$ century existence of solutions of differential systems in more general settings appeared (for example Picard's existence of solutions of ODE's), so the classical geometers were aware of constructions more general than analytic ones.

As a graduate student being involved in a program of the study of the classical geometry of surfaces I discovered over the course of roughly one year the beautiful (although difficult to read at the first cursory incursion; probably due also to the fact that for example I discovered in the Library first Vol **2**, then Vol **3,4** and then Vol **1** of Darboux [13]) Bianchi [4],[5] and Darboux [13],[14] and I was introduced to the theory of deformations of quadrics, one of the crowning achievements of the classical geometers. The main obstacle in understanding the classical geometers is the consistent use of imaginary numbers, as these numbers have been dropped from current courses dealing with the geometry of sub-manifolds in scalar product spaces. The current approach is to split this geometry into Riemannian and Lorentz, but this comes at a price: geometric connections between the two are lost and similarities remain only at the analytic level. Also results about *non-totally real* sub-manifolds (sub-manifolds of $\mathbb{C}^n$ which do not lie in totally real subspaces; however they may have real linear element) are lost. Consider for example the statement from Darboux ([14],§ 169): *'A surface of revolution can not only be deformed to a line, but further to an isotropic line'* (thus a 2-dimensional linear element can lose not only one dimension, but further two). Such a statement may seem false if one adheres to the strict current rule that applicability implies same dimension as manifolds, but if one replaces 'surface', 'line' and 'isotropic line' with the 2-dimensional collection of their tangent planes (with the points of tangency highlighted), it begins to make sense at the differential level, at least from a dimensional point of view. Now the condition that a 2-dimensional family of facets (a *facet* is a pair of a point and a plane passing through that point) is the collection of the tangent planes of a sub-manifold (leaf) does not distinguish between the cases when this sub-manifold is 0, 1 or 2-dimensional.

The deformation of a CGC -1 surface to a line explains the generation of the *real pseudo-sphere* (surface of revolution with a tractrix as generating curve) and *Dini surfaces* (helicoids with same type of generating curve) from a line via a B transformation; at the level of the sine-Gordon equation this is equivalent to generating (via a B transformation) 1-soliton solutions of the sine-Gordon equation from the vacuum soliton.

Note the metric-projective nature of the *tractrix*: it is the real curve individuated by the requirement that its real tangents meet a given line at constant distance from the tangency point; it is obtain by slowly pulling a cart along a line if initially the cart axis is perpendicular on the line.

The deformation of a CGC -1 surface to an isotropic line provides a simple point of view on the B transformation, due to Lie and apparent at the level of *confocal* ('with same foci') pseudo-spheres. Lie's ansatz is very simple: because the B transformation is of a general nature (independent of the shape of the seed), it must exist *not* only when the seed is a CGC -1 in $\mathbb{R}^3$, *but* also when the seed coincides with the pseudo-sphere, in which case the leaves degenerate to isotropic (imaginary) rulings on a confocal pseudo-sphere (note that there is no formulation of the Lie ansatz at the level of the sine-Gordon equation, since it deals with imaginary seed). This simplification allowed Bianchi to generalize the B transformation of CGC -1 surfaces to the B transformation of deformations of quadrics and it allows us to simplify the denomination *B transformation of deformations of quadrics* to *B transformation of quadrics*.

Thus the Bianchi-Lie ansatz states that the leaves (B transforms of the seed) are rulings of a ruling family on a confocal quadric when the seed coincides with the considered quadric.

In the famous letter *The Method*, sent to Eratosthenes of the Library-University of Alexandria and lost from the XIII$^{\text{th}}$ century until 1906 (the year of Bianchi's discovery, so it was unknown to Bianchi and Lie), Archimedes states: *'... certain things first became clear to me by a mechanical method, although they had to be proved by geometry afterwards because their investigation by the said method did not furnish an actual proof. But it is of course easier, when we have previously acquired, by the method, some knowledge of the questions, to supply the proof than it is to find it without any previous knowledge'*.

Note that the method at the level of points of facets was known to Bianchi and Lie; however, they had never used the full method, at the level of the planes of the facets too (this is where Archimedes enters into the picture), so Archimedes' part is the essential ingredient for generalizations in other

settings (especially in higher dimensions), since it contains more information. Thus if one assumes Theorem I of Bianchi's theory of deformations of quadrics a-priori to be true and to be the metric-projective generalization of Lie's approach, then by applying the Bianchi-Lie ansatz (the *'mechanical method'* in question) both at the level of points of facets and planes of facets one obtains in a natural and geometric way the necessary algebraic identities needed to prove Theorem I. The fact that Archimedes' opinion is the main ingredient in supplying the proof of Theorem I is confirmed by the fact that Bianchi's original proof is much more complicated, because it was initially discovered first at the analytic level and for the composition of two B transformations, so Bianchi had to retrace the steps back to the basic B transformation at the analytic level and guess the geometric picture; it took him 6 years of work on several particular cases to do so (of course Bianchi simultaneously worked on many other projects during this time) and thus it lacks the *'of course easier'* ingredient.

Of course Bianchi's proof is commendable for the fact that it solved an open problem at that time; just like Archimedes Bianchi showed that it works on several examples (again in each case Bianchi did computations from scratch, as was the standard of proof of the time), so one can safely deduct that it works for all quadrics (deformations of certain particular quadrics were previously found, so Bianchi's method was not seen as necessary in those cases).

However, without the power of a professional with good knowledge of the tools of the trade, the faith in finding a result at the end of long computations and the geometric intuition of Bianchi's, this could have been an open problem even today, as few years after its completion this type of problems fell into forgetfulness for more than 50 years. It is the word superhuman (*'sovrumana'*) that was chosen by Bianchi's contemporaries to describe his efforts and geometric intuition.

We shall call this method the *Archimedes-Bianchi-Lie* (ABL) method.

Note that although Archimedes and Bianchi-Lie have dealt with different problems, their approach was the same: $P \Rightarrow Q$ with P being either *'The area of a segment of parabola is an infinite sum of areas of lines'* or *'A line is a deformation of a quadric'*. Such sentences P were not accepted as true according to the standard of proof of the times, but they had valid relevant consequences Q which elegantly solved problems not solvable with other methods of those times; if only for this fact the standard of proof should change so as to accept these statements as true.

At some point I thought the *'mechanical method'* in question to be the rolling of the seed back on the considered quadric (used by Bianchi in his arguments and clearly a mechanical method), which gives a natural geometric explanation of the necessity of the existence of the *rigid motion provided by the Ivory affinity* (RMPIA) (this is how Bianchi found the *applicability correspondence provided by the Ivory affinity* (ACPIA)) and of the reflection in the tangent bundle of the considered quadric (one can roll a surface on both sides of one of its deformations). These were implicit in Bianchi (122,§ 11) and well known to me, but Archimedes was still not satisfied, because these were not the *'certain things'* which *'first became clear'* and they were geometric arguments which remained to be confirmed not by geometry, but by an analytical calculus (the double reduction ad absurdum, an analytical calculus today by means of the $\delta - \epsilon$ trick, was a geometric argument during Archimedes' times). Thus Archimedes further required that to the rolling of the seed back on the considered quadric one must also apply the ansatz that the seed coincides with the given quadric, in order to get the a-priori nonsensical *'certain things'* which *'first became clear'*, only *'to be proved by geometry afterwards because their investigation by the said method did not furnish an actual proof'*.

The classical geometers frequently used simple geometric arguments to justify statements as Darboux's from above (which provides itself a simple geometric picture of singular solutions of a certain differential system): Darboux in [13],[14] provides such arguments for almost any analytic computation; they are always shorter and more intuitive. However, without Cartan's simplifying notation (which led him to exterior calculus), their computations of moving frames took pages (as required by the standard of proof of the time) and thus the investigation of higher dimensional

problems lagged behind; for example Bianchi takes time in (122,§ 11) to write on a full page all components of the RMPIA, although this explicit formulation is never used (however the components of the rotation look strangely close to certain quadratic quantities with direct usefulness).

Another feature of the standard of proof of the time (namely explicit solutions) had a positive influence. It is one of the possible explanations of the success of the classical geometers in the study of integrable systems, as these are amenable to algebraic formulations of their solutions after a simpler auxiliary differential system associated to a given local solution of the initial integrable system is presumed solved. The algebraic formulation of solutions is possible due to the *Bianchi Permutability Theorem* (BPT), initially developed by Bianchi for the B transformation of CGC -1 surfaces in ([5],Vol **5**,(46)), but actively pursued by him in various settings throughout his career (there are also other results and methods, like the one mentioned in the introduction of Terng-Uhlenbeck [36] and due to Darboux).

With the development of calculus experts realized that most differential systems do not exhibit such behavior and they focused their attention on the qualitative behavior. Integrable systems were revived in the second part of the XX^{th} century, due mostly to the KdV equation and other problems from physics (see Rogers-Schieff [26]).

Because of the BPT Bianchi was able to put a local group action structure on the space of local solutions of an integrable system; thus he 'added' (via a non-linear commutative algebraic procedure, hence the *permutability* denomination) two solutions and obtained a new one. Although the classical geometers did not further investigate this group structure (see Terng-Uhlenbeck [36]), Bianchi was aware of the group structure at least for *n-solitons* of the sine-Gordon equation. The *n-solitons* of an integrable system are obtained by the iterated application n times of the B transformation to the vacuum soliton (practically one applies the BPT $n-1$ times to the vacuum soliton; Bianchi calls iterations of the BPT *moving Möbius configurations*). In the case of the sine-Gordon equation the above group can be identified with its (global) action on the vacuum soliton; its elements admit formulations as rational functions of exponential functions (see ([5],Vol **5**,(46,§ VI))). When the initial value data of the auxiliary differential system are taken into account, the permutability loses its commutativity connotation and the BPT for the B transformation of the pseudo-sphere is explained by two different ways of factoring quadratic terms in the group of rational maps on the unit sphere $\mathbb{S}^2 = \mathbb{C} \cup \{\infty\}$; the factors are Möbius transformations (see Terng-Uhlenbeck [36]).

These notes began with the purpose of understanding the theory of deformations of quadrics; in the process the proofs were reduced and simplified, thus we hope making it easier available to a wider audience. For example the proof of the existence of the B transformation of quadrics and of the ACPIA does not significantly surpass both in length and difficulty the actual statement and its prerequisites. The statements (with no major modifications) are due to Bianchi and their beauty, clarity and simplicity provided the motivation to go through Bianchi's original proofs. There are other classical monographs on the theory of deformations of quadrics, but they are incomplete or the approach is different than Bianchi's: for example Calapso [11] or the last chapter of Eisenhart [20].

While all main ideas are due to the classical geometers Archimedes, Bianchi, Darboux, Lie, etc, at the computational level there is something new: a complete, simplified and unified treatment of all quadrics in $\mathbb{C}^3$. For example Bianchi takes a quadric in $\mathbb{C}^3$, intersects it with various totally real subspaces to get real 2-dimensional (in)definite linear elements and then builds the B transformation in various totally real subspaces separately for each case (providing enough supporting examples according to the required standard of proof of the time). We provide a unified discussion of the B transformation and postpone the necessary splitting into totally real cases, as an application, to the very end (a similar discussion takes place in Terng-Uhlenbeck [36]). Because the initial surface (having $2, 3$ or 4 real dimensions) in the considered quadric or its deformations may be non-totally real, the most general configurations are obtained. In particular if the linear element is real, then the surfaces must have real dimension 2, so we obtain deformations (which may be non-totally real)

of totally real surfaces in quadrics (a surface in a quadric and having real linear element must be totally real).

In what follows we shall deal with quantities which may not be defined everywhere; instead of always pointing out the domain where the computations are true we shall mostly just assume them to be true on their domain of existence, without any further details (we discuss only the local theory). We shall require as many derivatives as needed; in fact since most of the current literature on integrable systems deals with solutions in the distribution sense, all computations can be interpreted in the distribution sense (there may be generalizations even in other settings, like p-adic (non-Archimedean) calculus, but at this point I am not familiar with such proceedings; note however that the correct interpretation of calculus in Leibnitz's notation gives for free the correct discretization). A global theory (if developed) must include singularities (along points or curves) and thus must probably be developed along lines similar to other integrable systems. While it is essentially the trend of the current geometers to prove global results, most such results are about rigidity and for global deformations of complete surfaces such results are few (see Spivak ([29], vol **5**)); thus one can safely assume that in order to have a consistent set of objects to study singularities must be allowed. In most current theories singularities are to be avoided; here they provide degeneracies, due to which certain general differential equations (with no method of integration) degenerate to ones that can be easier integrated. Again one can guess here the effect of Archimedes' method and probably a geometric solution will be either easier to obtain or, in the contrary case, the geometric explanation of the solution obtained by analytic computations will easily surpass and simplify the analytic computations.

As to the n-solitons of integrable systems known so far, they involve difficult formulae, usually involving (hyperbolic) trigonometric, exponential and even elliptic functions; the geometric projective picture should serve as an easier explanation of these strange laws governing these solitons, just like Jacobi's simple geometric interpretation on confocal quadrics of Abel's Theorem (see Darboux ([13],§ 464) and ([14], Note I)).

After the application of the ABL method the computations naturally split into *static* (algebraic) computations on quadrics confocal to the given one and *moving* (differential) computations in its tangent bundle. The flat connection form (a feature common to all integrable systems and always coming in a spectral 1-dimensional family) is provided by *rolling*: two rollable (applicable) surfaces can be rolled (applied) one onto the other, such that at any instant they meet tangentially and with same differential at the tangency point. Conversely, if two surfaces can be rolled one onto the other, then they must be applicable with the applicability correspondence being given by the correspondence between points whose tangent spaces coincide at some instant in the rolling. Note that it is less known today that Levi-Civita introduced the *parallel transport* denomination since it is the transfer by rolling of the usual Euclidean parallel transport in a plane, as the surface is rolled only along the given curve (along which the parallel transport needs to be done) on a fixed plane (thus as a plane rolls along one of its lines on a surface it describes a geodesic on the surface and all geodesics are born this way). The local minimizing property of geodesics is immediate from the geometric definition, since the shortest distance becomes at the infinitesimal level heading in the direction of shortest distance and by rolling this heading sticks (transfers) from the plane to the surface and conversely. An ant on an ellipsoid would do the same as what people did from the beginnings of mankind: it would choose a point far away as reference or it would look at the heavens (tangent space) and it would thus linearize its problem, rolling the heavens with it (changing the point of reference) as it proceeds on its journey.

Currently one would be inclined to take these geometric definitions which inspired analytic consequences as consequences of the analytic consequences (taken currently as definitions, instead of as confirmations), just as a trained mathematician would rather give an $\mathbf{L}^2$ proof of the discrete Cauchy-Schwartz inequality instead of the longer (but at high-school level) proof using induction or just as the Calculus II undergraduate student would give a L'Hospital proof to $\lim_{x\to 0}\frac{\sin(x)}{x}=1$.

Just as higher dimensional calculus is essentially 1-dimensional calculus in all directions plus the Clairaut integrability condition (commuting of mixed derivatives or the d-wedge condition), higher dimensional rolling (including of sub-manifolds in higher dimensional symmetric spaces) is essentially 1-dimensional rolling in all possible directions plus the same integrability condition.

As we have only 1-dimensional time to our imagination, it is somewhat difficult to imagine the 2-dimensional rolling of two applicable surfaces, so we have to imagine 1-dimensional rolling along all possible corresponding curves (which will thus admit an arc-length correspondence; since a correspondence of two surfaces which induces arc-length correspondence of arbitrary corresponding curves must be isometry, existence of rolling implies isometry). Since Leibnitz's notation does not have a memory of dimension (which is dealt with by another condition, namely *the independence condition*), it is the tool best suited to attack the rolling problem and the easy 1-dimensional rolling problem easily introduces the notation needed for higher dimensional rolling problems.

By application of the ABL method certain valid relevant algebraic identities naturally appear (from a geometric point of view) within the static part.

All necessary identities of the moving part boil down to these valid relevant algebraic identities of the static part (other terms of the differential identities dissolve into nothingness because of the flat connection form). A possible justification of this behavior (and thus the miracle of the ABL method) appears when one considers *discrete deformations of quadrics* (DDQ); in 1996 Bobenko and Pinkall have introduced in [6],[7] discrete isothermic and CGC -1 surfaces, although Newton was the first one to observe the discretization of integrable systems and of their solutions when he provided a mostly Euclidean explanation of Kepler's Laws of planetary motion. Discrete surfaces must obey two simple principles: on one hand their determination from the discrete Cauchy data involves only algebraic computations (precisely the algebraic computations required by the BPT; thus the last important step of the theory of deformations of quadrics becomes a basic axiom for DDQ) and on the other hand, just as their name suggests, discrete surfaces better approximate their corresponding surfaces as the discretization becomes finer; therefore they algebraically encode the differential structure of their corresponding surfaces (and thus the most important step of DDQ provides the building blocks of the theory of deformations of quadrics). It is thus no accident that Bianchi had in fact developed discrete versions of various integrable systems and of their solutions when he developed moving Möbius configurations (iterations of the BPT) a hundred years ago, but without being aware of this fact. One can even develop the discrete B transformation for discrete solutions of discrete integrable systems! (the simplest trick to do this is to preserve Leibnitz's notation but to interpret it differently). The infinitesimal becomes the discrete finite by iterating a $\mathbb{Z} \times \mathbb{Z}$ lattice of B transformations on an initial seed and picking a seed point on that seed and its corresponding $\mathbb{Z} \times \mathbb{Z}$ heirs; the finite becomes the infinitesimal by refining the $\mathbb{Z} \times \mathbb{Z}$ lattice of a DDQ (while preserving the subjacent curves of the discrete Cauchy data; since they must become asymptotes they have prescribed torsion by Enneper's). The discrete version of the Clairaut equality $(f_x)_y = (f_y)_x$ for $f = f(x, y)$ implies the BPT and as one refines the $\mathbb{Z} \times \mathbb{Z}$ lattice of B transforms the BPT becomes infinitesimal, so it precisely describes the above equality! Thus the theory of DDQ is the proof *'such as the ancients required'*, provided that all its algebraic computations are put in a synthetic language.

Why one would do such a thing? Basically Archimedes discovered in *The Method* integral calculus for some conics and quadrics of revolution (he discovered almost all important formulae; probably there are other formulae for quadrics which can be discovered with the same method) while Apollonius essentially discovered differential calculus for conics in *Conics* (part of the second half was available in Latin beginning unfortunately only with 1710; among the essentially differential notions introduced by Apollonius was the evolute of a conic). Up to Kepler some more results were added; these allowed him to do the discrete computations for his Laws. Note that basically all methods and results up to Leibnitz and Newton are useful not only from a historical point of view, but also for the fact that these hint an integrable system behavior (take for example Roberval's simple Euclidean arguments in computing tangents by kinematic arguments and the area of the

cycloid; it is probable due to the fact that nature was able to do computations for the cycloid using only simple mechanical methods and simple Euclidean arguments that it chose it as the solution for the *brachistochrone* and *tautochrone* problems (the resolution of the former among others by Newton was the first instance of *Calculus of Variations* or equivalently infinite dimensional calculus) and the masterly unification of the pendulum with the tautochrone by Huygens when he showed that the evolute of a cycloid is a cycloid). Beginning with Leibnitz and Newton calculus was the main theoretical tool, although discrete mathematics remained the main practical tool up to including today (because measurements, observations and computer memories are discrete). Beginning with the second part of the XX^{th} century the use of optimal discrete mathematics for problems with not so obvious solutions began to be again important, as more difficult problems (such as integrable systems) arose; this was the main motivation of Bobenko and Pinkall for their 1996 articles. Thus a synthetic language based on axioms and counting their use is a way to obtain not only the optimal proof, but also to implement the optimal discrete mathematics.

The static part requires basic knowledge of linear algebra and projective geometry; we use for example the Menelaus theorem (which is essentially a co-cycle theorem) to prove the associativity of the group action whose commutative 'addition' law is provided by the BPT; see also Arnold's observation from [2] about the link between the Jacobi identity (infinitesimal associativity, in itself a co-cycle theorem) and the intersection of the heights of a triangle. Note that the Jacobi identity for the R-matrix theory (Lie bracket deformation $[x,y]_R := [Rx,y]+[x,Ry]$ with R linear of the Lie bracket $[,]$; see Burstall, Ferus, Pedit and Pinkall ([10],§ 3)) may be the possible analogy of the Menelaus theorem, as I have noticed the R-matrix behavior (and consequently I was able to simplify the notation of the algebraic computations of the BPT) before seeing the Menelaus theorem. What are the chances of reading a book of history of mathematics, see a theorem, remember to have seen it in grade school and immediately realize that it is what one needs to answer the current questions?

In the first issue of the *Bulletin* on 2006 I read something about the Arnold-Jordan canonical form of matrix and I wanted to see if Arnold had already done what I call the symmetric Jordan canonical form; this is how I discovered [2].

So the R-matrix theory is probably the necessary tool in higher dimensions and the Jacobi identity of $[,]_R$ (which imposes certain conditions on R) is essentially a generalization of the Menelaus theorem in other settings.

For the moving part we shall only use rudiments of the absolute (Ricci) calculus of tensors, working mostly with $(0,1)$ and $(0,2)$ tensors. Since we would like d^2 to mean tensorial second derivative, we shall use the notation $d\wedge$ (d-wedge) for alternative (exterior) derivative. Take for example a plane $V\subset\mathbb{C}^3$ with normal $0\neq N\in\mathbb{C}^3,\ N\perp V$; if $|N|^2\neq 0$, then one can normalize N to $|N|=1$; otherwise N is an isotropic vector: $|N|=0$, so $N\in V$ and $<,>$ restricted to V is degenerate; thus the second fundamental form of most surfaces $x\subset\mathbb{C}^3$ is defined almost everywhere as N^Td^2x (Cartan never uses the notation d with the meaning $d^2=0$, as he probably wanted to preserve Leibnitz's notation: $\frac{d^2y}{dx^2}=\frac{d}{dx}(\frac{dy}{dx})$; see ([12], Ch III,§ 26)).

Most classical integrable systems were discovered while studying quadrics; thus they naturally admit, at least in particular cases, (geometric) links to quadrics. For example Darboux was studying in [16]-[18] rolling *congruences* generated by isotropic rulings of quadrics when he found the D transformation for special isothermic surfaces (a *congruence* in $\mathbb{C}^3$ is a 2-dimensional family of objects, presumed lines unless otherwise stated, but they can also be arbitrary curves or even surfaces; following Plücker a 3-dimensional family of objects in $\mathbb{C}^3$ is called *complex*). As the quadric rolls on its applicable surface in a 2-dimensional fashion (assume that the applicable surface is not ruled, in which case the rulings correspond under the applicability to a ruling family of the quadric and the rolling takes place in a 1-dimensional fashion), one if its isotropic rulings describes a 2-dimensional family of isotropic lines; hence the denomination 'rolling congruence'. The natural question thus arose if this transformation exists for general isothermic surfaces and Darboux provided an affirmative answer. The auxiliary differential system corresponding to the D transformation for isothermic surfaces admits a quadratic prime integral, which may explain the

link between special isothermic surfaces and quadrics; for the link between isothermic surfaces and quadrics we probably need quadrics in higher dimensional spaces.

Based on the classical and current literature (see for example Moser [24] or Rogers-Schieff [26]), we believe that any integrable system is bound to be linked at least in particular cases to quadrics, even in infinite dimensional spaces, where a spectral theory is well developed but without the corresponding geometrization; in fact confocal quadrics as particular confocal cyclides and *Dupin cyclides* (surfaces with circular lines of curvature) clearly appear in Darboux [14] as important examples and the notion of isoparametric sub-manifolds in finite dimensional spaces has already been generalized to isoparametric sub-manifolds in Hilbert spaces by Thorbergsson and collaborators. Note also that cyclides in pentaspherical coordinates have elliptic coordinates just like confocal quadrics (see Darboux ([13],§ 437 and Livre IV, Ch XII)).

We also believe that the above formulation (split the computations into static and moving by means of the ABL method) is generous enough to be amenable to generalizations in other settings; in particular we believe that the B transformation of deformations of space forms in space forms and with flat normal bundle (Tenenblat, Terng, Uhlenbeck and collaborators) can be generalized to the B transformation of higher dimensional quadrics in symmetric spaces and with flat normal bundle.

1.3. **Short historical perspective on quadrics.**
The interested reader is referred for more details to Boyer [8] or the website [39] (our main sources for this subsection up to the beginning of the XIX$^{\text{th}}$ century), to G. Scorza's and G. Fubini's eulogies in part 1 of Volume 1 of Bianchi [5] and to R. Calapso's introduction in part 1 of Volume 4 of Bianchi [5] (our main sources for this subsection in what concerns the history up to 1912 of the theory of deformations of quadrics).

Further this perspective (and its references) is definitely biased and incomplete, it being designed to suit a restricted point of view; besides that it has a personal touch (*'You want to make an omelet, you got to break some eggs'*). Problems of the XIX$^{\text{th}}$ century such as Monge's sphere (for any quadric there is a sphere and a moving orthonormal frame centered on that sphere such that the hyperplanes of the orthonormal frame are tangent to the given quadric) and Chasles's number of 3264 conics tangent to five given conics will not be discussed in detail here.

The primary organization of this subsection is the chronological one; the secondary one is focused on the subject at hand.

Quadrics are the simplest non-linear geometrical objects and thus they arose early on the interest of geometers: the ancient Greek geometry culminated with the work of Apollonius, Archimedes, Euclid, Pappus and others in particular on the geometry of conics.

Archimedes is credited to have been close to a theory of integral calculus: to find areas and volumes he used at the infinitesimal level an ingenious balancing argument and simple Euclidean arguments (the famous letter *The Method*, sent to Eratosthenes of the Library-University of Alexandria and lost from the XIII$^{\text{th}}$ century until 1906); to confirm the validity of the formulae thus obtained he used a double reductio ad absurdum, as was the standard of proof at the time. The discovery of *The Method* is important because it changed Archimedes' reputation from being unclear on the details and motivations of the methods employed in proving his results (presumably with the purpose of enhancing his own reputation) to being very generous in *disclosing the full method of investigation* to Eratosthenes, *'a capable scholar and a prominent teacher of philosophy'*, so with the stated purpose that the later should further investigate it and teach the curious trick to his students.

To find the area (volume) of an object Archimedes cut it into parallel lower dimensional slices (segments for area and plane figures for volume), then he placed all slices with their center of mass at one end of a balance and a simpler object at the other end; if equilibrium is obtained the area (volume) of the initial object can be found. The equilibrium is checked by comparing slices of the original object with corresponding slices of the simpler object and *not* only simple Euclidean geometric arguments are needed, *but* the choice of *the correct balance with fixed fulcrum* and of *the*

simpler Euclidean object is revealed by the simple geometric properties of the slices of the initial object. So basically *The Method* of Archimedes takes into account the *discrete* version of computing the *mass* and *center of mass* of an object from the knowledge of the *masses* and *centers of mass* of its component pieces and assumes it also to apply as a limiting case to the *infinite* (non-discrete) version of the same statement; the use of simple Euclidean statements about conics available at the time was realized with a balance and thus interesting results about conics and quadrics of revolution were added to the bag. Nowadays this would be proof enough (the slice theorem or Cavalieri's Principle for parallel slices), but Archimedes was unsure of the power of his method: he would set out to prove its validity by thickening the slices to regain the original measure (so he would go from the singular boundary of the space of solutions to his discrete principles back into the meat of the original space of solutions, where one can wiggle and still remain in the meat; this is what actually happens with the theory of deformations of quadrics in this notes and possibly with *all* integrable systems: we are unsure of the power of Archimedes' method and provide further explanation which may be considered unnecessary in the future) and a double reductio ad absurdum, as was the standard of proof at the time. Note that all through his method pervasive is the use of the balance with fixed fulcrum and left end where facets are *transferred and stuck with their center with* ∞ *multiplicity* (the variable right end of the balance is dictated by some of the variable mass of the object at the fixed left end, as it transfers and sticks to the apparently wrong unit of measure of length); since the balance works with any infinitesimal configuration, it will work with the integral version. One can look at the good metric properties of the Ivory affinity as being the required prerequisites of simple Euclidean relations by means of which one puts the quadric at the right end of the balance and rulings on a confocal quadric as leaves at the left one. Now the existence of seed provides a way to deform the quadric, that is a way to rearrange the object at the right end of the balance: the balance remains pervasive through this process and allows the rearrangement of the facets at the other end in a manner consistent with the initial arrangement (that is admitting leaves) so the balance wins and its principles (clearly valid at the infinitesimal level by means of the simple Euclidean properties) remain valid after integration. The integration process (proven by Archimedes to be correct by means of a double reduction ad absurdum) is realized by means of flat connection form: facets fit nicely at the infinitesimal level on the right; as a consequence facets fit nicely on the left. The discrete version of the theory of deformations of quadrics can be actually put by means of a $\delta - \epsilon$ argument (error estimate) to the form of a double reduction ad absurdum.

With this method Archimedes managed to compute the area and center of mass of the region bounded by a chord on a parabola, volumes of segments of quadrics of revolution, centers of mass of hemispheres and of the paraboloid of revolution. He also used a geometric argument reminiscent of what we call today 'trapezoid rule' to find the area bounded by a chord on a parabola; the argument does not work for ellipses or hyperbolas, as they cannot be realized as graphs of quadratic functions; however he was able to find the area of the ellipse.

Apollonius is mostly known for his *Conics*, which began as a project of collecting the results about conics from Euclid's *Elements* and other sources of the time, but stated in a more general setting and completed with corresponding proofs; in the process it became a subject of interest on its own and completely replaced all other sources, remaining the standard for almost two millenniums, just as Euclid's *Elements*. While it seems that Apollonius and his contemporaries were familiar with foci of conics and some of their properties, these were considered and referred to only indirectly at that time. It is Apollonius who first introduced the *hyperbola* (something is in excess), *parabola* (something is precisely) and *ellipse* (something is missing) denominations took over since then in many other domains (such as literature).

According to Boyer ([8], IX,§ 5): *'If survival is a measure of quality, the Elements of Euclid and the Conics of Appolonius were clearly the best works in their fields'.* It is chiefly because of the *Conics* that Apollonius was known as *'The Great Geometer'* of the antiquity. Understanding this treatise was regarded as the ultimate test an accomplished geometer should pass; this may be the possible explanation of the fact that the second half of the *Conics* has survived only partially and in

arabic (Jafar Muhammad ibn Musa ibn Shakir's version) until 1710, when, belated and apparently useful only from a historical point of view, Halley's first Latin translation appeared. Note however that this translation could have been done almost a century earlier, as the manuscript was available in one of Europe's libraries, but administrative reasons have prevented it.

According to a *'Nova'* program on Archimedes' palimpsest calculus could have probably been discovered at least one century earlier with *The Mehod* of Archimedes available at the time; consequently the small step for *'a man'* in the XX$^{\text{th}}$ century could have been on Mars (Florin pointed out also the other side of the coin, namely humankind could have self-destructed in the XIX$^{\text{th}}$ century).

Note also that the complete version of Apollonius's *Conics* could have helped in early XVII$^{\text{th}}$ century: among other essentially differential notions first introduced by him using only synthetic computations was the *evolute* of a conic. When a line rolls on the planar *evolute* a point of that line will describe an *involute*; thus the orthogonal family of a family of plane lines are involutes; in our situation the given conic is among the involutes. This picture admits generalization to curves in space by bending the plane of the curve with a cut made along the curve (although this is still not the correct picture, as one needs two copies of the original plane; one can make a contraption with few tangent lines attached to a thin metal sheet (a one sided tubular neighborhood of a plane curve)): the normals to a involute surface envelope focal surfaces (evolutes), the vector fields induced by normals on an evolute surface admit integral curves which are geodesics on the evolute surface; conversely a 1-dimensional family of geodesics on a surface makes that surface into an evolute of another 1-dimensional family of involute surfaces and each geodesic and its involutes (curves) is in the situation of a bent plane picture. Another possible generalization would be from a 1-dimensional family of plane lines to an $(n-1)$-dimensional family of lines in $\mathbb{C}^n$; it still admits $\leq n-1$ evolutes (focal hyper-surfaces or caustics), but it would not admit in general involutes, as the condition that such an $(n-1)$-dimensional family of lines are normals to a hyper-surface is special.

Pappus introduced the *focus-directrix* property of conics (in the plane geometry all conics are the locus of points such that the ratio of the distances to a given point and a given line is constant; the given point becomes focus and the given line directrix), but he is mostly known for a mechanical method of finding volumes of solids of revolution, which requires only knowledge of the area and center of mass of the generating plane figure; as a limiting case one can obtain the area of a surface of revolution from the length and center of mass of the generating plane curve. Replacing the generating plane figure with its center of mass and having as mass the area of the plane figure, the moment about the axis of revolution does not change; as a limiting case one can obtain the area of a surface of revolution from the length and center of mass of the generating plane curve (note again that the part of Archimede's method concerning the mass and center of mass provides the correct explanation if the slices are the positions of the original plane figure (curve) as it rotates as instructed).

At that time they endured the criticism of their opponents as to the usefulness of their results (see page xi of *Preface* to T. L. Heath [21] and the frontispiece); according to Apollonius *'They are worthy of acceptance for the sake of the demonstrations themselves, in the same way as we accept many other things in mathematics for this and for no other reason'.*

Up to the XVII$^{\text{th}}$ century the works of the wise ancients were required reading and their open questions constituted the main driving force in the development of mathematics; according to Leibnitz *'He who understands Archimedes and Apollonius will admire less the achievements of the foremost men of later times'* *.

* This quote (as most others of this subsection) was introduced in the early 2006 version of the notes, but at that time I thought it to seriously apply only up to the XVIII$^{\text{th}}$ century, since Leibnitz was not aware of Archimedes' simple point of view from *The Method* and also I was not aware of the range of applications of Archimedes' ideas. *'That's a dam' shame!'*

It is in trying to come closer to God that people chose Euclid as a model; thus Euclid's *Elements* were required reading for any important man regardless of his preoccupations and it is no accident that in *The Declaration of Independence* of 1776 one can easily see the simplicity of the proof beginning with simple axioms and easily drawing the corresponding conclusions, just as the balance is a universal principle (any action will have as effect a corresponding opposite reaction equal in size in a certain measurement).

Tartaglia (The Stammerer; he had his face sliced open by The French in his childhood) thought initially that the trajectory of a cannon ball is a broken segment; later on he changed his mind to a parabola. One could easily assume that Tartaglia was a naïve mathematician, only that de discovered the formula for the roots of a cubic equation, a thing above me without the help of Google. From the beginnings of mankind and probably up to its ends mathematicians are seeking sponsors and most sponsors are interested in measuring for taxing purposes (this is how geometry appeared in ancient Egypt and Gauß initially set out to make a geodetic survey of Hanover before he had his enlightening remarkable discussion; it is probably because of always noticing the same simplifications when compiling discrete measurements of angles and distances that he got the idea) and warfare so as to enhance and preserve the right of taxation: the first successful algebra book in Middle Aged Europe was successful due to its appendix in which it presented the method of *cooking the books*, so the first million is questionable not only for successful businessmen.

Using Tycho Brahe's detailed astronomical observations, Kepler found an important application:

I The orbit of a planet is an ellipse with the Sun situated in one of its foci.

II A line joining the planet and the Sun sweeps out equal areas in equal times as the planet describes its orbit.

III For any two planets the ratio of the squares of their periods is the same as the ratio of the cubes of the mean radii of their orbits.

Newton was able to explain this by the theory of universal gravitational attraction, thus discovering probably one of the first integrable system, for which Kepler's laws are conserved quantities. One could actually recognize the first integrable systems in some of the results of the wise ancients in the form of laws of a general nature. For example Apollonius's result that the tangent to a hyperbola cuts the asymptotes at points equidistant from the point of tangency is clearly a metric-projective result which easily individuates hyperbolas as the only real curves with this property; it remains valid if one replaces the asymptotes with another hyperbola having the same asymptotes (some such hyperbolas are homothetic to the given one; the whole family is the pencil of conics generated by the asymptotes and the line at ∞ counted twice; thus all such conics are tangent to the asymptotes at their corresponding point at ∞) and thus any line cuts a hyperbola and its asymptotes equidistantly (another result of Apollonius, which thus implies the previous statement). This can be easily generalized (by taking the slopes of the asymptotes to be conjugate purely imaginary) to homothetic ellipses (which have imaginary asymptotes); moreover if one considers the y coordinate purely imaginary such that the asymptotes have parametric equation $[t \ \ \pm it]^T \in \mathbb{R} \times (i\mathbb{R})$, then the above mentioned equidistant segments also preserve same value if the subjacent lines remain tangent to the same pseudo-circle (which thus becomes (isotropic) pseudo-tractrix); the equidistance gives the correct explanation of the a-priori counterintuitive fact that a plane with positive definite induced linear element in $\mathbb{R}^2 \times (i\mathbb{R})$ intersects any light (null) cone along a circle; one can apply this plane to a plane in $\mathbb{R}^3$ and actually see the circle. Darboux especially has the frequent custom of intersecting the tangent planes of a (presumably real) surface with a sphere of zero radius and obtaining circles.

Note however that Apollonius's result is not fully metric-projective, as it is not true for the pencil of homothetic parabolas (with equation $\lambda y^2 + xz = 0$; thus it is the pencil generated by the conic $xz = 0$ and the conic $y^2 = 0$); this can be easily seen by considering lines passing through the origin. Moreover the orthogonal curves of such pencils of conics are (excepting the homothetic parabolas) seldom pencils of homothetic conics. Thus one can safely deduct that an interesting metric-projective notion valid for all conics must involve more than a pencil (every pencil of conics

contains two degenerate ones) and the fact that the singular conic being a double line must be the line at ∞ for the trick to work will become clear later, since the information about the Euclidean scalar product is projectively encoded by a quadric on this line (thus two points).

Note that the pseudo-tractrix $\Psi = \Psi(s) := [\sinh(s)\ i\cosh(s)]^T$ and the tractrix $T = T(s) := [e^{-s}\ \int_0^s \sqrt{1-e^{-2u}}du]^T$ have the same metric definition, in the sense that one can roll the pseudo-tractrix on the tractrix such that the isotropic line $\Psi + \dot{\Psi}$ is captured and transported by the rolling, thus becoming the line $T + \dot{T}$. The rolling must be a curve $(R,t) = (R(s),t(s)) \subset \mathbf{O}_2(\mathbb{C}) \ltimes \mathbb{C}^2$ such that $(R,t)(\Psi, d\Psi) := (R\Psi + t, Rd\psi) = (T, dT)$. Now the rotation R is determined up to a reflection in the tangent bundle (the symmetry of the normal bundle) by $Rd\Psi = dT$; the translation $t := T - R\Psi$ satisfies $dt = -dR\Psi$. It thus becomes straightforward that $R(\Psi + \lambda\dot{\Psi}) + t = T + \lambda\dot{T},\ \forall\lambda = \lambda(s) \subset \mathbb{C}$.

As an indication of the standard of proof of the time (or according to the historian who must cover all possibilities, in order to avoid controversy), Newton kept the use of calculus at a minimum, using, in the spirit of Euclid, mostly geometric arguments. A big part of his *'Principia...'* deals with properties of conics, his chosen argument for the Pappus four-line locus problem being *'not an analytical calculus but a geometrical composition, such as the ancients required'*. Again the historian says that this may be an indirect attack on Descartes and his methods, who had previously solved this problem of Pappus, in the process revolutionizing mathematics as we know it today by the introduction of coordinates, although again the historian says that Apollonius was the first one to have introduced coordinates and formulae for changes of coordinates (invariant computations) at a synthetic level.

According to Newton *'If I have seen farther than Descartes, it is because I have stood on the shoulders of giants'*, as he probably referred to his predecessors (up to the first half of the $\mathrm{XX}^{\mathrm{th}}$ century one can find in the classical geometers direct references and pride in the lineage up to including the wise ancients).

Currently calculus as developed by Newton in *'The Method of Fluxions'* is mostly replaced by Leibnitz's elegant notation, so Newton remained with the reputation among physicists and Leibnitz among mathematicians (because in physics one needs computations in functions depending on the time variable; a mathematician may find oneself in trouble when one needs the introduction of the time variable, since experiment and observation come foremost and analytic confirmations later). The split that occurred when Newton, following *' the bitter quarrel between adherents of the two men'* decided to delete the reference to Leibnitz in the third edition of his *'Principia...'* was to remain on and off for centuries and even present today (see Arnold [2]).

The fact that Newton's methods were not the correct ones (no interesting mathematics happened in England until the $\mathrm{XIX}^{\mathrm{th}}$ century) is interpreted by the historian as an indication of the superiority of Leibnitz's notation (which was widely used in Europe).

But if it is the duty of the historian to appreciate the value of a book by its influence, it is the duty of the current mathematician to read the books of historical importance pertinent to his restricted field of expertise and to appreciate not only the value of the set of notations used, but also the value of the ideas and motivations behind those notations. It is not excluded that behind Newton's ideas one can see a symplectic structure and his struggle to define it, as such structures naturally appear from physics and Hamilton (the same with the hamiltonian and the four quaternions $1, i, j, k$, named after the four groups of four Roman soldiers each who guarded at some point St. Peter) was the first important scientist in England after Newton to produce significant research. The precursor of Cartan's exterior calculus and higher dimensional moving frames technique is clearly present in the classical French geometers.

The standard of proof for a differential problem decided by Newton was to remain for two centuries: consistent use of Euclidean geometrical arguments and finding solutions in closed form (that is involving mostly algebraic and differential computations; quadratures are used only if necessary). By referring to simple Euclidean geometric arguments not only one can easily see what is the course of investigation that must be followed and complicated analytic computations are

easily bypassed, but criticism as to the soundness of the argument is avoided (for example Kepler found the recent discovery of logarithms by Napier a great help for his complicated computations; when advised not to trust logarithms as being a contraption with possible hidden contradictions he set out to give a Euclidean proof of the soundness of logarithms and their properties).

By the mid-XIX$^{\text{th}}$ century a crisis has reached Euclid: the discovery of non-Euclidean geometries (although this particular rupture was produced by renouncing a single axiom). It seems that Gauß had previously discovered such contraptions, but did not dare stir controversy (as it is the case with most others contraptions of the XIX$^{\text{th}}$ century). This particular controversy was somehow closed when Beltrami showed that the hyperbolic 2-dimensional geometry can be realized as the real pseudo-sphere, so ultimately the 3-dimensional Euclidean space contained the crisis in 2-dimensions (one can easily generalize this to a piece of $\mathbf{H}^n(\mathbb{R})$ being realized as a 'real pseudo-sphere' in $\mathbb{R}^{2n-1}$; see Tenenblat-Terng [32]; it is Bianchi who showed first that space forms of same curvature and dimension are isometric).

However more abstract linear elements appeared: beginning with an object in a surrounding Euclidean space, one records the induced linear element, one deletes the surrounding ambient space and then one studies only the properties of the linear elements with properties as those obtained as above (Riemann). But one can safely assume the existence of the surrounding Euclidean space in order to get a good picture, as the only serious obstruction to the existence of the Euclidean surrounding space for these abstract linear elements was the difficulty of the corresponding differential equations; thus one can work at the level of the linear element with *parameterizations* instead of with *maps*.

This is a major rupture with Euclid's ideas, as Euclid was confined only to the infinitesimal world; however even in this case general laws and structures were inspired and best explained by a simple Euclidean picture at the infinitesimal level. Consequently most of them began the long difficult journey back to the finite world.

Although by mid-XX$^{\text{th}}$ century this crisis of abstract notions gave signs of being resolved, the current trends in teaching geometry have yet to recognize this fact (see Arnold [2]).

One cannot understand the classical geometers of the XIX$^{\text{th}}$-XX$^{\text{th}}$ centuries if one pays attention only to the two fundamental forms (the linear element and the shape) and to the normal connection (the geometry of the infinitesimally close surrounding space), since, in the spirit of Euclid, most geometric constructions of the classical geometers take place in a surrounding space and mostly at a respectable finite distance from the considered surface and the analytic version of such computations is meaningless at best at the first look (for this reason Bianchi had to struggle for 6 years to find the theory of deformations of quadrics, since equations initially discovered for particular cases behave better at the analytic level, so one naturally assumes that the same thing will happen for arbitrary quadrics: he had to guess the geometric picture at the analytic level).

Of course one can probably translate the B transformation to a language of abstract bundles over abstract surfaces, using the ∇ notation for the Levi-Civita connection, the symmetric shape operator instead of the second fundamental form, jets, etc, but that will not make it more illuminating, as even for representing such things we need to go back to the 3-dimensional space and draw a picture; moreover a quadric is naturally given by a quadratic equation in a surrounding space and its confocal family is not naturally defined in a bundle.

Ever since Cartan has shown that the geometry of sub-manifolds in symmetric spaces can be attacked only with Leibnitz's notation and with moving frames, I did not see any other simpler point of view. One needs not the notation x_* for x an immersion of a manifold in a symmetric space, when one can use instead dx or $x^{-1}dx$, etc for x being the actual sub-manifold in the symmetric space (or one of its representatives points and thus allowed to move only on the sub-manifold). Consequently we need local parametrizations of x and not local maps on x.

Note however that for arbitrary 2-dimensional quadrics we cannot use orthonormal frames, as other frames (rulings) of a projective nature are naturally chosen by the static picture; probably

for higher dimensional problems we need frames which are projective on quadrics and orthonormal in the normal bundle.

In fact one can find in the classical geometers extensive elementary treatments of various ideas and deep results a-priori known to us to be of a recent nature, like comparison geometry, geodesic flow and its rigidity (which can be extended so as to include a-priori non-arc-length correspondence of geodesics or just at the infinitesimal level of the flow), minimal surfaces, Dupin's cyclides as precursors of isoparametric sub-manifolds, etc: current monographs should at least mention these facts if they choose not to cover them. (Un)knowingly ignoring already known and established important facts will have as only effect the perpetual rediscovery of same and ubiquitous simple principles, disguised in an apparent aura of genius, modernity, rigorousness and firm footing. The very elementary, intuitive, unsound character and a-priori unwarranted generality of ideas and results of the classical geometers constitutes in fact the essence of the broad application of their ideas and results: decades before Einstein's theory of (special) relativity the classical geometers investigated and found interesting applications and properties of imaginaries in geometry, only to be apologetic of the fact that they did not see physical applications other than to apply various tricks and restrict the whole discussion back to the real domain. Darboux especially has the habit of applying a homothety when discussing non-zero CGC surfaces and reducing the argument to CGC -1 surfaces. Minding essentially did the same for geodesic triangles on spheres and found hyperbolic trigonometric formulae for geodesic triangles on pseudo-spheres (thus non-compact continuous groups of symmetries and corresponding symmetric spaces as counterparts of the compact ones); see (Darboux ([13],§ 666)). Current curvature rigidity results have in general analytic Riemanian manifolds with big continuous groups of symmetries and other important extra-structures as end objects of investigation, but the techniques are clearly split between positive and negative curvature, so one can apply complexification and possibly restrict to interesting totally real cases or methods of investigation.

Einstein's theory of general relativity as it appears in Levi-Civita [22] is still very down to earth (he never applied the abstractions, as the ideas of *'Gauß, Riemann, Christoffel and Ricci'* used by him were still very down to earth). It is in trying to explain Einstein's ideas that Levi-Civita came up with his connection and parallel transport, as the surrounding Euclidean space was non-existent, so Levi-Civita stuck the Euclidean picture to a sub-manifold and transferred it to the abstract space; another motivation of Levi-Civita's was mechanical problems. All interesting problems have elementary formulations (at least in particular cases, such as the special theory of relativity being the infinitesimal flat picture of the general theory of relativity; when one tries to integrate this infinitesimal picture one *naturally* gets the general theory) and these elementary formulations should be presented first within their simpler frame as a motivation for further generalizations; further each abstract notion should come with an interesting generalization for which the previous simpler apparatus was not generous enough to resolve (like the Lebesque integral being the natural generalization of the Riemann integral and built around the necessary natural axiom that limits commute with integrals). Most of the time the correct tools needed to attack the more general problem will turn out to be just the correct generalizations of the elementary tools used to attack the elementary problem; otherwise probably the general problem is a non-interesting one. There are rare exceptions and infinite dimensional spaces is one of them; however even here there are few basic results (in Banach, Hilbert and locally convex spaces) and most interesting applications boil down to these basic results and a-priori estimates; the interesting ones which do not yet boil down to these are still currently shown to actually do; thus methods to be taught to students should mainly focus on juicy consistent applications of these principles of a general nature instead of repeating dry abstract foundational issues for most part of the course and to present few elementary cases (usually involving the Euclidean space, (pseudo)-spheres and complex, quaternionic and Cayley projective (hyperbolic) spaces and amenable to elementary resolutions, like Desargues's theorem proving that there is no $\mathbf{P}^n(\mathbb{C}a)$ for $n > 2$ or Pappus's theorem not holding in $\mathbf{P}^n(\mathbb{H})$) as examples of culminations of the abstract theories. I should mention that I was not the best (at mathematics)

of my generation in my county; one colleague and friend went to the International Mathematics Olympiad (he is from the same city as the only other ialomiţean to have ever gone to an IMO), but he declared to me one year in college: *'Ion, I do not see the ideas. Where are the ideas?'*. I did not know at the time what he was talking about, as I saw simple multi-linear and *funktorial* operations and I was hoping to later see their interesting applications, but if he would have asked me a simple mechanical problem like the investigation of a water bag as it was thrown from the dorm window at the time of the questioning, then I would have understood him, as I would not have been able to apply Euler's equations (equivalent to the GW or their integrability condition; I have a vague memory at this point about this trick which can be generalized to infinite dimensional Lie groups), not being aware of the existence of such equations. Of course the natural question arises as to why one would study such an earthly mundane motion instead of abstract purified heavenly mathematics and the answer lies in the merry-go-around experiment of Einstein's: the change of frames must be valid also in the theory of relativity because nature does not favor any frames above others and therein lies the trick.

Not to mention again Archimedes' method of integration, for whose application one must completely renounce the current rigorous definition of isometry, a fact even the classical geometers of the late $\mathrm{XIX}^{\mathrm{th}}$-early $\mathrm{XX}^{\mathrm{th}}$ century were already uneasy to do and as a consequence they renounced it with too much restraint (they have the excuse of not having access to Archimedes' method being taught in high-school, as Archimedes is one of the handful whose words have such power so at to overturn deep rooted trends (even the mighty Earth itself with the proper choice of balance) and up to date Archimedes is the only one whose words rewrote and may still rewrite books of history of mathematics if the buzz around the palimpsest turns out to be true). An interpretation is not rigorous but partial if it removes important applications; it is not a simplification if it removes the very foundation of the original elementary intuitive concept (most of the times and with use of proper toys and experiments available at the level of grade and high-school) and replaces it with a concept only some college graduates would understand. Just as Archimedes dealt with a raw unsound principle at his time, one must probably carefully tread through all the current sound foundations of geometry and allow larger non-rigorous archaic unsound interpretations, at least initially at the level of elementary intuitive a-priori geometric arguments (again just like Archimedes) and in the direct interpretation as intended by their true discoverers before partial current interpretations (if any) were found, people were lost in translation and consequently the heritage of our ancestors was lost. The best free minds of the past had potential no different than that of the best free minds of our times and they delivered magnificently on the problems at hand. Chern was the best geometer of the second half of the $\mathrm{XX}^{\mathrm{th}}$ century (in fact it is Chern who raised Geometry from almost complete oblivion to its current status), but that happened because according to his own account he stood on the shoulders of the giants among others Blaschke and Cartan and one can find in the classical geometers proper justification for each new abstract notion and distillation by means of juicy applications (thus one must first at least cursory read Blaschke and Cartan and only later Chern; it is locally more expensive but on the long term is the optimal path; there is no way to fully understand and appreciate the value of Chern's contributions otherwise). Just as a human being passes thorough different stages in one's mother womb (egg-invertebrate-fish-amphibian-mammal-full developed human), a student in mathematics should be first and foremost rather trained in the history of mathematics as it was naturally developed and first and foremost taught rather each important change and why it came after difficult *'lupte seculare'* (centuries' worth struggles) and the motivations behind the scene: a full human being cannot be a full human being before spending some time as a fish, an amphibian or a mammal; otherwise it will lack and thus will have trouble seeing when it is presented in front of one's eyes respectively a spine, the four limbs or the warm blood; if chance has it that the student actually recognizes the simpler stage of the missed development, then the student has no choice but to fall back at that stage (memories of one's childhood) and redo everything from scratch. The Romanians have a strange sense of humor and consequently corresponding jokes; I can only think of one at this juncture: *A dentist*

from abroad pulled a tooth of a Romanian travelling abroad during Ceauşescu's regime through his ... (aka not through his mouth) because the Romanian was supposed to keep his mouth shut while abroad. I would say that Archimedes' balance should be a tool to be kept handy by all dentists, as it can move the mighty Earth itself with proper choice of fulcrum.

The fact that quadrics behave well *with respect to* (wrt) quadratic laws and conserved quantities (like the energy or the linear element for applicable surfaces) seems to be a tautology to the naïve, but to the not so naïve the difference is clear: 'quadratic' in quadrics is static, while 'quadratic' in quadratic laws and conserved quantities is moving (differential). But what is differential? According to Leibnitz and Newton it is Euclid's world (infinitesimal or finite) in motion. While at the infinitesimal level all things are simple and Euclidean arguments suffice, in the finite world Euclidean arguments apply to much less, as in the process of zooming out from the infinitesimal world to the finite world simple objects accumulate to objects out of the range of Euclidean geometry. Miraculously, for quadrics certain configurations in the Euclidean infinitesimal world (static) survive the trip to the finite world (moving) and the naïve turns out to be right, in the process reaping another success: the very same survival of the configuration from the static picture gives for free a quadratic law or conserved quantity. Conversely, a quadratic law or conserved quantity chooses the configuration which will survive the trip: since the law or conserved quantity is of a general nature, it must be true for any moving configuration at all instances (infinitesimals) in the moving picture and thus its very nature lies in the static picture. Once a configuration is chosen by a quadratic law or conserved quantity, it is very likely that it will be chosen by many others, since nature favors simplicity and thrift even more than man. These are some of the reasons for which the wise ancients were able to find so many interesting properties of conics without use of calculus and for which these properties can be found in Kepler's Laws and Newton's *'Principia...'*. But why 'quadrics' and what are 'certain configurations'? Consider an object with as many symmetries as possible: the more symmetries, the more chances are that that object will satisfy more laws, not only in the static Euclidean picture, but also in the moving picture. The first natural choice is a linear subspace and this leads to differentials, as they live in tangent spaces (the only objects common to both the infinitesimal and finite world and by means of which one jumps from one to the other). The next choice must be non-linear and simple: thus it is the sphere. In both cases Euclid provided a consistent body of results in the static picture and low dimensions. Now consider the moving picture: the tangency configuration enters into our consideration with differentials, as they live in tangent subspaces. Since 'tangency' is a projective notion, one can replace the sphere with a projective equivalent: an a-priori general quadric (it is known since Apollonius that the tangents to conics can be constructed with ruler and compass; Pascal showed that the compass can be dropped and according to the historian there are no other curves with this property). But keep in mind quadratic laws and conserved quantities: these are metric properties, so we need metric-projective tangency configurations of quadrics. Thus appears the family of quadrics confocal to a given one: its definition is mixed metric-projective and depends on a spectral parameter z.

Consider a simple 1-dimensional example: the rolling of an ellipse on a plane curve; this will provide a simplified analogy to the process of generalization of the B transformation from the pseudo-sphere to quadrics. At any instant the ellipse and the curve meet tangentially and with same differential at the tangency point. Since any two curves can be rolled one onto the other (there are no integrability restrictions on the rotation of the rolling), the problem is of a much too general nature to be useful, so we have to make an ansatz. If the ellipse is a circle, the natural ansatz is that the center should preserve constant 0 elevation, in which case the curve of rolling must be a horizontal line. With this in mind we go back to the case of the ellipse, when the center of the circle is replaced not with the center of the ellipse, but with one of the two foci, as they are the heirs of the metric-projective properties of the center of the circle.

Consider $e_1 := [1\ 0]^T$, $e_2 := [0\ 1]^T$, the focus $F := \sqrt{b-a}e_2$ of the ellipse $E : \frac{(e_1^T E)^2}{a-z} + \frac{(e_2^T E)^2}{b-z} = 1$, the point $E(s) := \sqrt{a-z}\cos\frac{s}{\sqrt{a-z}}e_1 + \sqrt{b-z}\sin\frac{s}{\sqrt{a-z}}e_2$ and the rolling $(R_u, t) = (R_u(s), t(s))$ (R_u is the rotation of decreasing angle $u = u(s)$ and t is a translation) of E on the curve $c(s) :=$

$c^1(s)e_1 + c^2(s)e_2 :\quad (R_u, t)E := R_uE + t = c,\ R_udE = dc$. We need the sum of the elevation of the a-priori inclined segment $R_uG,\ G := F - E$ and c^2 to be 0; since $|R_u(s)G(s)| = |G(s)| = \sqrt{b-z} - \sqrt{b-a}\sin\frac{s}{\sqrt{a-z}}$, the obvious choice of $c^2(s)$ (for which there is no inclination, just as for the circle case) is thus revealed in the static picture: $c^2(s) := -|G(s)|$. From $|dc|^2 = |dE|^2$ we get $c^1(s) = s$; now the rotation R_u is uniquely determined by $R_udE = dc$ and the translation is $t := c - R_uE$. This is the only choice, as the imposed 0 elevation ansatz boils down to a first order differential equation and thus a second preserved quantity is revealed. Conversely, if we impose the non-inclination ansatz $G = |G|R_{-u}e_2$, then we get the same solution. The rolled focus $(R_u, t)F$ describes the horizontal line $R_uG + c = se_1$ uniformly in s; again this provides another conserved quantity which generalizes the circle situation. Differentiating the non-inclination ansatz and using $R_udR_{-u} = -R_{\frac{\pi}{2}}du$ we get $|G|due_1 = -R_udE - d|G|e_2 = -dc + dc^2e_2 = -dse_1$, so $ds = -|G|du$. Note also that the non-inclination ansatz is similar to the polar coordinates at F: if the point E is seen under the increasing angle θ from the focus F such that $\theta = 0$ for $s = -\frac{\pi\sqrt{a-z}}{2}$, then $G = |G|R_\theta e_2$, so $u = -\theta$. As E rolls on c it is not at equilibrium wrt the gravitational influence (constant and pointing in the $-e_2$ direction), but this can be corrected by considering a second counter-balancing rolling ellipse attached to E by a rigid bar connecting their foci (thus of length $(2k+1)\sqrt{a-z}\pi,\ k \in \mathbb{N}$). Because of the non-inclination ansatz, the length of this rigid bar remains constant during rolling. There are further preserved quantities of the system of two ellipses which generalize the circle situation. The system (possibly with a weight attached at the middle point of the rigid bar) is in equilibrium wrt the gravitational influence: each ellipse pulls the rigid bar in opposite directions and with force a constant multiple of $\frac{\frac{d}{ds}|G(s)|}{1+(\frac{d}{ds}|G(s)|)^2}$. The gravitational energy of the system is preserved, as $G(s + (2k+1)\sqrt{a-z}\pi) = G(-s),\ \frac{F^TG(-s)}{|G(-s)|} = -\frac{F^TG(s)}{|G(s)|}$, so the heights of the centers of the two ellipses are equal in absolute value and of opposite signs.

From $|G(s)| + |G(-s)| = 2\sqrt{b-z}$ we get $\frac{|G(s)||G(-s)|}{a-z} = \frac{4(b-z)-|G(s)|^2-|G(-s)|^2}{2(a-z)} = 1 + (\frac{d}{ds}|G|)^2$, or $\frac{1}{2|G|^2}|\frac{d}{ds}E|^2 - \frac{\sqrt{b-z}}{a-z}\frac{1}{|G|} = -\frac{1}{2(a-z)}$; differentiating this we get $\frac{d^2}{ds^2}|G| = \frac{1}{|G|}(1+(\frac{d}{ds}|G|)^2) - \frac{\sqrt{b-z}}{a-z}$. The parameter s behaves well for the following reason: with a new parameter $\hat{s} := \frac{s}{\sqrt{a-z}}$ independent of z an affine correspondence (Ivory affinity) with good metric properties is established between confocal ellipses: $E = \begin{bmatrix}\sqrt{1-za^{-1}} & 0\\ 0 & \sqrt{1-zb^{-1}}\end{bmatrix}(\sqrt{a}\cos\hat{s}e_1 + \sqrt{b}\sin\hat{s}e_2)$. When z varies we get a hyperbola, as an orthogonal trajectory of the family of confocal ellipses.

Consider the variables a, b, z, s, θ linked by the algebraic relation $G = |G|R_\theta e_2$, but otherwise independent; we have $R_{-\theta}dG = -|G|d\theta e_1 + d|G|e_2,\ |G|d\theta = \frac{-1}{|G|}dG^TR_{\frac{\pi}{2}}G = ds - sd\log\sqrt{a-z} - \frac{\sqrt{a-z}\sqrt{b-z}}{\sqrt{b-a}}\cos\frac{s}{\sqrt{a-z}}d\log\sqrt{\frac{a-z}{b-z}}$. Thus the differential system $G = |G|R_\theta e_2,\ ds = |G|d\theta$ has as consequence the fact that $d(a-z) = 0$ *if and only if* (iff) $d(b-z) = 0$ iff s is independent of $a-z, b-z$ (in which case s can be taken as a universal variable; the system admits only the free parameter z, so $d(b-a) = 0$ and the rolling problem holds); if $d(a-z) \neq 0$, then one can functionally solve for $s = s(a-z, b-z, \frac{d(a-z)}{d(b-z)})$.

Note that $0, z < a < b,\ s \in \mathbb{R}$ for the real case, but the computations are valid over complex numbers, so one can play with the order of the real numbers $0, a, b, z$ and also choose $s \in i\mathbb{R}$; $(\sqrt{\epsilon_1}\mathbb{R}) \times (\sqrt{\epsilon_2}\mathbb{R}),\ \epsilon_1, \epsilon_2 := \pm 1$ instead of $\mathbb{R}^2$; $(\cosh(s), \sinh(s)) = (\cos(is), -i\sin(is))$ instead of $(\cos(s), \sin(s))$ and changing the Euclidean scalar product via the linear transformation $[x\ y]^T \mapsto [x\ iy]^T$ or by multiplication by i.

Kepler's second law can be stated: $2 = |G|^2\frac{d\theta}{dt} (= |G|\frac{ds}{dt})$; thus t is the new universal variable (independent of $a-z, b-z$) and $s = s(t, a-z, b-z),\ \theta = \theta(t, a-z, b-z)$; now $\frac{d^2}{dt^2}E = \frac{ds}{dt}\frac{d}{ds}(\frac{ds}{dt}\frac{d}{ds}E) = \frac{4}{|G|}\frac{d}{ds}(\frac{-1}{|G|}R_\theta(e_1 + \frac{d}{ds}|G|e_2)) = 4(1+(\frac{d}{ds}|G|)^2 - |G|\frac{d^2}{ds^2}|G|)\frac{R_\theta e_2}{|G|^3} = \frac{4\sqrt{b-z}}{a-z}\frac{1}{|G|^2}\frac{G}{|G|}$ and thus agrees with the inverse square distance law of universal gravitational attraction, on the condition that the mass of the sun $\frac{\sqrt{b-z}}{a-z}$ is independent of $a-z, b-z$. But this is where Kepler's third law states what

is needed: the mean radius of the orbit is $\sqrt{b-z}$; due to the metric properties of the foci this is true for any probability measure on the ellipse E having the symmetry which exchanges the foci; in particular the period of the orbit is $\int_0^T dt = \frac{1}{2}\int_0^{2\pi\sqrt{a-z}} |G| ds = \pi\sqrt{a-z}\sqrt{b-z}$. Note the conservation of energy $\frac{1}{2}|\frac{d}{dt}E|^2 - \frac{4\sqrt{b-z}}{a-z}\frac{1}{|G|} = -\frac{2}{a-z}$ and the fact that the mass of the sun $\frac{\sqrt{b-z}}{a-z}$ is the maximum of the curvature of the ellipse E, obtained for $s \in \pi(\frac{1}{2}+\mathbb{Z})$*.

Thus Newton's differential system can be interpreted as the second in a hierarchy of differential systems whose first one is the rolling problem and the properties of all such differential systems emerge from the static picture.

Most known integrable systems come in hierarchies.

Although any two plane curves can be rolled one onto the other, there exists a notion of *osculating rolling* of two space curves of same curvature (and thus the rolling is restricted) which naturally appears as a degeneration of rolling surfaces. The tangent bundle of a plane curve is identified with a mostly multiple cover of a certain region of the plane of the curve; if one deforms this mostly multiple covered flat region, then we get the tangent surface (which is flat) of a space curve having the same curvature as the initial plane curve. One can roll this flat surface on the plane such that the two curves osculate and roll one onto the other. If the space curve is plane, then it must be congruent to the original one. In our case if one rolls the ellipse on the other side of the same (fixed) ellipse (at any instant the two ellipses reflect in the common tangent line), then its foci describe circles centered at the other foci, just as in the circle case. The same statement remains true for hyperbolas and parabolas (parabolas have a finite focus and a focus at ∞; thus a circle has its centers on the line at ∞ and becomes a line; the other rolling focus on the line at ∞ describes the line at ∞). For the ellipse we have the normal direction $\hat{N} := \frac{\sqrt{b-z}}{\sqrt{a-z}}\cos\frac{s}{\sqrt{a-z}}e_1 + \sin\frac{s}{\sqrt{a-z}}e_2,\ \hat{N}^T dE = 0,\ \hat{N}^T G = -|G|$ and the rolling focus $F' := F - \frac{2\hat{N}^T G}{|\hat{N}|^2}\hat{N}$ satisfies $|F'+F|^2 = 4|F + \frac{|G|}{|\hat{N}|^2}\hat{N}|^2 = 4(|F|^2 + \frac{|G(s)||G(-s)|}{|\hat{N}|^2}) = 4\frac{b-z}{b-a}$.

The case when conics osculate and roll on space curves having the same curvature is discussed in § 6.5 (B transformations of singular quadrics).

While intuitively the metric-projective character of the confocal family of the ellipse is clear, it lacks the advantage of generality, as it must be separately stated for the parabola (the confocal family of the ellipse also includes hyperbolas for $a < z < b$) and for imaginary conics. The metric-projective definition thus must be independent of the chosen conic and it was found by Plücker. Earlier Poncelet introduced the imaginary points $[1, i, 0],\ [1, -i, 0]$ at ∞ to account for the fact that two circles must intersect at four points, just as any other two conics in general position (Apollonius and Bézout); now Cayley's absolute $C(\infty) := \{[1, i, 0],\ [1, -i, 0]\}$ encodes the metric structure of $(\mathbb{C}^2, <, >)$, as rigid motions and homotheties are the only *homographies* (projective transformations) which preserve $C(\infty)$. Note that $C(\infty)$ is the quadric $(x^1)^2+(x^2)^2 = 0$ in the line at ∞ : $[x^1, x^2, 0],\ (x^1, x^2) \neq (0,0)$ of $\mathbb{CP}^2$. The ellipse is given by the equation $\frac{(x^1)^2}{a} + \frac{(x^2)^2}{b} = 1,\ a, b > 0$, the hyperbola by the same equation with $a > 0 > b$ and the parabola by the equation with $\frac{(x^1)^2}{a} = 2x^2,\ a > 0$; all conics are given by the equation $x^T A x = 0,\ x = [x^1\ x^2\ 1]^T \simeq [x^1, x^2, 1]$ with certain matrices $A = A^T \in \mathbf{GL}_3(\mathbb{C})$. One can define the *adjugate* of the conic x : $x^T A x = 0$ to be the conic x^* : ${x^*}^T A^{-1} x^* = 0$; if $y = Hx$ for $H \in \mathbf{PGL_3}(\mathbb{C})$, then $y^* = (H^T)^{-1}x^*$ and the 'adjugate' denomination becomes clear. $C(\infty)$ is by definition a self-adjugate quadric in the line at ∞ (this is equivalent to bringing an arbitrary non-degenerate scalar product $< x, y >:= x^T Q y,\ Q = Q^T \in \mathbf{GL}_2(\mathbb{C})$ to the canonical form $< x, y >:= x^T y$).

The family $\{x_z\}_{z\in\mathbb{C}}$ of conics confocal to the given conic x is *the adjugate of the pencil of conics generated by the adjugate of the initial conic and* $C(\infty)$: $x_z^T(A^{-1} - zI_{1,2})^{-1}x_z = 0,\ I_{1,2} :=$ diag[1 1 0]; thus $x_0 = x$ and the mixed metric-projective character becomes clear.

---

* I was reminded by one of my Calculus III students that the area element for a graph in polar coordinates is $\frac{1}{2}r^2 d\theta$ instead of $r^2 d\theta$; thus the area element is $d \wedge (\frac{1}{2}r^2 d\theta) = r dr \wedge d\theta$.

For $z := \infty$ we obtain the line at ∞ as a singular conic x_∞ of the confocal family; it contains as a singular set the quadric $C(\infty)$. Similarly for z inverses of non-zero eigenvalues of the quadratic part of the conic ($z := a$ or b for ellipses or hyperbolas and $z := a$ for parabola) we obtain singular conics x_z of the confocal family (lines); again these contain as singular sets other quadrics: the two foci (the parabola has a finite focus and a focus on the line at ∞). Thus the family of confocal conics can be interpreted as the adjugate of the pencil of conics generated by the adjugate of the quadric consisting of the two foci and the quadric $C(\infty)$.

Plücker's characterization of confocal quadrics is: for ellipses and hyperbolas each conic of the family is tangent to the four lines joining the two foci and $C(\infty)$ (the two other points of intersection of these four lines are imaginary foci); for parabolas each conic of the family is tangent to the three lines joining the foci and $C(\infty)$. This characterization follows from simple projective arguments.

An easy consequence of this metric-projective definition is that if H is a homography such that the line containing $H^{-1}(C(\infty))$ intersects the line containing $C(\infty)$ at a point not in $C(\infty)$, then H takes the family of confocal quadrics with foci among $H^{-1}(C(\infty))$ to the family of confocal quadrics with foci among $H(C(\infty))$, while preserving all metric-projective properties and thus all integrable systems whose integrability boils down to these metric-projective properties (see also Darboux ([13],§ 603)).

Stepping on the path opened by Euler half a century earlier (namely differential geometry of surfaces, dynamics of a rigid body and fluids), Dupin, Ivory and Lamé met in the beginning of the XIX$^{\text{th}}$ century at confocal quadrics (some of the properties of confocal quadrics were also known to Monge, who integrated the equation of lines of curvature on the ellipsoid, in the process introducing the notion of *umbilic*; it is less known since Darboux that there are 12 umbilics on the general ellipsoid, among which four are the usual real ones; if one excludes the importance of the remaining imaginary 8, then one must also exclude the D transformation of special isothermic surfaces and thus Bianchi's theory of deformations of quadrics). They were not primarily interested in the geometrical properties of confocal quadrics, but in natural phenomena amenable to a mathematical formulation at the time: (optical properties of) light, gravity and heat.

Dupin studied light and its reflection in surfaces; geometrically this amounts to normal congruences. These congruences remain normal after reflection and refraction (Malus) in surfaces; a surface normal to the normal congruence is cut by the *flats* (developables) of the congruence along lines of curvature. Developables are important for the following reason: the normal congruence is tangent to its two focal surfaces (caustics) and thus it induces direction fields on them; the integral curves of these direction fields (geodesics) give the developables (each developable is generated from the lines of the congruence by keeping constant a parameter of an integral curve and varying the parameter of the other or equivalently they are the positions of the line in a plane as that plane rolls along the line on the focal surface).

Ivory studied the gravitational attraction of ellipsoids; in particular he transformed the problem of the gravitational attraction a homogeneous ellipsoid exercises upon an exterior point to the problem of the gravitational attraction a confocal homogeneous ellipsoid exercises upon a corresponding interior point; his work on the ellipsoidal equilibrium configuration of self-gravitating fluids was an extension of that of Laplace (Laplace kept Ivory in high esteem for his critical commentary of *Mècanique céleste*) and influenced the results of Jacobi and Liouville on the geodesic flow.

Lamé was interested in heat and confocal ellipsoids provide an example of level sets of the temperature for the stable heat equation with a singular ellipsoid of the family (the convex hull of the singular set which is ellipse) being the source of constant temperature; since the gradient of the temperature is normal on its level sets, this provides an easy physical justification of Lamé's result. The actual problem Lamé was interested is more general: *To find all triply orthogonal families formed by isothermic surfaces*. Bonnet completely solved it and showed that, excluding developables, only quadrics can appear as surfaces in such families (see Darboux ([13],§ 513),([14],Ch III)).

The Ivory affinity between two confocal quadrics can be easily seen now: the temperature flow from a point on the first quadric cuts the second quadric at the point which gives the Ivory affinity.

Consider the Euclidean scalar product on $\mathbb{C}^n$: $< x, y >:= x^T y,\ |x|^2 := x^T x$. A *quadric* $x \subset \mathbb{C}^n$ is given by the equation $Q(x) = 0$, where $Q(x) := \begin{bmatrix} x \\ 1 \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} x \\ 1 \end{bmatrix} = x^T(Ax + 2B) + C,\ A = A^T \in \mathbf{M}_n(\mathbb{C}),\ B \in \mathbb{C}^n,\ C \in \mathbb{C},\ \begin{vmatrix} A & B \\ B^T & C \end{vmatrix} \neq 0$; x is called *degenerate quadric* if $\begin{vmatrix} A & B \\ B^T & C \end{vmatrix} = 0$. A quadric x belongs to either of the types: *quadric with center* (QC) for $\ker(A) = 0,\ Q(-A^{-1}B) \neq 0$, the center being $-A^{-1}B$; *quadric without center* (QWC) for $\ker(A) = \mathbb{C}v,\ v^T B \neq 0,\ |v|^2 \neq 0$; *isotropic quadric without center* (IQWC) for $\ker(A) = \mathbb{C}v,\ v^T B \neq 0,\ |v|^2 = 0$. After applying suitable rigid motions, most quadrics are *diagonal* QC (A is diagonal, $B = 0, C = -1$).

One can complete a quadric x to a family of confocal quadrics $\{x_z\}_{z \in \mathbb{C}}$: the adjugate of the pencil of quadrics generated by the adjugate of x and the adjugate of Cayley's absolute (taken to be self-adjugate). Thus $x_z \subset \mathbb{C}^n$ is given by $Q_z(x_z) = 0$, where $Q_z(x_z) := \begin{bmatrix} x_z \\ 1 \end{bmatrix}^T (\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} - z \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix})^{-1} \begin{bmatrix} x_z \\ 1 \end{bmatrix}$, so $x = x_0$.

The condition that a *nonsingular quadric* (z is not inverse of non-zero eigenvalue of A and $z \neq \infty$) of a *general confocal family* (x does not have continuous rotational symmetries; the obvious choice for the origin is clear for QC; a similar choice will appear later for (I)QWC) contains a point $x \in \mathbb{C}^n$ is a polynomial of degree n in z which has in general distinct roots. Therefore through any point $x \in \mathbb{C}^n$ pass in general n quadrics of the confocal family: $x \in x_{z_j},\ j = 1, ..., n$; along the higher co-dimension set of $\mathbb{C}^n$ where normals are isotropic pass less than n confocal quadrics. With certain modifications the statement remains true for singular quadrics of the family or non-general quadrics. Replacing the cartesian coordinates of x with $(z_j)_{j=1,...,n}$ gives elliptic coordinates on most of $\mathbb{C}^n$ (Lamé). For non-general quadrics one can complete the set of such $(z_j)_{j=1,...,n-k}$ with k coordinates of the space obtained by factoring the continuous group of rotational symmetries with the subgroup fixing a general point of the quadric, since all confocal quadrics will admit the same continuous group of rotational symmetries. Thus all non-general quadrics are general in a non-isotropic subspace of $\mathbb{C}^n$ which intersects orthogonally all trajectories of points of $\mathbb{C}^n$ under the action of a continuous group of symmetries (lines passing through centers of spheres for spheres and planes passing through the axis of revolution for quadrics of revolution for $n = 3$) and conversely from general quadrics one can obtain non-general ones in higher dimensions by acting on the surrounding space of the initial general quadric with a continuous group of rotational symmetries of the bigger space so as to fill it. The sense in which this statement can be interpreted will be explained later; intuitively a quadric x is general iff all eigenvalues of A have geometric multiplicity 1 (including the metrically most degenerate case $\text{spec}(A) = \{0\}$ which thus must be an IQWC). Thus the use of elliptic coordinates (completed with other coordinates for non-general quadrics) is essential in the *metric classification* of all quadrics and in studying integrable systems related to them (as shown since Jacobi and Liouville, most known integrable systems boil down one way or another to the use of elliptic coordinates).

Up to homographies all possible degenerate quadrics appear as singular quadrics of a certain confocal family (one can find a non-degenerate quadric in a hyperplane such that the pencil generated by the adjugates (in a sense that will become clear later) is mostly formed by non-degenerate quadrics and consider the later quadric as Cayley's absolute).

The classical theorems about confocal quadrics are as follows:

(Lamé) If x is a general quadric, then the normal to x_z is proportional to $\partial_z x_z$ and the vectors $(\partial_z|_{z=z_j} x_{z_j})_{j=1,...,n}$ are orthogonal; thus the family of quadrics confocal to a given general quadric forms an orthogonal system.

(Dupin) General confocal quadrics cut each other along principal sub-manifolds; thus they admit lines of curvature parametrization (the elliptic coordinates).

Thus Dupin's indicatrix (essentially the shape) is encoded by the confocal family.

If the quadric is not general, then one can still complete the confocal family to an orthogonal family; for example confocal quadrics in $\mathbb{C}^3$ and of revolution around an axis can be completed with the family of planes passing through the axis of revolution. For confocal (pseudo-)spheres the indeterminacy is even higher, because the principal lines are not defined; however the level sets of the standard spherical coordinates (planes through an axis and circular cones with the same axis) appear as a natural choice.

This result subsists for surfaces applicable to quadrics only at the level of the second fundamental form missing mixt terms (*conjugate* system of coordinates; in this matter Dupin preserved the denomination chosen by Apollonius since at the level of Dupin indicatrix the two coincide). The fact that Apollonius was onto something is confirmed by the simple projective explanation of conjugate systems which preserves the property of normal conjugate diameters (corresponding to principal directions on a surface): the ruled surface generated by conjugate directions along a curve on a surface is developable (a projective notion, since asymptotic directions coincide on such surfaces and osculating planes of asymtotes are tangent to the surface) and any two pairs of conjugate diameters of a conic are in harmonic ratio; moreover Apollonius introduced the tangent to a conic with center among others as the parallel to a conjugate diameter through the given point (which thus becomes the endpoint of a diameter). Conversely if we bend a plane on a surface along a curve, then the rulings on the resulting developable are the conjugate directions of the curve. Thus the statement 'triply orthogonal system of quadrics' becomes 'triply conjugate system of surfaces containing a family of surfaces applicable to quadrics'; triply conjugate system of surfaces means surfaces cutting each other along conjugate lines (see Bianchi ([5],Vol **4**,(143),(146))). Because the principal lines on (pseudo-)spheres are indeterminate, the triply conjugate system of surfaces containing a family of CGC surfaces are actually triply orthogonal (the common conjugate system on a surface and on the unit sphere is realized by the lines of curvatures on the given surface and the correspondence is given by the Gauß map). Thus Calapso studied deformations of quadrics by imposing the natural condition that a conjugate system on the initial quadric is also a conjugate system on one of its deformations; by this *infinitesimal analytic* method (although more complicated than Bianchi's simple geometric picture) he managed to take Bianchi's results up a notch in 1912.

(Ivory) There exists an affine transformation of $\mathbb{C}^n$ which takes x_0 to x_z; it preserves the lengths of segments between confocal quadrics: with $V_0^1 := x_z^1 - x_0^0,\ V_1^0 := x_z^0 - x_0^1$ we have $|V_0^1|^2 = |V_1^0|^2$ for pairs of points $(x_0^0, x_z^0),\ (x_0^1, x_z^1)$ corresponding on (x_0, x_z) under this affine transformation.

Such an affine transformation is called *the Ivory affinity* because it is friendlier than an usual affine transformation. In the same vein we have the result mostly due to Bianchi (see (122,§ 10)):

(Bianchi I) The Ivory affinity preserves lengths of segments on rulings (segments on tangent cones) of confocal quadrics, the tangency configuration (TC), angles between segments (between confocal quadrics) and rulings, between rulings and between polar rulings.

These can be written as: $|w_z|^2 = |w_0|^2$ if $x_0^0 + tw_0 \in x_0,\ x_z^0 + tw_z \in x_z,\ \forall t \in \mathbb{C}$ and $w_0,\ w_z$ correspond under the Ivory affinity; $(V_0^1)^T \partial_z|_{z=0} x_z^0 = (V_1^0)^T \partial_z|_{z=0} x_z^1$, thus $x_z^1 \in T_{x_0^0} x_0$ iff $x_z^0 \in T_{x_0^1} x_0$; $(V_0^1)^T w_0^0 = -(V_1^0)^T w_z^0,\ (w_0^0)^T w_z^1 = (w_z^0)^T w_0^1,\ (w_z^0)^T \hat{w}_z^0 = (w_0^0)^T \hat{w}_0^0$ if $x_0^0 + sw_0^0 + t\hat{w}_0^0 \in x_0,\ \forall s, t \in \mathbb{C}$.

Thus for $x_0^0,\ x_0^1$ points on a quadric $x_0 \subset \mathbb{C}^{2n-1}$ and $W_0^0,\ W_0^1$ maximal sets of mutually polar rulings at $x_0^0,\ x_0^1$ (each containing $n-1$ rulings), with $F := [V_0^1\ W_0^0\ W_z^1]$ we have $F^T F = ((0 \leftrightarrow z) \circ F)^T ((0 \leftrightarrow z) \circ F)$ and there exists a RMPIA (R_0^1, t_0^1) such that $R_0^1 F = (0 \leftrightarrow z) \circ F,\ t_0^1 = x_z^0 - R_0^1 x_0^0 (= x_0^1 - R_0^1 x_z^1)$; moreover (R_0^1, t_0^1) is unique precisely when $\det F \neq 0$.

The preservation of lengths of segments on rulings under the Ivory affinity between confocal quadrics ($n = 3$) give 'almost' Tchebyshev coordinates which are further asymptotic; they are 'almost' because in general we cannot change these coordinates to make the rulings of unit length. Thus we get (up to rigid motions) the family of confocal quadrics if we deform the doubly ruled fabric of an initial quadric, changing the angles between the ruling families while preserving their

lengths. This construction of the articulated hyperbolic paraboloid is due to Henrici so some of these results may have already been folklore at the time (see Bianchi [4]).

In 1838 Jacobi integrated the equation of the geodesics on an ellipsoid; geometrically his result can be stated:

(Jacobi) The tangent lines to a geodesic on x_0 remain tangent to $n-2$ other confocal quadrics.

This is the first instance when the simple metric-projective definition of confocal quadrics has as consequence a deep metric-projective property, which hints at the fact that the confocal family of a quadric influences surfaces applicable to that quadric. Chasles proved its converse:

(Chasles) The common tangents to $n-1$ confocal quadrics form a normal congruence and envelope geodesics on the $n-1$ confocal quadrics.

Thus confocal quadrics not only *infinitesimally* cut each other orthogonally (Dupin and Lamé) but also *finitely*, that is as seen by the viewer from any point of the space (Darboux). This is another principle: *infinitesimal* laws for confocal quadrics admit corresponding *finite* laws (and according to Archimedes the converse is true); the theory of deformations of quadrics is one of its facets and its discretization realizes the complete unification of the *infinitesimal* and *finite*.

Consider the lines $x_{z_1}+sv,\ s\in\mathbb{C}$ tangent to the $n-1$ quadrics $x_{z_j},\ j=1,...,n-1$ at points $x_{z_j}:=x_{z_1}+s_{z_j}v$. As we shall see later, this implies that the normals along the common tangents are orthonormal: $N_{z_j}^TN_{z_k}=\delta_{jk},\ k=1,...,n-1$; thus 0, $\{z_j\}_j$ can be considered eigenvalues with corresponding eigenvectors $v,\ \{N_{z_j}\}_j$ for a certain symmetric operator which undergoes an isospectral deformation. The approach to integrable systems by means of isospectral deformation of a linear operator was first developed by Lax for KdV and it has since emerged into a powerful technique used to study most known integrable systems. Moser established in [24] and via the above considered symmetric operator links between confocal quadrics and other integrable systems, thus returning to the basics: confocal quadrics capture (encode) and geometrize spectral properties of symmetric operators.

While there are many transformations of $\mathbb{C}^n$ which take a family of confocal quadrics to another one, a transformation which preserves the previous properties of confocal quadrics must take lines to lines, so it must be a homography. The converse is mostly true and due to Bianchi (see ([5],Vol **5**,(90))):

(Bianchi II) For any general homography H of $\mathbb{C}^n$, there are two unique families of confocal quadrics in $\mathbb{C}^n$ such that H takes one family to the other. Moreover, it takes principal directions ($\partial_z|_{z=z_j}x,\ j=1,...,n$, where $\mathbb{C}^n\ni x\in x_{z_j}$) to principal directions (Lamé) and geodesics to geodesics (Chasles). Thus H induces a linear map on each tangent space of $\mathbb{C}^n$: $dH(x):T_x\mathbb{C}^n\mapsto T_{H(x)}\mathbb{C}^n$ and the singular value decomposition (SVD) of $dH(x)$ gives the principal directions in both the domain and the range of $dH(x)$. Moreover the Ivory affinities between (x_0,x_z) and $(H(x_0),H(x_z))$ (and thus the RMPIA) correspond under H; also H preserves the isospectral deformation character of the above considered symmetric operator.

Note that $A=A^T\in\mathbf{M}_n(\mathbb{C})$ cannot always be diagonalized via a conjugation with a rotation $R\in\mathbf{O}_n(\mathbb{C})$; we can hope at best to obtain a *symmetric Jordan* (SJ) form (a form as close as possible to a diagonal form and depending on as few as possible constants). Thus the SVD does not always reduce $A\in\mathbf{GL}_n(\mathbb{C})$ to a diagonal matrix: $A=R_1DR_2^T,\ R_1,\ R_2\in\mathbf{O}_n(\mathbb{C})$ and D is SJ (R_1,R_2,D can be mostly found from $A^TA=R_2D^2R_2^T,\ AA^T=R_1D^2R_1^T$). Note also that the two families of confocal quadrics must have the same continuous group of rotational symmetries.

The natural question about the existence of the SJ form appeared since I was using identities of the form $A\sqrt{I_n-zA}=\sqrt{I_n-zA}A,\ A=A^T\in\mathbf{M}_n(\mathbb{C}),\ z\in\mathbb{C}\setminus\mathrm{spec}(A)^{-1}$.

Note that according to Apollonius any line cuts two homothetic QC along two equal segments thus individuated (including the singular cases when the homothety is $0,\infty$, so the QC becomes a quadratic cone or the hyperplane at ∞). Since this is not a true metric-projective property, the natural question arises if the result still stands by replacing the pencil of homothetic QC with the family of confocal quadrics. The answer is no, but because of the symmetry of the TC the next natural question arises wether a segment between two points on a quadric and whose subjacent

line is tangent to a confocal quadric is taken by the Ivory affinity to a ray of light trajectory (that is reflecting in the initial quadric) and tangent to the given confocal quadric at two points. The answer is not only is yes, but also its converse (a ray of light tangent to a quadric and reflecting in a confocal quadric is still tangent to the given quadric; moreover the Ivory affinity takes the broken segment into a single one) is valid; applying Ivory's Theorem and the symmetry of the TC we get an intuitive a-priori idea about rays of light tangent to a quadric and reflecting on a confocal one (note that any such ray comes with a dual friend of same length obtained from the Ivory affinity). In this sense we have a result due to Chasles-Darboux about a ray of light whose trajectory is formed by polygonal segments on lines circumscribed (tangent) to $n-1$ given confocal quadrics and bouncing (reflecting) sequentially on another given sequence of quadrics $Q_1, Q_2, ...$ of the same confocal family as the initial one: if one such trajectory is closed, then all are; moreover all such trajectories have same length (see Darboux ([13],§ 465)). Note again the principle of correspondence between *infinitesimal* and *finite* by means of *discretization*: the rolling of an $(n-1)$ dimensional space along one of its lines on a quadric by means of which one can obtain Jacobi's geodesics becomes at the discrete level *precisely* the construction and motivation of Darboux's interpretation (due to the good metric properties of the Ivory affinity as pointed out by Ivory himself). One can replace the ray of light with a billiard ball; billiards on confocal quadrics is a current subject of research in integrable systems (due mostly to Serge Tabachnikov). Of course even such things are preserved by the homography of Bianchi II and one can investigate them first in the complex setting and then restrict, as an application, to totally real cases*.

Delaunay proved in 1841 that if we roll a conic on an line and then rotate around the line the curve described by its foci (focus), then we get all CMC (minimal) surfaces of revolution.

In 1859 the French Academy posed the problem:

To find all surfaces applicable to a given one.

It became the driving force which led to the flourishing of the classical differential geometry in the second half of the XIX$^{\text{th}}$ century and its profound study by illustrious geometers led to interesting results (see Bianchi [4],[5], Darboux [13], Eisenhart [19],[20], Sabitov [27] and its references for results up to 1990's or Spivak ([29], Vol **5**)). Today it is still an open problem in its full generality, but basic familiar results like the Gauß-Bonnet theorem and the Codazzi-Mainardi equations (independently discovered also by Peterson) were first communicated to the French Academy. A list (most likely incomplete) of the winners of the prize includes Bianchi, Bonnet, Guichard, Weingarten.

Weingarten proved in 1863 that the focal surfaces of the congruence of normals of a surface having a functional relationship between its curvatures are applicable to surfaces of revolution (which depend only on the functional relationship) and conversely, the tangents to geodesics corresponding to meridians on a surface applicable to a surface of revolution (meridians on the given surface) form a normal congruence, whose normal surfaces have a functional relationship between their curvatures. Such normal surfaces are called *Weingarten* (W). Note that in the correspondence established between the two focal surfaces by the normal W congruence to the meridians on a focal surface correspond parallels on the other, since on any normal congruence the developables of different families cut each other perpendicularly (Dupin and Malus). Since the two focal surfaces as above are applicable to surfaces of revolution, Darboux's statement about the deformation of a surface of revolution to a line has a simple geometric explanation: as a focal surface rolls on its applicable surface of revolution, the other focal surface (called *complementary* surface) rolls on the axis of revolution, with the arc-length of the meridians corresponding to the arc-length of the axis of revolution (each meridian on the second focal surface rolls on the axis of revolution precisely when the first focal surface rolls on the applicable surface of revolution along a parallel). Or: as

* At this point the fact that the Ivory affinity provides the correct explanation is pure speculation; Darboux uses differential computations in elliptic coordinates; should the trick with the Ivory affinity work one can use instead algebraic computations.

the first focal surface is deformed to its corresponding surface of revolution, the complementary surface is deformed to the axis of revolution, each meridian becoming the axis of revolution.

Ribaucour proved that the second fundamental forms of focal surfaces as above are proportional; therefore Bianchi calls a congruence on whose focal surfaces the second fundamental forms are proportional a *Weingarten* (W) congruence. This notion is invariant under homographies, since involves only tangency: the asymptotic directions (a projective notion) correspond.

W congruences play an important rôle in the infinitesimal deformation problem (Darboux proved that the later generates W congruences and Guichard proved the converse); thus the infinitesimal deformation problem is essentially projective.

As an example Weingarten classified the W surfaces with $2(r_2-r_1) = \sin(2(r_2+r_1))$, in which case the lines of curvature correspond via the Gauß map to geodesic confocal ellipses and hyperbolas on the unit sphere. By 1894 Darboux proved that the *evolutes* (focal surfaces of the normal congruence) of such W surfaces are applicable to imaginary paraboloids of revolution, completed the study of such W surfaces by replacing sin with sinh (in which case the paraboloids become real) and provided a simple geometrical construction of such W surfaces: they are surfaces of translation with the two generating curves having constant complex conjugate torsions (or opposite real); their evolutes give all surfaces applicable to (imaginary) paraboloids of revolution (see ([13],§ 745-§ 751)). Therefore Darboux posed the natural question (still open in its full generality): *'To find all algebraic curves of constant torsion'* (see ([13],§ 39 and Note 4 of Vol **4**)).

Bäcklund constructed in 1883 a transformation for CGC -1 surfaces (this point of view is due, according to Bianchi ([4],§ 374), to Lie): the tangent spaces to the unit pseudo-sphere x_0 cut a confocal pseudo-sphere x_z along circles, thus highlighting a circle in each tangent space of x_0. Each point of the circle, the segment joining it with the origin of the tangent space and one of the (imaginary) rulings on x_z passing through that point determine a facet. We have thus highlighted a 3-dimensional integrable distribution of facets: its leaves are the ruling families on x_z. If we roll the distribution while rolling x_0 on an non-rigidly applicable surface x (called *seed*), it turns out that the integrability condition of the rolled distribution is always satisfied (we have complete integrability), so the integrability of the rolled distribution does not depend on the shape of the seed. The rolled distribution is obtained as follows: each facet of the original distribution corresponds to a point on x_0; we act on that facet with the rigid motion of the rolling corresponding to the highlighted point of x_0 in order to obtain the corresponding facet of the rolled distribution. Note that the facets of the 3-dimensional distribution are differently re-distributed into 2-dimensional families of facets as tangent planes to leaves when the shape of the seed changes, but principles and properties independent of the shape of the seed remain valid for facets even in the singular picture: this is Archimedes' contribution to the Bianchi-Lie ansatz. The leaves of the rolled distribution (called the B transforms of x, denoted $B_z(x)$ and whose determination requires the integration of a Ricatti equation) are applicable to the pseudo-sphere. Moreover the seed and any leaf are the focal surfaces of a W congruence, so the inversion of the B transformation has a simple geometric explanation (the seed and leaf exchange places). For simplicity we ignore in the notation $B_z(x)$ the dependence on the initial value of the Ricatti equation.

The transformation originally constructed by Bäcklund states that if a CGC -1 seed x in general position and an angle $0 < \theta < \frac{\pi}{2}$ are given, then the 3-dimensional distribution formed by facets with centers on circles of radius $\sin\theta$ in tangent spaces of x, of inclination θ to these and passing through the origin of these is integrable; moreover the leaves are CGC -1 surfaces. Lie's point of view becomes clear now: because the B transformation is of a general nature (independent of the shape of x), it must exist (at least as a limiting case) also in the case when x in not in general position, but it actually coincides with the pseudo-sphere. In this case the 1-dimensional family of non-degenerated leaves (surfaces) degenerates to a 1-dimensional family of degenerated leaves (isotropic rulings on confocal pseudo-sphere) and thus the true nature of the B transformation is revealed at the static level of confocal pseudo-spheres.

The B transformation when the confocal pseudo-sphere is the isotropic cone (sphere of zero radius or equivalently $\theta = \frac{\pi}{2}$) was constructed even earlier (1879, upon some results of Ribaucour from 1870) by Bianchi in his PhD thesis and named the complementary transformation, since the seed and the leaf are the focal surfaces of a normal W congruence. Ribaucour did not look at his results as a method of generating CGC -1 surfaces from a given one (transformation) and thus Bianchi is credited with the idea of using transformations in the study of surfaces: one can obtain families of CGC -1 surfaces depending on arbitrary many constants by iterating the complementary transformation (each constant being introduced by the integration of a Ricatti equation).

Lie proved the inversion of the complementary transformation; thus its iteration is realized by quadratures since a Ricatti equation degenerates to a linear one once a solution is known. He also discovered the 1-dimensional (spectral) family of CGC -1 surfaces, but this was only *infinitesimally* determined (the fundamental forms are known), with no explicit procedure of constructing it without the integration of Ricatti equations and quadratures; moreover this Lie transformation cannot be iterated. According to a formula recently discovered by Sym in [30], one can replace the quadrature step with a derivative in the spectral parameter; however the most difficult part (the integration of a family of Ricatti equations) remains. According to Bianchi the B transformation is a conjugation of a complementary transformation with a Lie transformation.

Bianchi proved in 1892 the BPT for the B transformation; it is with the BPT that the B transformation can be iterated using only algebraic computations (after first integrating a 1-dimensional family of Ricatti equations and taking some derivatives to find the rolling).

Up to Weingarten's prize winning article of 1896 ([38]), Darboux's intrinsic Monge-Ampère equation of isometric immersion of an abstract linear element already appeared in conjunction with the deformation problem, but Weingarten made the essential remark that the problem already assumes the existence of a given surface (see Darboux ([13],vol **4**, ch XIII and XIV) and Eisenhart ([19], ch X)). However, few applications of Weingarten's method have been since found; most of them give the already found results for surfaces of revolution or of Goursat's deformations of certain imaginary quadrics. Probably one must modify Weingarten's approach so as to take into consideration not only the existence of the given surface, but also that of a deformation (*seed*). The only question from an audience of graduate students after a talk about Weingarten's method in Spring 2004 (where I mentioned, just as Darboux does, an equation of Euler's in conjunction with the propagation of the sun, clearly an example of integrable systems) was where am I heading with this program. Looking back I would say that that is an important question; the answer would still elude me for more than two years. It has been an open question ever since the classical times to find all surfaces for which an interesting theory of deformations can be build, just as for quadrics (see § 5.17 of R. Calapso's introduction to Vol **4**, part 1 of Bianchi's *Opere*).

In 1899 Guichard considered Q(W)C of revolution around the focal axis: when such a quadric rolls on an applicable surface, its foci (focus) describe(s) CMC (minimal) surfaces. Note that since a surface of revolution can be rolled on a line with the arc-length of the meridian corresponding to the arc-length of the line, Guichard's result is a generalization of Delaunay's. Note also that according to Bonnet when the unit sphere rolls on an applicable surface its center describes a CMC $\frac{1}{2}$ surface, so Guichard's result is a metric-projective generalization of Bonnet's result (compare with the rolling problem for the ellipse). Note also that when a Q(W)C of revolution around the focal axis rolls on itself (at each instant the fixed quadric and the rolling quadric reflect in the common tangent space) its foci (focus) describe(s) CMC (minimal) surfaces: spheres (planes), so Guichard proved that this property is essentially metric.

With Guichard's result the race to find the deformations of general quadrics was on.

It ended in 1906 with Bianchi's discovery; in the process the differential geometry underwent a fundamental change through the results of geometers involved in the study of quadrics: mostly Bianchi, Calapso, Darboux and Guichard, but a list of geometers with results related to quadrics and not mentioned so far is longer: Chieffi, Peterson, Ţiţeica (aka Tzitzeica), etc. In particular Peterson discovered certain surfaces (later on extensively studied by Bianchi and thus called *Bianchi*

surfaces; note however that The Arnold Principle applies here only partially, as Bianchi gave due credit to Peterson in [4] and he never calls them Bianchi surfaces; also Peterson's work became widely known in western Europe only after 1903 (see Darboux ([13],§ 111))) and Ţiţeica (see [37]) discovered the affine sphere (which led Blaschke to the development of the affine geometry) while studying certain deformations of quadrics. The *'illustre geometra rumeno'* Ţiţeica was a student of Darboux's (he got only a Master degree in France) and the founder of the modern geometry school from Romania a couple of decades after its independence, but in Romania he is known even to the non-mathematician, mostly due to his high-school level book of problems in Euclidean geometry (which is the standard even today). Upon his untimely death from a cerebral vascular accident in 1938 the two main pretenders to the position of Chair of the Geometry Desk at the Mathematics Department of the Bucharest University were Dan Barbilian and Gheorghe Vrânceanu. Dan Barbilian (mostly known for the *Barbilian geometries*, a generalization of Cayley's point of view) is also known in Romania as Ion Barbu, the pseudonym under which he published successful poetry; again his poetry is required subject of study in Romanian high-schools, so kids are tormented to comment on his messed up poetry (what other type of poetry a mathematician could create?). One could easily understand why the apparently messed up moving picture *'Pulp Fiction'* (the name itself explains: juicy raw rough difficult to believe underground story) is generous enough to admit interpretations in other settings (for this reason *'Pulp Fiction'* is actually a good model of *'Generic Essence of Pure Reality'*), because besides the natural flow of logical deductions (*dialogue, environment* and even *lyrics* further the story, in itself a patch of more than two decades' worth of pop culture; the facets of the infinitesimal picture (facets and bits of the story) fit perfectly and consistently integrate (by means of a flat connection form, namely QT) to the finite world thus providing the correct explanation of the moving picture as a whole) and episodes comparable with the elegance and simplicity of the proofs of the wise ancients (but not apparent at the first cursory incursion: just like good medicine and sometimes the bare truth tastes bitter, *the simplest possible explanation is always the true one*, no matter how confounding it is), Butch explaining the character of his day (*'without a doubt the single weirdest ... day of my life'*), whose chopper was his current chopper (*'I had to crash that Honda ... might've broke my nose'*) and when inquired *'Who's Zed?'* he replies with the apparently wrong answer *'Zed's dead, baby, Zed's dead'* (according to The Wolf *'time is a factor'* when we have *'a body in a car minus a head in a garage'* (no need for parenthesis, change of order of words or commas, as the whole sentence is associative, *'court'* and self-contained and on top of that the garage can be counted with either $\pm$), but the inquirer does not know it yet) is the actual scientist of the XVII$^{\text{th}}$ century in the first few days after being hit with the apple on his head, when the whole world collapsed with that apple and must then be rearranged anew: *interesting things are simple, already exist and can at most reveal themselves. 'Dead'* ain't no friend I ever heard of, just as *'What ain't no country I ever heard of'*; should such a country exist the natural question *'Do they speak English in What?'* immediately arises. First he does not believe and tries to disprove himself, but in doing so he already changed the laws of the solar system and sees no error yet and everything is *'Flowers on The Walls'* (*'Playing Solitaire till dawn with a deck of 51'* is a good picture of distributions of facets; if the croupier is fast enough to throw the last card before the first one touches the casino table and a snapshot is taken at that instant, then the perfect picture of a space distribution of facets appears; also if all cards are counted with their centers in one point and with 51 multiplicity, then a perfect picture of Archimedes' method appears); then it turns out that things are not so simple and rosy, so within minutes (a longer time for the scientist) the first *shadow of a doubt* appears; he tries to fend it off immediately (*'It's good to see you, I must go, I know I look a fright'*), but that turns out not to be so easy, so he feels the bounds of his own sanity within the reach of his bound hands and only then, in order to preserve one's sanity, he lets go of the old views as unimportant details, changes his own beliefs and unties his hands with a supreme effort and ultimately rides chop-chop out of town with the eery *'Twilight Zone'* theme omnipresent and providing the natural smooth change in time-line and on the first available chopper (*'Grace'*), not before paying to the Caesar of the land (always bearing

a Latin name; while bestowing *'oak'* on friends also the prefix *'Uncle'*) the debt he owed, just as everybody else (for example Jules) also does before leaving town, so as to be *'Kool and The Gang'*. But then he remains with fear of God for the remainder of his life, as he cannot find other rational explanation for the falling of the apple *precisely* when he was studying the movement of the moon as it was explained by his predecessors (or equivalently, while turning back for the forgotten heritage of his predecessors, studying an unusual strange object at hand and finding *without even thinking* its immediate and natural usefulness *precisely* when the toast ripened and was thus released, *The Brothers Startled* and one of them got smoke-toasted, thus fulfilling in the process the previously established *'grease spot'* denomination as a result of his beliefs in a *'freak occurrence'*; most of the *hyperbolic* denominations stick to their object and *parabolically* fulfill their purpose, even if Butch apparently has no meaning in English). So he finds it likely that God has created the world in a small number of days making thrift use only of the ruler, the compass, the balance and other simple mechanical methods (it seems that even God Himself took some recreational breaks from this endeavor to play some dice; in doing so some of the dice spilled and stuck to this endeavor and the resulting ambiguity had to be decided among others by means of *'eenie, meeny, miney, moe'*) and provides the explanation along these lines, passing through several interpretations until he finds the one which he believes finally fits the moving picture, just as Jules's *last minute* interpretation of *Ezekiel 12:57* in the actual *bony situation* (the story where some characters *'get into character'* and *'have character'* returns in a full circle to its true beginnings). Playing with the dice and things sticking and transferring to their apparently wrong place is common not only to DNA (where time and nature take over the resulting ambiguity and decide its rightfulness) or to human language (all languages are inherently ambiguous and continuously changing; as a consequence the closest relative in Romanian to the English *kitchen*, the French *cuisine* and the Italian *cucina* is *cocina* (pig sty; most of the time the correct description of my kitchen and probably the same explanation *parabolically* applied in the Middle Ages), a fact revealed to me on a trip to Chicago in Summer 2005 when I saw an Italian restaurant and began laughing instead of in the first year of the study of French and English), but in all aspects and facets of reality. When instructed *'Say something!'* or in the morning *'Say goodnight, Raquel!'* the instructed obeys the command shown in front of one's eyes and replies (*'quite rationally, in my opinion'*) *'Something!'* or *'Goodnight, Raquel!'*. There is another structured and mathematical story with a *Jack Rabbit's Slim hole* (it is mostly things that shock that are not forgotten, probably as a side effect of evolution): the *'Matrix Trilogy'* is a good model of the *infinitesimal world*, where Delaunay's surface (closely resembling UEFA's Delaunay trophy; strange things appear in front of one's eyes when one asks *Google, show me!* instead of *Open, Sesame!*) clearly appears as the trail of a bullet to make the point out of it and to provide the link to *gas dynamics*; also some information sticks (probably due to entropy) and a-priori inexplicably consistently transfers from The One to Agent Smith. Consequently I have a vague idea at this point about the mathematical definition of *equivariancy*, but I remember a talk where the speaker approached a member of the audience in the first row, winked (always winking with the right eye is the trick; understanding this trick puts the *I* in *Robot* and by entropy conversely) and identified oneself: *'I am your friend and you are my friend; thus I treat your other friends right and in return you will do the same'*; *consequently* the TC can be interpreted as something to do with tangent cones (as seen from x_z) taken by RMPIA's to cones of tangents (as seen from x_0), so the RMPIA may be the wink by means of which *cones* and *tangents* for quadrics befriended and sealed their friendship with the symmetry of the B transformation (that is Archimedes' balance) and the BPT. And by the way, *'Antwan should've ... better known better'* because Archimedes *'is The One that says 'Bad ...' '* (Marsellus himself had to learn the hard way that one cannot defeat Archimedes directly, so he *'took advantage of a poorly guarded fortification which he had seen during diplomatic negotiations'*; it may be a possible explanation of Archimedes' unfortunate premature death or so they say *'around the campfire'*): getting *'the technique down, I don't tickle or nothing'* is useless if one loses the focus and purpose of the idea: *'sticking'* facets with their centers at a point and with an (close to) ∞ multiplicity may provide the required *'charming ... pig'* as the correct substitution

to the initial unattractive one. Also even if the Dutch decided to put mayonnaise on French fries (*'I've seen them do it, man. They ... drown them in that ...'*), the choice of *'ketchup'* should be the natural one (there is a Romanian equivalent of the joke with three tomatoes: when the junior one is warned by a senior one to move from the road to the sidewalk because of the danger posed by cars it replies *etee, fleoşc!*; this can be translated as either *'eat my shorts!'* or *'eat my splashhh!'*; on the other hand the Romanians also have another saying: *'Orice pasere pre limba ei piere'*; this can be translated as either *'Every bird dies singing its song'* or, by entropy and not paying corresponding due attention to the cat, *'Every bird dies because of its song'*). Then he realizes that when God said *'Let there be light!'* it meant that God decided to disclose His Plans to man (*'at the request of'* man, since out of his own vanity (the devil's favorite sin) man chose the apple and later on went on A Mountain, repented and made *A Covenant*), so he begins looking in religious texts available at the time in hope of finding hidden messages from the wise ancients. I would say that this is the most logical and scientific step a deeply religious scientist of those times would have made, since God or a Librarian of Alexandria or a combination thereof, knowing that the age of the wise ancients was coming to an end, would have most likely chosen a religious text to partially hide and thus protect the knowledge in order to better preserve it for the next generations. Also at that time the wise ancients were regarded with the proper due respect (even during his lifetime there was a general consensus that the wise ancients reached a level of development of scientific investigation much higher than the current one; in particular this became clear with the almost complete version of Apollonius's *Conics* of 1710) and it was still believed (as inferred from the available literature) that an important part of their results was missing. It turns out that indeed the Word of God (wisdom: note that Archimedes was also hit with the apple; as a consequence he would run naked on the streets shouting like a person out of one's mind) was hidden in the Word of God (religious text), so *reading between the lines of a religious text of the time was indeed the best way to reach the wisdom of the wise ancients*. Indeed the finding is simple: to the inquiry *'You 'Flock of Seagulls' ... why don't you tell my man Vince here where you got the ... hid at'* the answer is of course the simple and logical *'It's in the cupboard'*, but the final delivery is a little bit more complicated: *'I cannot give this case to you because it don't belong to me'*. Vrânceanu (probably the most famous Romanian geometer) had been a student of Levi-Civita's and by that time even Cartan had already built on his ideas. The next oral story still circulates today at Bucharest University: it seems that Barbilian formally filed his request before Vrânceanu; after hearing about Vrânceanu formally filing the request (at that time Vrânceanu was Professor at Cernăuţi University), Barbilian warned the Committee charged to choose Ţiţeica's successor that the only way he would not become Chair and would not contest the findings of such a Committee is for Vrânceanu to become Chair, so the Committee had no choice but to put Vrânceanu Chair; this position was held until his retirement in 1970.

Bianchi generalized the B transformation of the pseudo-sphere to the B transformation of quadrics: the same statement remains true if in Lie's interpretation 'pseudo-sphere' is replaced with 'quadric' and 'circle' with 'conic'. The completion of the study of the B transformation was realized by Calapso in 1912 by taking into consideration the singular case $z = \infty$ for diagonal QC (see [11]).

A simple heuristic argument in favor of the existence of the B transformation for general quadrics goes as follows: the B transformation for CGC -1 surfaces is of a general nature (independent of the shape of the seed), so the cancellations of the shape of the seed in the integrability condition of the considered rolled distribution (equivalent to the complete integrability condition of the Ricatti equation) occur because of general equations (the GCM equations of the seed and of the pseudo-sphere) coupled with algebraic consequences of the TC (apparent in the static picture of confocal pseudo-spheres): these are valid for general quadrics. Thus the fact that a general quadric has less symmetries does not influence the existence of the B transformation.

However, because of the smaller group of symmetries, the applicability correspondence on CGC surfaces (which can be found in ∞^3 ways) does not have an easy generalization to quadrics. Without knowledge of the applicability correspondence it is very difficult in general to check if two surfaces are applicable, the test being similar to and much older than the Cartan-Ambrose-Singer theorem: basically one must find an arc-length correspondence between same level curves of the Gauß curvatures such that their orthogonal trajectories also admit an arc-length correspondence (see Eisenhart ([19],§ 136)). This test answers a problem posed by Minding in 1839: *'To find a necessary and sufficient condition that two surfaces be applicable'* (see Eisenhart [19]). Once the Minding problem was solved, the problem posed by the French Academy naturally appeared.

A first attempt is to prove that the correspondence between the seed and the leaf given by the W congruence is the applicability correspondence: the differential of the leaf depends on the shape (but not on its derivatives) of the seed, but in the linear element of the leaf the Gauß theorem does not cancel all the shape of the seed. Thus to follow the general method and find the applicability correspondence directly between the seed and the leaf (or between a surface in the initial quadric and the leaf) is out of question. According to Bianchi's own account from (122,§ IV), should the applicability correspondence exist in general, it can be presumed to be of a general nature and thus independent of the shape of the seed: therefore the answer lies not in the picture with the seed and leaf, but in the static picture with confocal quadrics and we have to roll back (or one can apply the ABL method). Thus to find the applicability correspondence one must look for a natural correspondence between confocal quadrics: the Ivory affinity provides such a correspondence and proving that the ACPIA is valid remained a matter of computations. This applicability correspondence is realized as follows: if we roll the seed x^0 on the applicable surface $x_0^0 \subseteq x_0$, then the point on the rolled B transform $x^1 = B_z(x^0)$ corresponding to the point of tangency of x_0^0 and the rolled surface x^0 is $x_z^1 \in x_z$. Now x_z^1 is taken by the Ivory affinity between x_0, x_z into a point x_0^1 on the initial quadric x_0: this is the applicability correspondence. Although the applicability correspondence is restricted, the structure becomes richer (a quadric with less symmetries has less degeneracies; for example a segment of B transformations degenerates to the complementary transformation for a quadric of revolution or the geodesic flow on an ellipsoid degenerates to the one on a sphere).

The necessary algebraic relations of the static picture do not naturally appear in the static picture from a geometric point of view, but are naturally chosen by the moving picture. For example if we assume the B transformation to exist and we roll the given quadric on a side of the seed, then we get two families of leaves, corresponding to the two ruling families on the confocal quadric. If we roll the initial quadric on the opposite face of the seed, then we get the same two families of leaves (otherwise we would have three or four ruling families on the confocal quadric), so the two rolled distributions reflect in the tangent bundle of the seed and this property is independent of the shape of the seed. According to the ABL method, the reflection in the tangent bundle of the seed property must be true also in the static picture, when the seed coincides with the initial quadric and the leaves are the ruling families on the confocal quadric. If one applies the same ABL method to the ACPIA, then one naturally obtains the RMPIA.

Now it becomes clear why the initial attempt to prove the applicability correspondence directly between the seed and the leaf will be unsuccessful: the leaf may be applicable to a surface $x_0^1 \subseteq x_0$ different from x_0^0. For example by applying the B transformation to a real seed x^0 applicable to the real ellipsoid $x_0^0 \subseteq x_0$ (and $z \in \mathbb{R}$ chosen so that the confocal quadric x_z intersects $\mathbb{R}^3$ along a hyperboloid with one sheet), the leafs are real surfaces, but they are not applicable to the given ellipsoid: they are applicable to another totally real region of x_0 (the ACPIA becomes *ideal*, in Peterson's denomination). To get back real surfaces applicable to the given ellipsoid we need to apply the B transformation to the leaf also. This is the reason why the B transformation for the sphere was developed much later, as the first application of the B transformation to real CGC 1 surfaces gives imaginary surfaces and one needs iteration of the B transformation to get back real CGC 1 surfaces.

Such is the simplicity of the answer to a question which eluded geometers for a long time! Bianchi's contemporaries acclaim *'Qui la geometria vive!'* and even in the 1950's, when Bianchi's *Opere* were published, geometers still considered it one of the most important discoveries of his career and the culmination of the golden age of classical differential geometry. It is very sad that the undergraduate student hears today only of Bianchi's identities (appearing in a four page article; it seems that Ricci had previously discovered them while doing computations for his newly discovered absolute calculus of tensors contraption, but he did not see them important so as to publish them): G. Fubini, in his eulogy appearing in part 1 of Bianchi ([5],Vol **1**) mentions Bianchi's identities only in a sentence on page 60, crammed among four other results (including the list of spaces from Thurston's geometrization program), but mentions in a consistent footnote on page 70 a beautiful theorem where Bianchi described all normal W congruences by putting an Euclidean configuration into motion and under certain tangency requirements (so basically *integrable systems* means nothing more and nothing less than *Euclid in motion*; as to what Euclidean configurations are chosen to behave well in the moving picture is decided by *Archimedes' method of integration*).

There are tree main theorems in the theory of deformations of quadrics, all due to Bianchi (see (122,§ I-XII)):

-the existence and inversion of the B transformation of quadrics and the ACPIA:

Theorem I

Every surface $x^0 \subset \mathbb{C}^3$ applicable to a surface $x_0^0 \subseteq x_0$ (x_0 being a quadric) appears as a focal surface of a 2-dimensional family of W congruences, whose other focal surfaces $x^1 = B_z(x^0)$ are applicable, via the Ivory affinity, to surfaces x_0^1 in the same quadric x_0. The determination of these surfaces requires the integration of a family of Riccatti equations depending on the parameter z. Moreover, if we compose the inverse of the RMPIA with the rolling of x_0^0 on x^0, then we obtain the rolling of x_0^1 on x^1 and x^0 reveals itself as a B_z transform of x^1;

-the Bianchi Permutability Theorem:

Theorem II

If $x^1 = B_{z_1}(x^0),\ x^2 = B_{z_2}(x^0)$, then one can find only by algebraic computations a surface $B_{z_2}(x^1) = x^3 = B_{z_1}(x^2)$; thus $B_{z_2} \circ B_{z_1} = B_{z_1} \circ B_{z_2}$ and once all B transforms of the seed x^0 are found, the B transformation can be iterated using only algebraic computations;

-the Hazzidakis transformation:

Theorem VIII

If a surface $x^0 \subset \mathbb{C}^3$ is applicable to a surface $x_0^0 \subseteq x_0$ and the homography H of Bianchi II takes the confocal family x_z to another confocal family $\tilde{x}_{\tilde{z}},\ \tilde{z} = \tilde{z}(z),\ \tilde{z}(0) = 0$, then one infinitesimally knows a surface $\tilde{x}^0 = H(x^0)$, called the Hazzidakis (H) transform of x^0 and applicable to a surface $\tilde{x}_0^0 \subseteq \tilde{x}_0$. Moreover the H transformation commutes with the B transformation ($H \circ B_z = B_{\tilde{z}} \circ H$) and the $B_{\tilde{z}}(\tilde{x}^0)$ transforms can be algebraically recovered from the knowledge of $\tilde{x}^0$ and $B_z(x^0)$.

If we apply Theorem VIII to QC of revolution around the focal axis, then this induces some transformation of CMC surfaces described by the foci (Guichard). This transformation was discovered earlier by Hazzidakis and is the reason why Bianchi still calls the transformation for general quadrics Hazzidakis. The current denomination for Hazzidakis type (algebraic) transformations of local solutions of same or different integrable systems is *Miura* transformations, as Miura discovered a transformation between local solutions of the KdV equation and local solutions of the *modified* KdV (mKdV) equation (see Roger-Schieff ([26]).

An important tool in proving Theorem VIII is the notion of *surfaces conjugate in deformation*: pairs of non-flat non-homothetic surfaces on which the asymptotic coordinates and all *virtual asymptotic coordinates* (coordinates which can become asymptotic on an applicable surface) correspond. Thus for any surface applicable on the first one one infinitesimally knows a surface applicable to the other. An immediate consequence of the Codazzi-Mainardi equations, the change of Christoffel symbols for the change of coordinates and the equation of geodesics is the fact that 'conjugate in deformation' is equivalent to correspondence of asymptotic coordinates and geodesics (and we know that this is realized by the homography of Bianchi II); conversely Bianchi in ([5],Vol **5**,(87),(90))

and Servant in [28] proved that only deformations of quadrics can appear as surfaces conjugate in deformation (see also Darboux ([13],§ 603)); Bianchi proved the statement for quadrics of revolution and Servant for the remaining quadrics.

The notion of surfaces conjugate in deformation has another important application (Bianchi (122,§ XI,§ 56)): if we consider x_z for a particular value of z as the Cayley's absolute, then the Euclidean confocal family of x_0 remains confocal family in this new geometry. Lines become geodesics (so remain lines) and all properties of confocal quadrics are preserved; in particular Euclidean geodesics and asymptotic coordinates on x_0 remain geodesics and asymptotic coordinates on x_0 in the new geometry, so all virtual asymptotic coordinates on x_0 in the new geometry still correspond to virtual asymptotic coordinates of x_0 in the initial Euclidean geometry. Thus deformations of quadrics in other geometries are reduced to deformations of quadrics in the Euclidean space.

With this construction the deformation of all quadrics in $\mathbf{O}_3(\mathbb{C})$ except the flat Clifford torus can be reduced to the deformation of quadrics in $\mathbb{C}^3$; all flat surfaces in $\mathbf{O}_3(\mathbb{C})$ appear as rotations of rollings of flat surfaces in $\mathbb{C}^3$.

Darboux split the prize of the French Academy between Bianchi and Guichard for solving the problem for quadrics (Guichard had also other results, including some transformations G of deformations of general quadrics) and asked if there are any relationships between the transformations found by Bianchi and Guichard. Bianchi showed that certain G transformations are compositions of two B transformations of different ruling families $B_z \circ B'_z$ with finite z; Calapso showed in 1912 that the remaining G transformations are compositions of two B transformations $B_\infty \circ B'_\infty$ (see [11]). Thus most transformations of deformations of quadrics were reduced to compositions of B and H transformations. The problem was solved in the sense that solutions depending on arbitrarily many constants have been found: beginning with a seed surface we integrate a 1-dimensional family of Ricatti equations; after that only algebraic computations are required to produce solutions (BPT). Bianchi communicated to Calapso (see [11]) that the B_∞ transforms are generated by some trivial infinitesimal deformations (isotropic translations); therefore we can find them by quadratures.

It thus becomes apparent that certain metric-projective properties of confocal quadrics (most of them established in the first half of the XIX$^{\text{th}}$ century) *carry out* (stick and transfer) by rolling to and influence deformations of quadrics and surfaces geometrically linked to these, therefore providing a wealth of integrable systems and projective transformations of their solutions (the B transformation boils down to a Ricatti equation, the cross-ratio of whose four solutions is constant), directly linked to quadrics or closely related to these: see Moser [24] for similarities between quadrics and other integrable systems, Bianchi ([5],Vol **4**,(108)) or Burstall [9] and collaborators for isothermic surfaces, Bianchi surfaces (first studied, according to Bianchi ([4],§ 294-§ 295), by Peterson), Rogers-Schieff [26] for Ţiţeica surfaces, etc. For example the sine-Gordon equation (with its many real forms) describes many types of surfaces (with positive definite metric or not) in spaces of constant curvature (with positive definite metric or not) and all these are generated just by a particular type of confocal quadrics: the complex spheres. In particular the deformation of the (pseudo-)sphere is equivalent to the deformation of Darboux quadrics (quadrics tangent at one point to $C(\infty)$; they cannot be realized as quadrics in $\mathbb{R}^3$, although their deformations can: see Darboux [15]), of paraboloids and of quadrics of revolution (see Eisenhart ([20],§ 144)); deformations of general quadrics are put in relation with special coordinates on the (pseudo-)sphere (Eisenhart [20]).

There is literature of deformations of 2-dimensional quadrics in higher dimensional spaces including up to the 1930's, due mostly to Calapso and Guichard.

Cartan (see [12]) studied in 1918 deformations of space forms in space forms; in particular he proved that there are no deformations of the hyperbolic space form $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{n+p}$, $p < n-1$ and all deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ appear only with flat normal bundle. In 1972 Moore (see [32]) introduced generalized Tchebyshev coordinates on deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$. Upon a suggestion of S.S. Chern and using Cartan-Moore, Tenenblat and Terng (see [32]) developed in 1980 the B transformation for deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ (and Terng developed in [33] the BPT for such a B transformation). Note that a simple corollary of the ABL method is the

$\mathbf{O}_{n-1}(\mathbb{R}) \times \mathbf{O}_{n-1}(\mathbb{R})$ symmetry of the isoclinic facets for the Tenenblat-Terng B transformation of deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ (the rolling can be realized with the symmetry $\mathbf{O}_{n-1}(\mathbb{R})$ of the normal bundle and one takes into consideration the symmetry of the tangency configuration). More recently Tenenblat, Terng, Uhlenbeck and collaborators (see [36] and its references for earlier results and names) have developed in particular the B transformation of space forms (with flat normal bundle) in space forms and in general have adapted a machinery, Lie algebraic in character, to deal with such B and Darboux transformations (different from the D transformation of isothermic surfaces) for integrable systems, obtaining for example variants of the Bianchi Permutability Theorem and n-soliton formulae. Bobenko and Pinkall have developed in 1996 (see [6],[7]) a theory of discrete isothermic and CGC -1 surfaces, at the heart of which lies the BPT and Bianchi's notion of Möbius configurations (composed with a Lie contact transformation for discrete isothermic surfaces). While Bianchi developed in ([5],Vol **5**,(117)) moving Möbius configurations for *Bäcklund* (B) transformations of general surfaces (the two focal surfaces of a W congruence are considered B transforms one of the other), he was not aware of the fact that pieces of discrete surfaces are subsets of Möbius configurations. Probably discrete mathematics (present since the wise ancient's times; it is by discrete mathematics that Apollonius and Kepler completed their computations) was not seen important, as there were no computers and much need for it.

Since most of Cartan's arguments are projective, the natural question arises if a rich family of deformations in $\mathbb{R}^{2n-1}$ of space-like regions in n-dimensional quadrics exists. Such regions can appear as the ones in n-dimensional quadrics of the Lorentz space of signature $(n, 1)$ along which the normal is time-like (thus this linear element is incomplete except for the pseudo-sphere, since there are isotropic normal directions) or as quadrics in $\mathbb{R}^{n+1}$. Berger, Bryant and Griffiths provided an affirmative answer in 1983 (see [3]); in fact such regions of quadrics are the only n-dimensional Riemannian manifolds whose family of deformations in $\mathbb{R}^{2n-1}$ is as rich as possible: it depends on $n(n-1)$ functions of one variable. Once this question was settled, the natural problem of generalizing Tenenblat-Terng's B transformation to such deformations appears. Note that the $\mathbf{O}_{n-1}(\mathbb{R}) \times \mathbf{O}_{n-1}(\mathbb{R})$ symmetry deduced from the ABL method remains true for general quadrics, since it has to do with the normal bundle. One can study a more general question by allowing the linear element of the quadric to be indefinite, every component of its signature being of course smaller than or equal to the corresponding component of the signature of the ambient Lorentz space of dimension $2n-1$, but once the issues are clarified for a general quadric all other cases should follow immediately.

1.4. **Statement of results.**
First a short history of the theory of deformations of quadrics from my point of view; since at this point I have a vague memory about its early part, the only sources are the Library's stamped due dates (assumed to be roughly half an year after the day books were borrowed), the talks I gave and some of my emails. I became aware in early Fall 2002 of Eisenhart [19]; except for the first and last chapters and most of the exercises (which are sort of a short list of important facts from Bianchi [4] and Darboux [13]) it was in general accessible. By early Spring 2003 (after the qualifier) I also found Eisenhart [20] and the theory of deformations of quadrics began to loom large, as it always appeared in the last chapters and thus clearly beyond my level of expertise. The find of Eisenhart [20] brought to my attention isothermic surfaces (and their *Darboux* (D) transformation) and W congruences (and their B transformation) as important objects of investigation, but that book is very technical and does not allow a discontinuous reading (it builds an impressive apparatus from the beginning and provides precise proofs later just by sending the reader to previously proved results; by the time the reader reads the applications the impressive apparatus is already forgotten): after a serious reading of the first chapters I just jumped to a cursory reading of the remaining ones and from the theory of deformations of quadrics I was not able to understand much. During this time I was also involved in studying some current literature, but the current techniques and tools clearly appeared insufficient (I was involved in a parallel reading of the classical literature and the current one up to Summer 2005, after which time I mostly dropped the later from my readings).

By mid Spring 2003 I also found Darboux ([13], Vol **2**) and I realized that it is much cheaper to read Darboux than to solve the problems of Eisenhart [19], at least until one gets used with the techniques (whole sections of Darboux's are compressed as exercises in Eisenhart). In late Spring 2003 the article Bianchi ([5],Vol **4**,(108)) was brought to my attention. Since at that time I was involved in a dual classical-current literature reading, I searched on the same afternoon *'isothermic surfaces'* on Altavista (for some time I use Google). In the internet age the best research is done by use of search machines on the internet; of course there are also leads which *apparently* have less to do with mathematics, but in the first 20 entries one also finds the relevant literature (this is how I found Moser [24] and later Arnold [2]). Today it is very difficult to find Burstall [9] among the first hits, mainly due to the size of the literature on isothermic surfaces. After a cursory reading of Bianchi ([5],Vol **4**,(108)) and the last section of Burstall [9], I realized again the importance of the theory of deformations of quadrics; Bianchi seemed to provide in ([5],Vol **4**,(108,§ 15)) an approach simpler than Eisenhart's and also the correspondence between the D transformation of *special isothermic surfaces* (the closure of *constant mean curvature* (CMC) surfaces under conformal transformations of $\mathbb{C}^3$) and deformations of quadrics provided the necessary bridge to show that quadrics put their hands in a lot of cookie jars. The most successful expedition in the Library turned out to be when I searched Bianchi as author and I found his *Opere*; on the same day I grabbed a few of his volumes (including the two parts Vol **4** concerning the theory of deformations of quadrics), Vol **2** of Bianchi [5] and Vol **3,4** of Darboux [13]. I do not believe there has been more booty in an expedition, except probably when the Crusaders first broke Constantinopole's walls, bringing economic decline and thus easying the rise of the Ottoman Empire and the fall of but the last remnants of the Eastern Roman Empire two hundred years later . It is on the same day that I read and understood Bianchi's theorems of the theory of deformations of quadrics; understanding their proofs would take me one more year and repeated unsuccessful attempts of scaling the walls of the fortification (I was still trying at the time to use current tools in finding alternative simpler proofs or new results, but I was never leaving the infinitesimal neighborhood of the seed; in fact my purpose at that time was to provide an explanation according to the current tools (loop groups, solitons, etc) of the theory of deformation of quadrics; as I have delved deeper into the classical geometers I found other interesting things to do). Since Guichard's result appeared to be easier in Bianchi's interpretation than in that of Eisenhart [20], by late Fall 2003 I had a good understanding of it; in Spring 2004 I decided to seriously read Weingarten's method (one of the last chapters of Eisenhart [19]), since it involved a particular case of deformations of quadrics. At that time I already had a good understanding of the rolling problem (including the trick with infinitesimal deformations of the rotation of the rolling generating the corresponding pairs of applicable surfaces) and a small result of cyclic systems in space forms. I distinctly remember considering the next point of view on the rolling and the deformation problem: imagine a piece of paper; one deforms it while preserving a virtual copy of the initial planar position such that a given planar curve remains planar and nonsingular in the initial plane (the crumpled piece of paper is allowed to develop singularities outside a small tubular infinitesimal neighborhood of the curve); the same trick but in higher dimensions would be used in the deformation problem. However too many variables were introduced and it seemed to me to always boil down to the same difference of the two GCM equations (again I was never leaving the infinitesimal neighborhood of the seed, but the difference of the two GCM equations got a simple geometric explanation). And even understanding deformations of sheets of paper is still an open question: I have a vague memory of a talk where a sheet of paper was deformed while keeping a linear boundary fixed (two strong opposite laws): *parabolas* of singularities developed near the linear boundary and in a *fractal* (fractured) pattern, so I can honestly say that *deformation into parabolic patterns* naturally appears in mathematics as in every facet of reality and nature; when such a pattern appears it usually comes in abundance and singular manner (thus finite energies are discharged and distributed on infinitesimal domains creating ∞ multiplicity), so as to equilibrate the strong opposing forces without breaking the usual smooth rules (the smooth part of the problem notices nothing interesting, but according to Archimedes the simplest explanation possible and thus

the truth lies in the singular behavior; when in doubt nature always *runs home to mama* and chooses quadrics in order to simplify and expedite the non-linear problem). There is a famous joke with a famous current mathematician stating that he studies quadratic equations (to the dismay of the inquirer): it is in studying iterations of quadratic functions over complex numbers that the most famous fractal patterns naturally appear.

Being involved with teaching a course in Summer 2004, I cut down on research and I concentrated instead on reading Bianchi's proofs, which by that time I almost understood; as soon as I went through his proofs I decided to write a set of notes to myself. Still due to that Summer 2004 there is a *'The Tonight Show'* show of Jay Leno's to whom I may have thank for my current research. So Jay Leno decided one night to invite a guest which was a Professor of Mathematics; that fella came riding a car with square wheels; of course it was a bumpy ride, but the car was lifted and fitted on a special road: on that road the car had a smooth ride. The road looked like a cycloid (I had just covered the cycloid in the Calculus II course I was teaching at the time); thus I wanted to see if that is true; it turns out that it uses the most difficult and unnatural to remember integral formula for trigonometric functions, namely $\int \sec(s)ds = \ln(\sec(s)+\tan(s))+\mathcal{C}$, because the road is given by $(\ln(\sec(s)+\tan(s)), -\sec(s)),\ 0 \leq s \leq \frac{\pi}{4}$ and then it extends by reflections in the mirrors $x = 0, x = \ln(1+\sqrt{2})$. So after the fact one can easily see that the road cannot be the cycloid, as it turns at 90° (as it should) instead of at 180°, but doing the computations and a Maple moving picture (I applied the rolling I already knew to curves) helped me later immediately do the Maple moving picture for the elliptical wheel. While preparing the final exam for the course taught in Summer 2004 and working in the same time on the notes I noticed the first simplification in Bianchi's proofs, which was something of the form $du_1 = u_{1u_0}du_0 + u_{1v_0}dv_0 + u_{1v_1}dv_1$ for $u_1 = u_1(u_0, v_0, v_1)$; so Bianchi was doing computations with the *right hand side* (rhs) and I did them with the *left hand side* (lhs) (the sinister side of the equation; also I thought at some point that Bianchi was a member of numerous Academies having to do with silviculture, since *forestiere* can be interpreted not only as foreign, but also as something having to do with a forest; thus *foreign* has a Latin root, a fact previously unknown to me; having good knowledge of Romanian and none of Italian I realized however the correct use of *forestiere*, as opposed to *nostrane*; not to talk about *concorso banditi* being probably an *admission contest* in a respected elite group, such as *Scuola Normale Superiore*). As a consequence by late August 2004 I saw the relations (3.12) and the one line proof of the ACPIA and I decided to stay on the theory of deformations of quadrics, which I did since then. I remember being very optimistic in Fall 2004 about finishing a simplification of Bianchi's computations and thus having the correct tools at hand to attack the higher dimensional problem (by that time I already found Bobenko-Pinkall [6],[7] and Tenenblat-Terng [32]). It is then that I realized that by considering imaginaries one can put together all QC in a single case (Bianchi was using parametrization by trigonometric functions for real ellipsoids and hyperboloids with two sheets). However, I got bogged down in a project of proving the algebraic computations of the BPT for QC (3.3); after completing the difficult computations and going further to prove the cross-ratio property I realized that the cross-ratio property can be used to simplify the algebraic computations of the BPT. By Spring 2005 I headed to the H transformation; thus I followed Bianchi's example of a Darboux quadric and realized that all quadrics can be treated in the tree cases QC, QWC and IQWC and that the last two were similar. In late Spring 2005 in trying to discuss the BPT for the B'_z transformation I had to redo all computations for the other ruling family: it is this how I found the simple geometric proofs of the necessary algebraic relations (3.12) (the simple geometric interpretations were clear enough from Summer 2004, but their proof was not). Then I have jumped however to the most difficult part of the algebraic discussion of the static picture in higher dimensions (namely Bianchi II) without having the SJ canonical form; just as in the Romanian *Mănăstirea Argeşului* story what was built one day was destroyed over night. It is then that I realized the need of the SJ canonical form, since even the proofs of the other classical theorems about confocal quadrics would succumb if $\sqrt{I_n - zA}$ does not commute with A (this is an obvious statement for z close to 0 but even the existence, definition and good name of $\sqrt{I_n - zA}$

is questionable for z away from 0). Few examples in dimension $n = 4, 5$ showed the proper attack strategy: a global one, since the eigenvalue-by-eigenvalue strategy for non-isotropic eigenvalues was the story with destroying what was previously built. Once the attack strategy established everything fell into place and all possible information was squeezed and completely used (as expected). With the SJ canonical form in place the proof of Bianchi II was amenable to attack, but again false leads took their toll (even at this point there is the question of a diagonalization process). By the end of Fall 2005 the proof of the Möbius configuration $\mathcal{M}_3$ was attacked; as a consequence the simpler notation of the Möbius configuration $\mathcal{M}_2$ revealed itself and consequently the algebraic computations of the Möbius configurations $\mathcal{M}_n$ also immediately revealed themselves (there is still the question of validating those formulae). For the most degenerate quadric from a metric point of view IQWC (3.9) these computations boil down to a determinant formulation, a mixture of Vandermonde determinant and homographies (separate linearity in rulings); from this example one can deduce the invariant algebraic formula for $\mathcal{M}_n$ and assume that it is true for all other quadrics. With use of Menelaus's theorem a simple geometric argument for all cases is immediate. Still due to the identities involved in the argument with Menelaus's theorem a proof of the fact that the cross-ratio property is the consequence of the second iterated tangency configuration was immediate; as a consequence what up to that point was a non-rigorous argument on top of which a complicated identity with $u_1 := \infty$ still had to be checked became rigorous and the checking became unnecessary. All specific algebraic computations were done with Maple; consequently I knew I was on the wrong path (or I was using the wrong notation) when my computer was crunching numbers and did not even want to shut down when I was pressing the corresponding button and I knew I was on the correct one when the computer was polite enough to confirm the result in a few seconds for $n = 3$ and up to 10 minutes for $n = 6$ for QC (3.3); moreover the computer clearly showed me many tricks on many other occasions and verified most of algebraic identities appearing in these notes. In Spring 2006 with the algebraic computations of the higher dimensional static picture completed (of course it was and is still missing the algebraic computations specifically needed for the theory of deformations of quadrics, namely those obtained by the ABL method) I decided to investigate Cartan [12]; up to that point it was unclear to me if for example $\mathbf{H}^n(\mathbb{R})$ can appear as a sub-manifold of $\mathbb{C}^{n+p}$ with $p < n-1$ other than the pseudo-sphere itself. It is after seeing Cartan's mostly projective arguments that I realized that the dimension $2n-1$ cannot be lowered even if one relaxes the restriction on not using imaginaries. The strategy is clear: just like Lie find the leaves when the seed $\mathbf{H}^n(\mathbb{R})$ actually coincides with the actual pseudo-sphere in $\mathbb{C}^{n+1} \subset \mathbb{C}^{2n-1}$, then just like Bianchi take their metric-projective generalization (thus the leaves must be cones in the tangent cone including possibly the most degenerate case of lines) and in trying to prove the higher dimensional theory of deformation of quadrics the necessary algebraic identities of the static picture will reveal themselves. However, just like the sine-Gordon equation is not best suited for the Lie ansatz (I was calling it at that time approach), similarly the Tenenblat-Terng approach is not best suited for the Lie ansatz since it assumes real seed: one gets only 1-solitons similarly to getting 1-solitons from the axis of the tractrix for CGC -1 surfaces. The find of the Berger-Bryant-Griffiths [3] article not only confirmed the program, but also removed its most difficult part (I should mention that this article was first mentioned to me in a course on the isometric embedding problem in Fall 2004, but I did not hear the word *quadric* on that occasion, just the fact that it deals with some difficult particular cases where the dimension of the surrounding space is below $\frac{n(n+1)}{2}$). However any hope of understanding that article without serious prerequisites of Algebraic Geometry is unfounded in my opinion (*'I lost it, Lou!'* when I was hit with that Serre's stuff taken for granted), so a simpler proof of the weaker part (the program I needed and intended to prove) that n-dimensional totally real quadrics admit rich families of deformations in $2n-1$ Lorentz spaces of corresponding signatures and with flat normal bundles should be provided first (the use of elliptic coordinates on the quadric and lines of curvatures on its deformation should render it a perfect example of Cartan's calculus, moving orthonormal frames and counting certain numbers to provide solutions of the corresponding differential system and the dimensionality of the solutions space).

Just like Terng's generalized sine-Gordon equation, use of proper coordinates should reduce the problem of existence of seeds and of the B transformation to a Frobenius (complete) integrability, thus avoiding the use of the more complicated Cartan-Kähler theorem. The Ivory affinity still provides the applicability correspondence and the $n, 2n-1 = (n-1)+1+(n-1)$ dimensions fit perfectly to uniquely provide in general the rigid motion taking a facet centered on the confocal quadric (but not tangent to it; it even in general lives outside the space $\mathbb{C}^{n+1}$ where the quadric lives, so the easy generalization of the RMPIA to higher dimensions is not the correct one) to the tangent facet corresponding by the Ivory affinity and does the reverse trick to the other pair of facets. The discussion of totally real cases should close the 2-dimensional discussion in Euclidean surrounding space, but again the stronger statement that the only surfaces in quadrics and having real linear element are totally real took its toll and the discussion is incomplete (only the most degenerate cases are completed; I believe there is a simpler unifying approach using rolling, but that still eludes me at this point). So this was the setting when I found Archimedes' opinion on the matter: this provided further insight into the higher dimensional discussion and a method to find *all* necessary algebraic identities needed for the moving picture from the static picture, without actually trying to complete the computations of the moving picture in order to see what are those algebraic relations. Once those algebraic relations are found again their geometric proof should be easiest.

In § 2 we are interested in bringing a quadric to the canonical form (depending on as few constants as possible), finding its confocal family and proving the classical theorems (only Bianchi II presents some challenge). The computations are on occasion lengthy, although they always require only elementary linear algebra notions. While part of these results with proofs are present in the quoted bibliography, we believe that the unified complete treatment appears here for the first time. For example Bianchi in ([5],Vol **5**,(90)) proves Bianchi II with $n > 3$ only for diagonal QC; also although most (if not all) of the totally real quadrics of $\mathbb{C}^3$ were known to the classical geometers, we have not found a reference with an exhaustive and organized discussion.

First we bring a symmetric complex matrix, via conjugation by a rotation $R \in \mathbf{O}_n(\mathbb{C})$, to a form depending on as few as possible constants (SJ form); it inherits the properties of matrices in Jordan canonical form. Such a SJ matrix is formed by SJ diagonal blocks $aI_p + J_p,\ p \geq 1,\ J_1 := 0 = 0_{1,1}$; the SJ blocks $J_p,\ p \geq 2$ are formed by augmenting either (according to p being either even or odd) of the blocks $J_2 := \frac{1}{2}\begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix},\ J_3 := \frac{1}{\sqrt{2}}\begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & -i \\ 1 & -i & 0 \end{bmatrix}$, diagonally on the upper left and until the dimension p is reached, with blocks $\begin{bmatrix} 0_{2,2} & M \\ M^T & 0_{2,2} \end{bmatrix},\ M := \frac{1}{2}\begin{bmatrix} 1 & i \\ -i & 1 \end{bmatrix}$ (the right lower $0_{2,2}$ block is superimposed by addition on the left upper 2×2 block of J_p in order to get J_{p+2}). The normalization $f_1 := \frac{e_1 - ie_2}{\sqrt{2}}$ instead of $f_1 := e_1 - ie_2$ for the first standard isotropic vector (the $e_j,\ j = 1, ..., n$ are the standard basis: $e_j^T e_k = \delta_{j,k}$) is preferred in order to avoid cumbersome powers of 2 when working with SJ blocks.

Although a SJ matrix as defined above efficiently encodes all metric invariants of a symmetric complex matrix A (its eigenvalues λ and orders of cyclic vectors of $A - \lambda I_n$), we prefer when convenient to work with the *SJ type* (the block J_p is replaced with a polynomial $P(J_p)$ in J_p with $P'(0) \neq 0$), since it is closed under needed algebraic operations; the stronger SJ notion would require a rotation after each such algebraic operation.

For $z = \infty$ or inverses of non-zero eigenvalues of A we have singular quadrics of the confocal family; the singular set $\mathcal{S}(x_z) \subset x_z$ for x_z singular is a lower dimensional quadric. These singular sets play an important rôle in the proof of Bianchi II, in the course of which we provide a classification of $\mathbf{PGL}_n(\mathbb{C})$ suited to it.

We close § 2 with a parametrization of confocal IQWC (this allows us to consider them as metrically degenerated QWC and thus to treat later all (I)QWC as a single case) and with a discussion of totally real quadrics.

In § 3 we restrict our attention to quadrics in $\mathbb{C}^3$; in § 3.1 we present a list of all confocal quadrics in $\mathbb{C}^3$, their ruling families parametrization invariant under the Ivory affinity and homographies of Bianchi II. Bianchi in ([4],§ 425) and (122,§ 51-§ 57) uses only the homographies which take QC (3.3) to QC (3.3) and QC (3.4) to QWC (3.6), although the general geometric method from ([5],Vol **5**,(90)) applies to all cases.

In § 3.2 we discuss totally real quadrics; the part of the argument where one proves that a surface in a quadric and having real (valued) linear element must be a totally real quadric may be replaced in § 5.6 with a simpler unifying argument using rolling.

While currently it is well known that there is a single type of quadrics from a complex projective point of view, two types of quadrics from a complex affine point of view (namely with or without center), two types of real quadrics from a real projective point of view (namely ruled or not ruled), 5 types of real quadrics from a real affine point of view (namely three with center: connected ruled, connected not ruled and disconnected not ruled; two without center: ruled and not ruled) and 12 types of real quadrics from a real metric point of view (all subcases of the previous 5), it is less known since the classical times that there are 11 types of quadrics from a complex metric point of view (all subcases of the 7 cases QC (3.3)-(3.5), QWC(3.6)-(3.7), IQWC (3.8)-(3.9)) and numerous types of totally real quadrics. The number of types of totally real quadrics is even since multiplication by i changes the signature, both of the linear element of the totally real surface and of the ambient totally real space, so it is enough to count the 12 types of real quadrics plus the types of quadrics in $\mathbb{R}^2 \times (i\mathbb{R})$ and double this number.

A simple heuristic argument in favor of counting all types of totally real linear elements of a complex metric type which is not a complex sphere is to double the number of components of singular quadrics of the confocal family (including the singular quadric at infinity), since each type of totally real linear element comes with its totally real confocal family, which contains singular linear elements. The change in the type of linear element occurs along the singular quadric; thus the change of the type of linear element due to the multiplication by i occurs at infinity. Since most families of confocal quadrics are generated by finite conics c (the family being the adjugate of the pencil generated by the adjugate of c and the adjugate of $C(\infty)$), it is expected that all information about such a family is recorded by the pair $(c, C(\infty))$.

Take for example the real confocal family of a general ellipsoid; the change in the real confocal family from ellipsoids to hyperboloids with one sheet occurs along a real singular quadric of the real family: a mostly doubly covered real principal plane. In that real singular quadric the singular part (covered once) is an ellipse; it separates the interior convex region (parameterized by complex conjugate parameters and thus having imaginary rulings) from the exterior concave region (parameterized by real parameters and thus having real rulings); the two parametrizations coincide on the ellipse. Therefore the ellipsoids of the real confocal family admit parametrization by complex conjugate parameters and have imaginary rulings, while the hyperboloids with one sheet of the real confocal family admit parametrization by real parameters and have real rulings.

Note that not only the types of totally real quadrics are distinct (one cannot be obtained from another by means of a complex rigid motion), but also their linear elements are distinct, since two quadrics with same linear element can be rolled one onto the other and the rolling with the determinant of its rotation being 1 must be a rigid motion. Thus all real symmetries of the linear element of a totally real quadric are in correspondence with rigid motions preserving the quadric; the complex symmetries are accounted by changing the type of the totally real quadric within the same complex metric type of quadric and they must preserve the character of real or imaginary rulings.

Since the B transformation is essentially projective, we must use projective coordinates (either asymptotic or conjugate systems) either on x_0^0 or on x^0 (the sine-Gordon equation $\omega_{uv} = \sin\omega$ for deformations of the pseudo-sphere is obtained by considering the net of arc-length asymptotes on x^0). Due to the essential rôle played by the Ivory affinity in the B transformation of quadrics, the use of rulings on x_0 (asymptotic coordinates on x_0^0) is more appropriate.

Beginning with § 3.3 we present the algebraic identities necessary for the theory of deformations of quadrics. While they and their proof are simple and on occasion elementary, the natural character of these relations is definitely not obvious; these relations and their necessity become obvious once certain natural ansatzs are made in the moving part (such as the ABL method, the ACPIA or ruled surfaces for an a-priori valid BPT).

The algebraic identities (3.12), their equivalency to the TC and their simple geometric interpretation do not appear in Bianchi's account of deformations of quadrics, although he uses equivalent algebraic identities (for example (29) in (122,§ 4)) and the place of use is the same: bring the differential system of the B transformation to a Ricatti equation, check the complete integrability of this Ricatti equation and prove the ACPIA. Bianchi's approach to the ACPIA in (122,§ 7,§ 8,§ 9) is to take dv_1 from the (5.3), du_1 from (3.26) and to show that $|dx^1|^2-|dx^1_z|_{v_1=\text{const}}|^2 = |dx^1_0|^2-|dx^1_z|_{v_1=\text{const}}|^2$ as symmetric quadratic forms in $du_0,\ dv_0$; in (122,§ 31,§ 32) he takes du_1 from (3.26), imposes the ACPIA and obtains the Ricatti equation (5.3) (in both instances he uses the Gauß theorem to cancel the influence of the shape of x^0 with the influence of the shape of x^0_0). Bianchi's choice to prove $|dx^1|^2-|dx^1_z|_{v_1=\text{const}}|^2 = |dx^1_0|^2-|dx^1_z|_{v_1=\text{const}}|^2$ instead of just $|dx^1|^2 = |dx^1_0|^2$ becomes clear for the following reasons:

(I) For a function $f = f(u_0, v_0, v_1(u_0, v_0)),\ du_0 \wedge dv_0 \neq 0$ he considered for example the explicit derivative $\partial_{u_0} f = \frac{\partial f}{\partial u_0}$ of f wrt u_0 as opposed to the implicit derivative $\partial_{v_1} f v_{1u_0}$ of f wrt u_0; the total derivative is $f_{u_0} = \frac{df}{du_0} = \partial_{u_0} f + \partial_{v_1} f v_{1u_0} = f_{u_0}|_{v_1=\text{const}} + \partial_{v_1} f v_{1u_0}$.

(II) When x^0 coincides with x^0_0 the leaves of the considered integrable distribution are the rulings $v_1 = \text{const}$, so all formulae already found for x^1 in general position apply to x^1_z by considering $v_1 = \text{const}$ and replacing the shape of x^0 with the shape of x^0_0.

To simplify the notation we shall use f_u with the meaning of $\partial_u f$ when the explicit dependence of f on u is clear (other variables appearing in the definition of f do not depend on u); for example in the above situation we assume first $du_0 \wedge dv_0 \wedge dv_1 \neq 0$, derive $df = f_{u_0} du_0 + f_{v_0} dv_0 + f_{v_1} dv_1$ and after restricting to a surface $v_1 = v_1(u_0, v_0)$ we have further $dv_1 = v_{1u_0} du_0 + v_{1v_0} dv_0$.

We propose (in § 5) a different approach: show that $|dx^1|^2$ and $|dx^1_0|^2$ are equal as symmetric quadratic forms in $du_1,\ dv_1$; in doing so the computations are reduced almost to a tautology and the necessary relations (3.12) naturally appear from an analytic point of view; they will be consistently used throughout the remaining sections. Note that it is very likely that Bianchi saw the simpler proof using only $du_1,\ dv_1$ and without expanding these as 1-forms in $du_0,\ dv_0$, since the needed rudiments of Cartan's exterior calculus and of Ricci's absolute differential calculus of tensors were present at the time for almost a decade, but he probably felt more secure in providing an explanation according to the standards of proof of the time; it took another decade for Ricci's absolute differential calculus of tensors and several more decades for Cartan's exterior calculus to become mainstream mathematics.

The simplest proof of the ACPIA is not almost a tautology, but an actual tautology; it is essentially due to Bianchi, although he preferred the security of an analytic confirmation to the power of his geometric arguments. To prove the inversion of the B transformation in (122,§ 11), Bianchi defines the rolling of x^1_0 on x^1 as the composition of the RMPIA with the rolling of x^0_0 on x^0; he also checks that the v_1 ruling goes to the correct place (if a family of rigid motions takes a surface and its tangent spaces to another surface and its tangent spaces, it is not a-priori a rolling, as it may rotate the tangent spaces into themselves). In doing so he used the ACPIA and found a simple formula for the rolling x^1_0 on x^1, but he actually found the simplest proof for the ACPIA, since existence of rolling implies applicability correspondence (one needs only check the complete integrability of the Ricatti equation (5.3), since we need the leaf x^1 to exist).

While Bianchi knew in (122,§ 11) that by changing the ruling family on x^1_z the rotation of the RMPIA must acquire -1 determinant (and thus must be a rotation of determinant 1 composed with a reflection), he was not aware of the simple observation that if the required reflection takes place in the tangent space of x^0_0, then the rotation does not otherwise change at all (to change the rulings $u_0 \leftrightarrow v_0$ he uses instead a reflection r_{e_2} in the $e_2^T x = 0$ plane for x^0_0 being QWC

(3.6)). In fact initially we were also not aware of this observation and thus provided an analytic confirmation of (3.12); it is only after considering the algebraic computations of the BPT that we realized that a good accounting of all RMPIA's for all rulings is needed. Using this observation a simple geometric proof of (3.12) is immediate; as a result we have gained much respect for what Bianchi calls *'elementary properties of the Ivory affinity'* and consistent use of symmetries is carried through the remaining sections*. We introduce four quadratic quantities $\Delta^{\pm}$, $\Delta'^{\pm}$ which are roughly the angles between the normal fields m_0^1, $m_0'^1$ of the considered distributions and the rulings of x_0 at x_0^0 with the denominator parts removed; the normal fields m_0^1, $m_0'^1$ themselves have the usual definition (involving a cross product) with the denominator part removed; further we derive some useful algebraic identities. Note that these analytic computations are not necessary for the B transformation for finite z, but we need them for the case $z = \infty$, since in that case the geometric picture disappears and we can only use analytic computations.

Bianchi derives the properties of the *second iteration of the tangency configuration* (SITC) and proves the BPT for ruled deformations of QWC (3.6) in (122,§ 21-§ 25); as a corollary he proves in (122,§ 26) the beautiful cross-ratio property (the BPT is essentially a cross-ratio theorem; in fact it roughly boils down to the product of a cross-ratio property and same cross-ratio property composed with a simple symmetry). We propose a change in Bianchi's approach: consider the cross ratio property from the beginning, to be proven together with the other algebraic properties of the SITC; this interlacing simplifies the computations of the SITC (we need this simplification especially for QC (3.3), where even with the use of symmetries computations will become involved at some point). Bianchi uses directly (3.29) and a counterpart as a homography between u_0, u_3, which is inconvenient because it depends quadratically on v_1. With moderate effort and use of simple symmetries we propose to replace (3.29) and its counterpart with a single relation (3.31), which is separately linear in all variables, just like the cross-ratio property (3.32). The fact that both (3.31) and (3.32) must be true imposes the homography (3.33) which reveals the necessary 'algebraic computations' of the BPT. Our formula (3.33) for QWC (3.6) coincides with (III) in Bianchi (122,§ 23); as for the remaining quadrics Bianchi just claims that such formulae exist by similar computations. He is not far from the truth, since any projective property (like the cross-ratio property) which is valid for a particular quadric is valid for all other quadrics (all quadrics are projectively equivalent), but since in building the confocal family of a given quadric Cayley's absolute $C(\infty)$ plays an important rôle, it remains to prove that the cross-ratio property is also independent of a choice of a conic in a hyperplane of $\mathbb{CP}^3$ as Cayley's absolute (probably a purely synthetic proof unifying all cases exists). It is this what we did; in fact we did more: we found the necessary 'algebraic computations' of the BPT for all quadrics. It may be the case that Bianchi, who first proved the BPT for the pseudo-sphere in 1892 (in which case the proof is easier), has noticed by chance (or looking on purpose for projective properties) the cross-ratio property. Once the cross-ratio property for pseudo-spheres is found it is natural to conjecture that it is true for all quadrics and indeed it is just a matter of computations to prove its validity for general quadrics. Bianchi (122,§ 25) assumes the use of the algebraic identity (3.41) in the proof of the BPT without explicitly stating it: *'Mais toutes ces conditions, ... , ne peuvent être remplies que si la deformation se réduit à un mouvement invariable, ...'*. Its use in the proof of the BPT can be easily justified with a differential symmetry (it is probably what Bianchi had in mind), but to err on the safe side we prefer to use only algebraic symmetries. In fact should we accept differential symmetries just as easily as algebraic symmetries, the BPT would require no proof beyond algebraic computations, since it is essentially a statement about differential symmetries.

We conclude § 3 with further iterations of the TC: Möbius configurations $\mathcal{M}_n$ (the TC is $\mathcal{M}_1$ and the SITC is $\mathcal{M}_2$). Again Bianchi's computations from ([5],Vol **5**,(117)) proves their existence and assures that they can be obtained only from algebraic computations; we provide a simple geometric argument for their existence (using for example the Menelaus theorem) and provide the actual algebraic computations. As chance has it, I was reading at leisure Boyer [8] during my office

* After noticing the link to *The Method* of Archimedes this observation lost most of its initial purpose.

hours around the days I was trying to find the algebraic computations supporting the existence of $\mathcal{M}_3$; it is then that I found the Menelaus theorem and I realized that it was what I needed. The algebraic identity (3.41) naturally appears and plays an important rôle in the analytic computations of Möbius configurations, thus closing the list of *all* algebraic identities necessary for the moving part. Möbius configurations will play an important rôle in the discussion of DDQ and thus in creating computer images of deformations of quadrics.

Beginning with § 4 we present basic results and formulae about the moving part. While most of these results are due to the classical geometers, the compact notation makes them more comprehensible and other issues become more clear.

On occasion there are small observations which were not apparent to the classical geometers; for example Bianchi did not notice that fact that pairs applicable surfaces in $\mathbb{C}^3$ are in correspondence with infinitesimal deformations of surfaces in $\mathbf{O}_3(\mathbb{C})$ (which become the rotations of the rollings), although he was in possession of that differential equation and of other geometric interpretations.

In § 5, 6 we present the important issues (in my opinion at this time) of the moving part for deformations of quadrics in $\mathbb{C}^3$ and respectively in higher dimensions. Note that many important issues from Bianchi are not covered; for example the *permanent conjugate system* (the conjugate system common to the surface in the considered quadric and to its seed deformation; it is also the permanent conjugate system on the leaf). The permanent conjugate system is best suited to describe the B transformation at the analytic level (this was Bianchi's original approach to the theory of deformations of quadrics: he began with Guichard's result and found the analytic formulation of the B transformation for quadrics of revolution and diagonal paraboloids; take for example the article ([5], Vol **4**,(107)) of 1905 where everything was in place except the static picture of confocal quadrics and the Ivory affinity). Also it is best suited to describe triply conjugate systems of surfaces containing a family of deformations of quadrics and it may be the correct tool in attacking the higher dimensional problem of conjugate systems containing a family of deformations of quadrics (orthogonal systems are clearly present in the current literature as examples of integrable systems; see for example [35]).

2. Algebraic preparatives for quadrics in $\mathbb{C}^n$

2.1. **Confocal quadrics in canonical form.**

Consider a non-degenerate symmetric bilinear form (scalar product) on $\mathbb{C}^n$: $< x, y >:= x^TQy,\ Q = Q^T \in \mathbf{GL}_n(\mathbb{C})$. Since the map $\mathbf{GL}_n(\mathbb{C}) \ni A \mapsto A^TA$ has kernel $\mathbf{O}_n(\mathbb{C})$, it has $\frac{n(n+1)}{2}$-dimensional connected image ($\mathbf{GL}_n(\mathbb{C})$ is connected), so $Q = A^TA$ for some $A \in \mathbf{GL}_n(\mathbb{C})$ (the set of symmetric matrices in $\mathbf{GL}_n(\mathbb{C})$ is connected); equivalently A is the product of row (and thus A^T the product of same column) operations which bring Q to the reduced echelon form. Under the linear transformation $\mathbb{C}^n \ni x \mapsto Ax$, the scalar product can be considered the Euclidean one: $< x, y >:= x^Ty,\ |x|^2 := x^Tx$. Consider the standard basis $\{e_j\}_{j=1,...,n},\ e_j^Te_k = \delta_{jk}$ of $\mathbb{C}^n$ and $f_j := \frac{e_{2j-1}-ie_{2j}}{\sqrt{2}},\ j = 1, ..., [\frac{n}{2}],\ I_{j,k} := \Sigma_{l=j}^k e_l e_l^T$.

Consider a quadric $x \subset \mathbb{C}^n$: $Q(x) := x^T(Ax + 2B) + C = 0,\ A = A^T \in \mathbf{M}_n(\mathbb{C}),\ B \in \mathbb{C}^n,\ C \in \mathbb{C},\ \begin{vmatrix} A & B \\ B^T & C \end{vmatrix} \neq 0.$

If $\ker(A) = 0$, then $\begin{vmatrix} A & B \\ B^T & C \end{vmatrix} \begin{vmatrix} I_n & -A^{-1}B \\ 0_{1,n} & 1 \end{vmatrix} = \begin{vmatrix} A & 0_{n,1} \\ B^T & Q(-A^{-1}B) \end{vmatrix}$, so we have $Q(-A^{-1}B) \neq 0$. These are QC, since they have center of symmetry o (which remains center of symmetry under rigid motions and homotheties): $x \in Q \Leftrightarrow 2o - x \in Q$, equivalent to $Q(2o) = Q(0),\ Ao + B = 0$ and $o := -A^{-1}B$ satisfies both.

If $M = M^T \in \mathbf{M}_n(\mathbb{C}),\ \ker(M) = F\mathbb{C}^j \oplus E\mathbb{C}^k,\ F := [f_1...f_j],\ E := [e_{2j+1}...e_{2j+k}]$, then M restricted to $\bar{F}\mathbb{C}^j \oplus I_{2j+k+1,n}\mathbb{C}^n$ and co-restricted to $M(\mathbb{C}^n) = F\mathbb{C}^j \oplus I_{2j+k+1,n}\mathbb{C}^n$ is invertible. Note $F^TF = 0,\ \bar{F}^TF = I_j,\ \bar{F}F^T + F\bar{F}^T = I_{1,2j},\ I_{2j+1,n}F = 0$ (the last $n - 2j$ rows of F are 0). Denote $M^{\circ -1}$ the inverse of the restricted and co-restricted M, extended as

0 on $\bar{F}\mathbb{C}^j \oplus E\mathbb{C}^k$: $M^{\circ -1} := (M + \bar{F}\bar{F}^T + EE^T)^{-1} - FF^T - EE^T$, $MM^{\circ -1} = I_n - \bar{F}F^T - EE^T$, $MM^{\circ -1}M = M$, $M^{\circ -1}MM^{\circ -1} = M^{\circ -1}$ and $Mv = e^{-a}v$, $v \in \mathbb{C}^n \setminus \{0\} \Leftrightarrow M^{\circ -1}v = e^a v$. As we shall see later, the kernel of any symmetric matrix can be brought to the form above by applying rotations $R \in \mathbf{O}_n(\mathbb{C})$, so we can define $M^{\circ -1} := R^T(RMR^T)^{\circ -1}R$ for general $M = M^T \in \mathbf{M}_n(\mathbb{C})$, $\ker(RMR^T) = F\mathbb{C}^k \oplus E\mathbb{C}^p$. Although $M^{\circ -1}$ may depend of the choice of R, we still have $MM^{\circ -1}M = M$, $M^{\circ -1}MM^{\circ -1} = M^{\circ -1}$ and $Mv = e^{-a}v$, $v \in \mathbb{C}^n \setminus \{0\} \Leftrightarrow M^{\circ -1}v = e^a v$.

If $\ker(A) = \mathbb{C}v$, $v \in \mathbb{C}^n \setminus \{0\}$, then $v^T B \neq 0$ (otherwise $[v^T\ 0]^T \in \ker \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$). Such Q does not have center and is thus (I)QWC according to $|v|^2 (= 0) \neq 0$. However, it has another set of interest, the vertex o: for QWC it satisfies $A^{\circ -1}(Ao + B) = 0$, $Q(o) = 0$, equivalent to $o = -A^{\circ -1}B + \frac{B^T A^{\circ -1}B - C}{2v^T B}v$ (o is a well defined point and remains vertex under rigid motions: $o \circ (R,t) = (R,t) \circ o$ and homotheties). A similar formula for IQWC fails, since $A^{\circ -1}$ (and thus o) depend on the rotation R with $\ker(RAR^T) = \mathbb{C}f_1$, but since $A^{\circ -1}$ always has isotropic 1-dimensional kernel (in fact as we shall see later, if $|w|^2 = 0$, $w^T v \neq 0$, then we can make $\ker(A^{\circ -1}) = \mathbb{C}w$), we can define the vertex of an IQWC to be the co-dimension 1 sub-manifold of Q along which normal directions are isotropic: $\mathcal{O} := \{o \in \mathbb{C}^n |\ |Ao + B|^2 = Q(o) = 0\}$. $\mathcal{O}$ remains vertex under rigid motions and homotheties; a canonical choice of a vertex $o \in \mathcal{O}$ will appear later.

If $\dim(\ker(A)) \geq 2$, then $\begin{vmatrix} A & B \\ B^T & C \end{vmatrix} = 0$.

We would like to reduce x, by applying rigid motions $(R,t) \in \mathbf{O}_n(\mathbb{C}) \ltimes \mathbb{C}^n$, $(R,t)x := Rx + t$, to the canonical form (its equation depending on as few constants as possible); equivalently prescribing a coset of the action of rigid motions on quadrics. Once we prescribe the rotation R such that RAR^T depends on as few constants as possible (SJ form) the translation will be $-o$, so as to bring the center (vertex) to the origin.

Complete a maximal set of orthonormal eigenvectors of A to an orthonormal basis of $\mathbb{C}^n$ and consider the vectors of this basis as the rows of the rotation R. RAR^T is decomposed into two blocks on the diagonal, one block being a diagonal matrix and the other a symmetric matrix with isotropic eigenspaces; we have thus reduced the problem to symmetric matrices A with isotropic eigenspaces.

Consider an isotropic subspace $W\mathbb{C}^p \subset \mathbb{C}^n$, $W := [w_1 ... w_p]$ of rank p; since $W^T W = 0$ we have $w_1^T W = 0$. Because of the transitive action of $\mathbf{O}_n(\mathbb{C})$ on the isotropic cone $|x|^2 = 0$, we can make $w_1 = f_1$; replacing $WI_{2,p}$ with $W - f_1(\bar{f}_1^T W) = I_{3,n}W$ we make $WI_{2,p} = I_{3,n}WI_{2,p}$; by induction hypothesis we can make $W = F := [f_1 ... f_p]$.

Just as for the diagonalization process for symmetric matrices with non-isotropic eigenvectors, we would like to find rotations which preserve isotropic subspaces of $\mathbb{C}^n$ (the translations are the subspaces themselves). A rotation R preserving $F\mathbb{C}^p$ must satisfy $RF = FL$, $L \in \mathbf{GL}_p(\mathbb{C})$; with $R' := I_{1,2p} + I_{2p+1,n}RI_{2p+1,n} \in \mathbf{M}_n(\mathbb{C})$, $M := -\bar{F}^T R^T I_{2p+1,n} \in \mathbf{M}_{p,n}(\mathbb{C})$, $N := \bar{F}^T R^T (F\bar{F}^T + \frac{1}{2}I_{2p+1,n})R\bar{F} \in \mathbf{M}_p(\mathbb{C})$ we have $R'R'^T = I_n - I_{2p+1,n}(FL\bar{F}^T R^T + R\bar{F}L^T F^T)I_{2p+1,n} = I_n$, $N + N^T = \bar{F}^T R^T R\bar{F} = 0$, $R^T F = FL^{-1}$, $LMR' = -L\bar{F}^T R^T (I_n - F\bar{F}^T - \bar{F}F^T)RI_{2p+1,n} = \bar{F}^T RI_{2p+1,n}$, $L(N - \frac{1}{2}MM^T) = L\bar{F}^T R^T F\bar{F}^T R\bar{F} = \bar{F}^T R\bar{F}$, so $R = FL\bar{F}^T + \bar{F}(L^T)^{-1}F^T + FL(N - \frac{1}{2}MM^T)F^T + FLMR' - M^T F^T + I_{2p+1,n}R' =: R(L,M,N,R')$. Conversely, given $L \in \mathbf{GL}_p(\mathbb{C})$, $MI_{2p+1,n} = M \in \mathbf{M}_{p,n}(\mathbb{C})$, $-N^T = N \in \mathbf{M}_p(\mathbb{C})$, $R' \in \mathbf{O}_n(\mathbb{C})$, $R'I_{1,2p} = I_{1,2p}$ we have $R(L,M,N,R') \in \mathbf{O}_n(\mathbb{C})$, $R(L,M,N,R')F = FL$. If $R_j := R(L_j, M_j, N_j, R'_j)$, $j = 1,2$, then $R_1 R_2 = R(L,M,N,R')$, $L := L_1 L_2$, $M := M_2 R_1'^T + L_2^{-1}M_1$, $N := N_2 + L_2^{-1}N_1(L_2^T)^{-1} - \frac{1}{2}(L_2^{-1}M_1 R'_1 M_2^T - (L_2^{-1}M_1 R'_1 M_2^T)^T)$, $R' := R'_1 R'_2$, $I_n = R(I_p, 0, 0, I_n)$, $R^{-1}(L,M,N,R') = R(L^{-1}, -LMR', -LNL^T, R'^T)$.

We have the decomposition $R(L,M,N,R') = R(L)R(M)R(N)R(R')$,
$R(L) := FL\bar{F}^T + \bar{F}(L^T)^{-1}F^T + I_{2p+1,n}$, $R(M) := I_n + FM - (I_n + \frac{1}{2}FM)M^T F^T$,
$R(N) := I_n + FNF^T$, $R(R') := R'$ and the relations $R(L_1)R(L_2) = R(L_1 L_2)$,
$R(M)R(L) = R(L)R(L^{-1}M)$, $R(N)R(L) = R(L)R(L^{-1}N(L^T)^{-1})$, $R(R')R(L) = R(L)R(R')$,

$R(M_1)R(M_2)=R(M_1+M_2)R(\frac{1}{2}(M_2M_1^T-M_1M_2^T))$, $R(N)R(M)=R(M)R(N)$,
$R(R')R(M)=R(MR'^T)R(R')$, $R(N_1)R(N_2)=R(N_1+N_2)$, $R(R')R(N)=R(N)R(R')$,
$R(R'_1)R(R'_2)=R(R'_1R'_2)$.

At the level of the Lie algebra $\mathbf{o}_n(\mathbb{C})$ we have $r\in\mathbf{o}_n(\mathbb{C})$, $rF=Fl$, $l\in\mathbf{M}_p(\mathbb{C})\Leftrightarrow r=r(l,m,n,r'):=Fl\bar{F}^T-\bar{F}l^TF^T+Fm-m^TF^T+FnF^T+r'$, $mI_{2p+1,n}=m\in\mathbf{M}_{p,n}(\mathbb{C})$, $n\in\mathbf{o}_p(\mathbb{C})$, $I_{2p+1,n}r'=r'\in\mathbf{o}_n(\mathbb{C})$, $[r(l_1,m_1,n_1,r'_1),r(l_2,m_2,n_2,r'_2)]=r([l_1,l_2],l_1m_2-l_2m_1+m_1r'_2-m_2r'_1,l_1n_2+n_2l_1^T-(l_2n_1+n_1l_2^T)+m_2m_1^T-m_1m_2^T,[r'_1,r'_2])$, $R(e^l)=e^{r(l)}$, $R(m)=e^{r(m)}$, $R(n)=e^{r(n)}$, $R(e^{r'})=e^{r(r')}$. Note $r(m)^3=r(n)^2=0$; this allows simple integral formulations of $M(t)$, $N(t)$: if $R(t):=e^{rt}$, then $rdt=R^TdR$, $ldt=\bar{F}^TR^TdRF=L^{-1}dL$,
$r'dt=I_{2p+1,n}R^TdRI_{2p+1,n}=R'^TdR'$, $mdt=-\bar{F}^TR^TdRI_{2p+1,n}=L^{-1}d(LM)R'$,
$n=\bar{F}^TR^TdR\bar{F}=dN+(lN+Nl^T)dt+\frac{1}{2}((L^{-1}d(LM)M^T)^T-L^{-1}d(LM)M^T)$, so $L(t)=e^{lt}$, $R'(t)=e^{r't}$, $M(t)=\int_0^t e^{-(t-s)l}me^{-sr'}ds$,
$N(t)=\int_0^t e^{-sl}(\frac{1}{2}\int_0^{t-s}(me^{-r'u}m^Te^{-l^Tu}-e^{-lu}me^{r'u}m^T)du+n)e^{-sl^T}ds$.

Example

$R\in\mathbf{O}_n(\mathbb{C})$ *which preserve* $f_1^Tx=0$ *are of the form* $R=R(c)R(v)R(R')$, $R(c):=e^{ic}f_1\bar{f}_1^T+e^{-ic}\bar{f}_1f_1^T+I_{3,n}$, $c\in\mathbb{C}$, $R(v):=I_n+f_1v^T-(\frac{f_1v^T}{2}+I_n)vf_1^T$, $v=I_{3,n}v\in\mathbb{C}^n$, $R(R'):=R'\in\mathbf{O}_n(\mathbb{C})$, $I_{1,2}R'=I_{1,2}$. *Further* $R(c_1)R(c_2)=R(c_1+c_2)$, $R(v)R(c)=R(e^{-ic}v)R(c)$, $R(R')R(c)=R(c)R(R')$, $R(v_1)R(v_2)=R(v_1+v_2)$, $R(R')R(v)=R(R'v)R(R')$ *and* $\{R(c)R(v)\bar{f}_1|\ c,\ v\}=\{x\in\mathbb{C}^n|\ |x|^2=0,\ f_1^Tx\neq0\}$.

In what follows up to obtaining the SJ form we shall always have $k,l\in\{1,...,m\}$, $j\in\{2,...,m\}$. Consider m orthogonal isotropic subspaces $W_k\mathbb{C}^{j_k-j_{k-1}=:p_k}\subset\mathbb{C}^n$, $W_k:=[w_{j_{k-1}+1}...w_{j_k}]$ of rank p_k, $j_0:=0$, $j_m:=p$, $W:=[W_1...W_m]$ of rank p, $W^TW=0$. By applying rigid motions $R\in\mathbf{O}_n(\mathbb{C})$ we would like to make $W_k=F_k:=[f_{j_{k-1}+1}...f_{j_k}]$ and thus $W=F:=[F_1...F_m]$. As above we can make $W_1=F_1$ and $W=F_1(\bar{F}_1^TW)+I_{2p_1+1,n}W$. We have orthogonal isotropic subspaces $I_{2p_1+1,n}W_j\mathbb{C}^{p_j}\subset I_{2p_1+1,n}\mathbb{C}^n\simeq\mathbb{C}^{n-2p_1}$, $I_{2p_1+1,n}W$ of rank $p-p_1$; by the induction hypothesis there exists $R'_1\in\mathbf{O}_n(\mathbb{C})$, $R'_1I_{1,2p_1}=I_{1,2p_1}$ which makes $I_{2p_1+1,n}W=FI_{p_1+1,p}=I_{2p_1+1,n}F$. We can finish by choosing $M_1:=-\bar{F}_1^TW\bar{F}^TI_{2p_1+1,n}:\ R(M_1)W=F$.

Consider the distinct eigenvalues a_k of $A=A^T\in\mathbf{M}_n(\mathbb{C})$ and their isotropic eigenspaces $\ker(A-a_kI_n)=W_k\mathbb{C}^{p_k}$, W_k as above. Thus we can make $W=F$ and with $a:=\mathrm{diag}[a_1I_{p_1}...a_mI_{p_m}]$, $\mathcal{A}'^T=\mathcal{A}':=\bar{F}^TA\bar{F}\in\mathbf{M}_p(\mathbb{C})$, $\mathcal{H}I_{2p+1,n}=\mathcal{H}:=\bar{F}^TAI_{2p+1,n}\in\mathbf{M}_{p,n}(\mathbb{C})$, $\mathcal{A}I_{2p+1,n}=\mathcal{A}^T=\mathcal{A}:=I_{2p+1,n}AI_{2p+1,n}\in\mathbf{M}_n(\mathbb{C})$ we have $A=Fa\bar{F}^T+\bar{F}aF^T+F\mathcal{A}'F^T+F\mathcal{H}+\mathcal{H}^TF^T+\mathcal{A}$. Since $A-a_kI_n$ has kernel $F_k\mathbb{C}^{p_k}$ and eigenspaces $F_l\mathbb{C}^{p_l}$ for eigenvalues a_l-a_k, $l\neq k$, $A-a_kI_n$ restricted to subspaces of $(I_n-F\bar{F}^T)\mathbb{C}^n=\bar{F}\mathbb{C}^p+I_{2p+1,n}\mathbb{C}^n$ and co-restricted to its image is invertible.

In order to further simplify A we would like to consider rotations R which preserve each $F_k\mathbb{C}^{p_k}$. As above these have the form $R(L,M,N,R')$, but we further require that $RF_k=F_kL_k$ for each k (there are no other restrictions on M,N,R'), so a block decomposition of L,M,N appears: $L=\mathrm{diag}[L_1...L_m]\in\mathbf{GL}_p(\mathbb{C})$, $M=[M_1^T...M_m^T]^T\in\mathbf{M}_{p,n}(\mathbb{C})$, $-N^T=N=(N_{lk})_{l,k}\in\mathbf{M}_p(\mathbb{C})$, $R'\in\mathbf{O}_n(\mathbb{C})$, $R'I_{1,2p}=I_{1,2p}$.

Under such an R we have $aL=La$, $Fa\bar{F}^T+\bar{F}aF^T$ is preserved and $\mathcal{A}$, $\mathcal{H}$, $\mathcal{A}'$ respectively become $R(\mathcal{A}):=R'\mathcal{A}R'^T$, $R(\mathcal{H}):=L(\mathcal{H}R'^T+MR(\mathcal{A})-aM)$, $R(\mathcal{A}'):=L(\mathcal{A}'+Na-aN-\frac{1}{2}((MR(\mathcal{A})-aM)M^T+M(MR(\mathcal{A})-aM)^T))L^T+R(\mathcal{H})(LM)^T+(LM)R(\mathcal{H})^T$. In particular $R(M)(\mathcal{H})=\mathcal{H}+M\mathcal{A}-aM$, $R(N)(\mathcal{A}')=\mathcal{A}'+Na-aN$; if further $M\mathcal{A}-aM=0$, then $R(L,M)(\mathcal{H})=L\mathcal{H}$, $R(L,M)(\mathcal{A}')=L(\mathcal{A}'+\mathcal{H}M^T+M\mathcal{H}^T)L^T$.

By induction hypothesis we can prescribe R' so that $R(\mathcal{A})$ is SJ (note that $\det(A-\lambda I_n)=\prod_{k=1}^m(a_k-\lambda)^{2p_k}\det(I_{1,2p}+\mathcal{A}-\lambda I_{2p+1,n})$ and since all eigenvalues and eigenspaces of A are accounted for, eigenvalues of the restriction and co-restriction of $\mathcal{A}$ to $\mathbb{C}^{n-2p}\simeq I_{2p+1,n}\mathbb{C}^n$ are among the eigenvalues of A, while its eigenspaces are $\subseteq I_{2p+1,n}\mathbb{C}^n$ and thus disjoint from the eigenspaces of A, which are $\subset I_{1,2p}\mathbb{C}^n$).

Let $\mathcal{H}=:[\mathcal{H}_1^T...\mathcal{H}_m^T]^T$, $\mathcal{H}_kI_{2p+1,n}=\mathcal{H}_k\in\mathbf{M}_{p_k,n}$; if $\ker(I_{1,2p}+\mathcal{A}-a_1I_{2p+1,n})=F_{m+1}\mathbb{C}^{p_{m+1}}\oplus E_1\mathbb{C}^{q_1}$, $F_{m+1}:=[f_{p+1}...f_{p+p_{m+1}}]$, $E_1:=[e_{2(p+p_{m+1})+1}...e_{2(p+p_{m+1})+q_1}]$, then take $M_1:=-\mathcal{H}_1(I_{1,2p}+$

$\mathcal{A}-a_1I_{2p+1,n})^{\circ-1}$, which makes $\mathcal{H}_1=\mathcal{H}_1(F_{m+1}\bar{F}_{m+1}^T+E_1E_1^T)=:\mathcal{H}_1'[\bar{F}_{m+1}\ E_1]^T$, $\mathcal{H}_1'\in\mathbf{M}_{p_1,r_1:=p_{m+1}+q_1}$; we can apply the same simplification to $\mathcal{H}_2,...,\mathcal{H}_m$. With $\mathcal{A}':=(\mathcal{A}_{kl}')_{kl}$, $\mathcal{A}_{kl}'\in\mathbf{M}_{p_k,p_l}(\mathbb{C})$ take $N_{kl}:=\frac{\mathcal{A}_{kl}'}{a_k-a_l}$, $k\neq l$, which makes $\mathcal{A}_{kl}'=0$, $k\neq l$. In order to preserve the properties so far obtained we can further arbitrarily prescribe L and M with the restriction $M\mathcal{A}-aM=0$: for such M we have $M_1=:M_1'[F_{m+1}\ E_1]^T$, $M_1'\in\mathbf{M}_{p_1,r_1}$ and similar formulae for M_j; as $\ker(I_{1,2p}+\mathcal{A}-a_kI_{2p+1,n})$, $\ker(I_{1,2p}+\mathcal{A}-a_lI_{2p+1,n})$, $k\neq l$ are subspaces of non-isotropic orthogonal subspaces of $\mathbb{C}^n$ we have $\mathcal{H}_kM_l^T=0$, $k\neq l$. We still have to simplify $\mathcal{H}_k'$, $\mathcal{A}_{kk}'$; by induction it is enough to prescribe L_1, M_1' so as to simplify $\mathcal{H}_1'$, $\mathcal{A}_{11}'$. We have $\mathcal{H}_j(F_{m+1}\mathbb{C}^{p_{m+1}}\oplus E_1\mathbb{C}^{q_1})=0$ (again because eigenspaces for different eigenvalues of $\mathcal{A}$ are supported in orthogonal non-isotropic subspaces of $\mathbb{C}^n$); since $A-a_1I_n$ restricted to $F_{m+1}\mathbb{C}^{p_{m+1}}\oplus E_1\mathbb{C}^{q_1}$ and co-restricted to its image $F_1\mathcal{H}_1'\mathbb{C}^{r_1}$ is invertible, $\mathcal{H}_1'$ must have maximal rank $r_1\leq p_1$. Take L_1 the product of the row operations which bring $\mathcal{H}_1'$ to its reduced echelon form $\mathcal{H}_1'=[e_1...e_{r_1}]$; we can further prescribe L_1 with the restriction $L_1\mathcal{H}_1'=\mathcal{H}_1'$ (equivalently L_1 is the product of further arbitrary row operations not involving the pivot rows). Now choose $[M_1'\ 0_{p_1,p_1-r_1}]:=-I_{r_1+1,p_1}\mathcal{A}_{11}'I_{1,r_1}-\frac{1}{2}I_{1,r_1}\mathcal{A}_{11}'I_{1,r_1}$, which makes $\mathcal{A}_{11}'=I_{r_1+1,p_1}\mathcal{A}_{11}'I_{r_1+1,p_1}$; since $A-a_1I_n$ restricted to $\bar{F}_1I_{r_1+1,p_1}\mathbb{C}^{p_1}$ and co-restricted to its image $F_1\mathcal{A}_{11}'$ is invertible, $\mathcal{A}_{11}'$ must have maximal rank p_1-r_1. Further choose L_1 the product of row (and thus L_1^T the product of same column) operations which bring $\mathcal{A}_{11}'$ to the reduced echelon form I_{r_1+1,p_1}. We now permute the coordinates of $\mathbb{C}^n$ in order to follow each eigenvector of A on the flag structure A, $\mathcal{A},...$ along its eigenvalue:

Any matrix $A=A^T\in\mathbf{M}_n(\mathbb{C})$ can be reduced, via conjugation by a rotation $R\in\mathbf{O}_n(\mathbb{C})$, to a SJ matrix: a block diagonal matrix formed by SJ blocks aI_p+J_p, $J_1:=0$, $J_p:=V_pI_p'\bar{V}_p^T$, $p\geq 2$, where $I_p':=\Sigma_{j=1}^{p-1}e_je_{j+1}^T\in\mathbf{M}_p(\mathbb{C})$, $p\geq 1$ is the usual upper diagonal in a Jordan block, $\hat{I}_p:=\Sigma_{j=1}^pe_je_{p+1-j}^T$, $V_{2p}:=[F\ \bar{F}\hat{I}_p]$, $V_{2p+1}:=[F\ e_{2p+1}\ \bar{F}\hat{I}_p]$, $F:=[f_1...f_p]$. Moreover $\bar{V}_p^TV_p=I_p$, $V_p\hat{I}_p=\bar{V}_p$, so $J_p^T=J_p$, $J_p^{\circ-1}=\bar{J}_p$; J_p has $\bar{f}_1$ as cyclic vector of order p, so J_p has only eigenvalue 0 with only eigendirection f_1 and minimal polynomial x^p.

Define the type of a SJ matrix A to be the set of all matrices M with same block diagonal decomposition as that of A, each block J_p being replaced with a polynomial $P(J_p)$ in J_p, $P'(0)\neq 0$. Then two matrices of the same type commute, a matrix with same block diagonal decomposition as that of a SJ matrix A and which commutes with A has the type of A (without the restriction $P'(0)\neq 0$), M^{-1} has the type of M when it exists, $f(I_n-zM)$ has the type of M if f is analytic near 1 and $z\in\mathbb{C}\setminus(\mathrm{spec}(M))^{-1}$ is in the domain of convergence of f (we are interested mostly in $\sqrt{I_n-zM}$).

Although a SJ matrix as defined above efficiently encodes all metric invariants of a symmetric complex matrix, we prefer when convenient to work with the SJ type, since it is closed under certain algebraic operations; the stronger SJ notion would require a rotation after each algebraic operation.

Given $\sum_{j=1}^{n-1}b_jJ_n^j$, $b_1\neq 0$ we know that there is $R\in\mathbf{O}_n(\mathbb{C})$ such that $R(\sum_{j=1}^{n-1}b_jJ_n^j)R^T=J_n$; moreover $RJ_nR^T=J_n$, $R\in\mathbf{O}_n(\mathbb{C})\Rightarrow R=\pm I_n$. With $S:=\bar{V}_n^TRV_n$ we have $\hat{I}_nS^T\hat{I}_nS=I_n$ and we need $S(\sum_{j=1}^{n-1}b_jI_n'^j)\hat{I}_nS^T\hat{I}_n=I_n'$. If $S:=b_1^{-\frac{1+n}{2}}\sum_{j=1}^nb_1^je_je_j^T$, then $\hat{I}_nS^T\hat{I}_nS=I_n$, $b_1SI_n'\hat{I}_nS^T\hat{I}_n=I_n'$, so we can assume $b_1=1$. With $\mathcal{K}:=(k_1,...,k_{n-1})$, $k_1,...,k_{n-1}\in\{0,...,n-1\}$, $s_1(\mathcal{K}):=\sum_{j=1}^{n-1}k_j$, $s_2(\mathcal{K}):=\sum_{j=1}^{n-1}jk_j$, $a_{kj}:=\sum_{s_1(\mathcal{K})=k,s_2(\mathcal{K})=j}\binom{k}{k_1,...,k_{n-1}}b_2^{k_2}...b_{n-1}^{k_{n-1}}$ we have $(a_{kj})_{k,j=1,...,n-1}-I_{n-1}$ strictly upper triangular and $(\sum_{j=1}^{n-1}b_jI_n'^j)^k=\sum_{j=k}^{n-1}a_{kj}I_n'^j$, $k=1,...,n-1$. Thus $\sum_{j=k}^{n-1}a_{kj}SI_n'^j\hat{I}_nS^T\hat{I}_n=I_n'^k$, $k=1,...,n-1$ and $SI_n'^k\hat{I}_nS^T\hat{I}_n=\sum_{j=k}^{n-1}b_{kj}I_n'^j$, where $(b_{kj})_{k,j=1,...,n-1}-I_{n-1}$ is strictly upper triangular and $\sum_{j=k}^qa_{kj}b_{jq}=\delta_{kq}$. Now $\sum_{j=1}^{n-k}(Se_j)(Se_{n+1-k-j})^T=\sum_{j=k}^{n-1}b_{kj}\sum_{q=1}^{n-j}e_qe_{n+1-q-j}^T$. By induction after decreasing $k=n-1,...,1$, $S-I_n$ is strictly upper triangular and $s_{jk}:=e_j^TSe_k$ can be found by induction after decreasing k from $s_{j\ n-k}=b_{k\ n-j}-\sum_{l=1}^{n-k-j}s_{1\ l+1}s_{j\ n-k-l}$, $k=1,...,n-1$, $j=1,...,n-k$.

We are interested in the rotations which preserve the SJ form of A; since $\ker(A-a_jI_n)^k$ increases with k and stabilizes to the subspace of $\mathbb{C}^n$ on which the SJ blocks corresponding to the eigenvalue

a_j of A are supported, such a rotation preserves these subspaces, so it admits a diagonal block decomposition of rotations of such subspaces. We have thus reduced the problem to A having only eigenvalue $a_1 = 0:\ A = \bigoplus_{j=1}^k J_{p_j},\ p_1 \geq ... \geq p_k \geq 1,\ \sum_{j=1}^k p_j = n$ and we need $R \in \mathbf{O}_n(\mathbb{C})$ with $RAR^T = A$. If $k = 1,\ p_1 = 2$, then $n = 2k,\ R = R(L)R(N) = FL\bar{F}^T + \bar{F}LF^T + FLNF^T,\ F := [f_1...f_k],\ L \in \mathbf{O}_k(\mathbb{C}),\ -N^T = N \in \mathbf{M}_k(\mathbb{C})$. If $k = 2,\ p_1 = 3,\ p_2 = 2$, then let A be formed of k_1 blocks J_3 and k_2 blocks $J_2,\ 3k_1 + 2k_2 = n$. Let $F_1, E_1 \in \mathbf{M}_{n,k_1}(\mathbb{C}),\ F_2 \in \mathbf{M}_{n,k_2}(\mathbb{C})$ respectively be the matrices with columns the f_1's of J_3's, e_3's of J_3's, f_1's of J_2's, considered as vectors in $\mathbb{C}^n$; thus $A = F_1E_1^T + E_1F_1^T + F_2F_2^T$. Using $\ker(RA^j) = \ker(A^jR)$ and $AR = RA$ we get $R = F_1L_1\bar{F}_1^T + (F_2 - F_1M_1^T)L_2\bar{F}_2^T + (\bar{E}_1 + F_2M_1 + F_1(N_1 - \frac{1}{2}M_1^TM_1))L_1E_1^T + (\bar{F}_2 - E_1M_1^T + F_2(N_3 - \frac{1}{2}M_1M_1^T) - F_1(M_1^TN_3 + N_1M_1^T + M_2^T))L_2F_2^T + (\bar{F}_1 + \bar{F}_2M_1 + E_1(N_1 - \frac{1}{2}M_1^TM_1) + F_2M_2 + F_1(N_2 - \frac{1}{2}(M_2^TM_1 + M_1^TM_2 + (N_1 - \frac{1}{2}M_1^TM_1)^T(N_1 - \frac{1}{2}M_1^TM_1))))L_1F_1^T,\ L_j \in \mathbf{O}_{k_j}(\mathbb{C}),\ M_j \in \mathbf{M}_{k_2,k_1}(\mathbb{C}),\ -N_j^T = N_j \in \mathbf{M}_{k_1}(\mathbb{C}),\ j = 1, 2,\ -N_3^T = N_3 \in \mathbf{M}_{k_2}(\mathbb{C})$ arbitrary. While there is a precise determination of R for $p_1 \geq 4$ by similar computations, it increases in complexity (also the relations among such rotations). Again the picture at the level of the Lie algebra should be simpler: $rA - Ar = 0,\ r \in \mathbf{o}_n(\mathbb{C})$ and it becomes clear that for $A = J_n$ such an r must be a polynomial in J_n (thus $r^T = r$ and $r = 0$).

Returning to the quadric $x:\ \ Q(x) = x^T(Ax + 2B) + C = 0$, once A is SJ and $\ker(A)$ spanned by $\emptyset,\ e_n$ respectively for Q(W)C, choose the translation $t := -o$ and we have respectively canonical Q(W)C: $B = 0, -e_n,\ C = -1, 0$ (still denote by A the matrix $-Q(o)^{-1}A,\ \frac{-1}{e_n^TB}A$, although another rotation may be required to make the new A SJ). Therefore for Q(W)C the rigid motions preserving the canonical form are rotations preserving A, so $\mathrm{spec}(A)$ is preserved by such rotations. For IQWC once A is SJ, $\ker(A) = \mathbb{C}f_1$ and the SJ block for the eigenvalue 0 of A is J_p, choose the translation $t := -o$ to get $B = e^{-ic}\bar{f},\ C = 0$. If we conjugate A with a rotation $R \in \mathbf{O}_n(\mathbb{C}),\ RI_{p+1,n} = I_{p+1,n},\ R(J_p \oplus I_{p+1,n})R^T = (e^{2ic_1}J_p) \oplus I_{p+1,n}$ (we use the $\oplus$ to denote the blocks in a block diagonal decomposition; such a rotation R and $-R + 2I_{p+1,n}$ are the only ones with this property), then with $c_1 := -\frac{c}{p+1}$ the vector B becomes $-\bar{f}_1$ and A becomes $e^{-2ic_1}RAR^T$: this is the canonical form (of course, after a further rotation preserving the first p coordinates to bring the remaining part of A to the SJ form). Two canonical forms differ only by a rotation and the corresponding A's must differ in the first p coordinates by a rotation as above with $c_1 \in \frac{\pi}{p+1}\mathbb{Z}$, in which case the spectra of these A's differ by a $(p+1)^{\text{th}}$ root of unity. Thus for canonical IQWC $\mathrm{spec}(A)$ is not preserved by rotations preserving the canonical form, but the collection of spectra $\{e^{\frac{2\pi ij}{p+1}}\mathrm{spec}(A)\}_{j=0,...,p}$ is; for simplicity we shall make a choice of $\mathrm{spec}(A)$ and work with it.

We have the diagonal Q(W)C respectively for $A = \Sigma_{j=1}^n a_j^{-1}e_je_j^T,\ A = \Sigma_{j=1}^{n-1}a_j^{-1}e_je_j^T$; the diagonal IQWC come in different flavors, according to the block of $f_1:\ A = J_p + \Sigma_{j=p+1}^n a_j^{-1}e_je_j^T$; in particular if $A = J_n$, then $\mathrm{spec}(A) = \{0\}$ is unambiguous. Thus general quadrics are those for which all eigenvalues have geometric multiplicity 1; equivalently each eigenvalue has an only corresponding SJ block.

With $R_z := I_n - zA$ the family of quadrics $\{x_z\}_z$ confocal to x_0 is given by $Q_z(x_z) = x_z^TAR_z^{-1}x_z + 2(R_z^{-1}B)^Tx_z + C + zB^TR_z^{-1}B = 0$.

Consider the IQWC $x^T(Ax + 2B) + C = 0,\ \ker(A) = \mathbb{C}v,\ |v|^2 = 0,\ v \in \mathbb{C}^n \setminus \{0\}$ and $o \in \mathcal{O}$. Applying the translation $-o$ and a rotation $R,\ Rv = f_1,\ R(Ao + B) = -e^{-ic}\bar{f}_1,\ e^{-ic} := -v^TB$ we can make $\ker(A) = \mathbb{C}f_1,\ B = -e^{-ic}\bar{f}_1,\ C = 0$. If we begin with a different point in $\mathcal{O}$, we get a similar situation; the difference between the two is a rigid motion (R, t) which preserves the form of $A,\ B,\ C:\ \ker(RAR^T) = \mathbb{C}f_1,\ e^{-ic}R\bar{f}_1 + RAR^Tt = e^{-i(c+c_1)}\bar{f}_1,\ Q(-R^Tt) = 0$, so $Rf_1 = e^{ic_1}f_1$ and $t = e^{-i(c+c_1)}(I_n - f_1\bar{f}_1^T)RA^{\circ -1}R^T\bar{f}_1$ is obtained by plugging into the formula $o = -A^{\circ -1}B + \frac{B^TA^{\circ -1}B - C}{2v^TB}v$ the values $RA^{\circ -1}R^T,\ -e^{-i(c+c_1)}\bar{f}_1,\ 0,\ f_1$ respectively instead of $A^{\circ -1},\ B,\ C,\ v$. In this simplified setting it is easy to see that the formula of o is well defined up to rotations of the type $R(c_1)R(R')$ (which also fix $\mathbb{C}\bar{f}_1$); the remaining type of rotations $R(v)$ give a parametrization of $\mathcal{O}:\ \mathcal{O} = \{e^{ic}(I_n - f_1\bar{f}_1^TR(v)^T)(R(v)AR(v)^T)^{\circ -1}R(v)\bar{f}_1|\ v = I_{3,n}v \in \mathbb{C}^n\}$.

There is a unifying point of view in the projective space: the action of the group of rigid motions and homotheties $\mathbb{C}^n \ni x \mapsto (e^a(R,t))x := e^a(Rx+t),\ (R,t) \in \mathbf{O}_n(\mathbb{C}) \ltimes \mathbb{C}^n,\ a \in \mathbb{C}$ can be extended to $\mathbb{CP}^n \supseteq [\mathbb{C}^{nT},1] \simeq \mathbb{C}^n$ as $(e^a(R,t))[x^T,x^{n+1}] := [(\begin{bmatrix} R & t \\ 0^T & e^{-a} \end{bmatrix} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix})^T] = [(Rx + x^{n+1}t)^T, e^{-a}x^{n+1}]$ and is transitive on Cayley's absolute $C(\infty) := \{[Y^T,0] |\ Y \in \mathbb{C}^n \setminus \{0\},\ |Y|^2 = 0\}$ (circle at infinity for $n=3$); in fact it is the maximal subgroup of $\mathbf{PGL}_n(\mathbb{C})$ which preserves $C(\infty)$. Any quadric in $\mathbb{CP}^{n-1} = [(\mathbb{C}^n \setminus \{0\})^T, 0]$ can be brought, via a homography, to $C(\infty)$, thus confirming that any symmetric non-degenerate bilinear form on $\mathbb{C}^n$ is isometric to the Euclidean scalar product.

We have $\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} = \begin{bmatrix} A^{\circ -1} & b \\ b^T & C \end{bmatrix}$, where A is SJ and respectively for canonical (I)Q(W)C $\ker(A) = 0,\ \mathbb{C}e_n,\ \mathbb{C}f_1,\ B = 0, -e_n, -f_1,\ b = 0, -e_n, -f_1,\ C = -1, 0, 0$. If $z \in \mathbb{C},\ \det R_z \neq 0$, then $\begin{bmatrix} AR_z^{-1} & R_z^{-1}B \\ B^TR_z^{-1} & C + zB^TR_z^{-1}B \end{bmatrix} = \begin{bmatrix} A^{\circ -1} - zI_n & b \\ b^T & C \end{bmatrix}^{-1}$.

A quadric $[x^T, x^{n+1}] \subset \mathbb{CP}^n$ satisfies $\begin{bmatrix} x \\ x^{n+1} \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix} = 0,\ A^T = A,\ \begin{vmatrix} A & B \\ B^T & C \end{vmatrix} \neq 0$ and its family of confocal quadrics $\begin{bmatrix} x_z \\ x_z^{n+1} \end{bmatrix}^T (\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} - z \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix})^{-1} \begin{bmatrix} x_z \\ x_z^{n+1} \end{bmatrix} = 0.$

If we define the adjugate of the quadric $\begin{bmatrix} x \\ x^{n+1} \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix} = 0$ to be the quadric $\begin{bmatrix} x \\ x^{n+1} \end{bmatrix}^{*T} \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix}^* = 0$, then $(\begin{bmatrix} R & t \\ 0^T & e^{-a} \end{bmatrix} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix})^* = (\begin{bmatrix} R & t \\ 0^T & e^{-a} \end{bmatrix}^T)^{-1} \begin{bmatrix} x \\ x^{n+1} \end{bmatrix}^*$; the family of quadrics confocal to $[x^T, x^{n+1}]$ is the adjugate of the pencil generated by the adjugate of $[x^T, x^{n+1}]$ and $C(\infty)$.

Conversely, given a quadric $Q \subset \mathbb{CP}^n$, a hyperplane $P \subset \mathbb{CP}^n$ and a quadric $C(\infty) \subset P$, we can consider the Euclidean metric introduced by $C(\infty)$ on $\mathbb{CP}^n \setminus P$; P becomes the hyperplane $\mathbb{CP}^{n-1}$ at ∞ and we can recover the construction above. If P is not tangent to Q, then Q is QC; if P is tangent to Q (that is $P \cap Q$ is a quadratic cone), then Q is (I)QWC; if $C(\infty)$ does not pass through the point of tangency of P and Q (that is the vertex of the cone $P \cap Q$), then Q is QWC, otherwise Q is IQWC.

If $\ker(A) = 0$ or is not isotropic, A SJ, then we can define $\sqrt{A}$ to be a symmetric matrix of the same type as A: $\sqrt{e^a I_p + J_p} = \sqrt{e^a} \Sigma_{j=0}^{p-1} e^{-ja} (\frac{\frac{1}{2}}{j}) J_p^j,\ p \geq 1$; (the sign of $\sqrt{\cdot}$ does not matter, because any quadric is symmetric wrt the principal spaces, but in order to keep a strict account of these symmetries (to be used for other reasons), we shall always choose $\sqrt{re^{i\theta}} := \sqrt{r}e^{i\frac{\theta}{2}},\ r \geq 0,\ -\pi < \theta \leq \pi$ and we shall avoid simplifications of the form $\sqrt{a}\sqrt{b} = \pm\sqrt{ab} \simeq \sqrt{ab}$). Consider the unit sphere $X \subset \mathbb{C}^n,\ |X|^2 = 1$ and the equilateral paraboloid $Z \subset \mathbb{C}^n,\ Z^T(I_{1,n-1}Z - 2e_n) = 0$; then we have parametrization $x_0 = (\sqrt{A})^{-1}X$ of QC and $x_0 = (\sqrt{A + e_n e_n^T})^{-1}Z$ of QWC. A natural parametrization of IQWC will appear later; because of this we should rather require that $A^{\circ -1}$ is SJ.

For z inverses of nonzero eigenvalues of A (which make R_z singular) we have finite singular quadrics; for $z = \infty$ we have the singular quadric at ∞; all are obtained as sets by letting z tend to the singular value in the closure of $x_z \simeq [x_z^T, 1] \subset \mathbb{CP}^n$. If A is of SJ type with $\ker(R_a)$ spanned by $f_1, ..., f_k, e_{2k+1}, ..., e_p$, then by L'Hospital $x_a = \{x \in \ker(R_a) | |x|^2 = 0\} \times I_{p+1,n}\mathbb{C}^n,\ x_\infty = \mathbb{CP}^{n-1} = [X^T, 0] \cup C(\infty)$; for example if $A = aI_n$, then x_a is the isotropic cone $|x|^2 = 0$. Singular quadrics have extra projective structure: given a quadric $Q \subset \mathbb{C}^n$ and $p \in \mathbb{C}^n$ the (quadratic) tangent cone of Q at p is $C_pQ := \{v \in \mathbb{C}^n | Q(p+sv) = 0 \text{ has a double root } s\}$ and degenerates to the singular quadratic cone T_pQ if $p \in Q$. Thus $\mathbb{C}^n$ can be seen as a quadratic cone bundle over itself, factorized by the relation of equivalence $p \sim q$ iff $p = q$ as points in $\mathbb{C}^n$. The factor set has singular set Q (which can again be seen as quadratic cone bundle over itself factorized by

the same relation of equivalence, but this time the factor set has no singular points, since all cones $Q \cap T_pQ,\ p \in Q$ are projectively equivalent). Similarly singular confocal quadrics can be seen as quadratic cone bundles over themselves, factorized by equivalence relationships whose singular sets are lower dimensional quadrics and this (singular) projective structure of singular quadrics of the confocal family is the limit of the (non-singular) projective structure of non-singular quadrics of the confocal family; the whole confocal family can be recovered from the knowledge of the singular set of certain finite singular quadrics of the family (including those for which the eigenvalue has geometric multiplicity 1). The singular set of x_∞ is $\mathcal{S}(x_\infty) := C(\infty)$; for $z = a_1,\ A$ of the SJ type with $\ker(R_{a_1}) \neq 0$ the singular set $x := \mathcal{S}(x_{a_1})$ of the singular quadric x_{a_1} satisfies: $x^T(AR_{a_1}^{\circ -1}x + 2R_{a_1}^{\circ -1}B) + C + a_1B^TR_{a_1}^{\circ -1}B = 0,\ x = R_{a_1}R_{a_1}^{\circ -1}x$. For a family of confocal quadrics, the spectrum is the set of singular values of z (thus it is not well defined, since it depends on the choice of the initial quadric); the relative spectrum is the set of differences of values in the spectrum (thus it is well defined).

2.2. **Proofs of Lamé, Dupin, Ivory, Bianchi I, Jacobi and Chasles theorems; a ray of light.**
Assume that the quadric x_0 is in the canonical form: A SJ, $\ker(A) = 0,\ \mathbb{C}e_n,\ \mathbb{C}f_1,\ B = 0, -e_n,\ -\bar{f}_1,\ C = -1, 0, 0$ respectively for (I)Q(W)C. The Ivory affinity for QC should preserve the center 0 of the quadrics x_z, so it is given by a linear transformation $x_z = M(z)x_0$, where $M(z) \in \mathbf{GL}_n(\mathbb{C})$ satisfies $M(z)^TAR_z^{-1}M(z) = A$; the simplest choice $M(z) := \sqrt{R_z}$ is the correct one.

For (I)QWC we make the ansatz $x_z = \sqrt{R_z}x_0 + C(z)$ and further require it to satisfy Lamé's: $0 = -2dx_w^T\partial_z|_{z=w}x_z = dx_0^T\sqrt{R_w}(A(\sqrt{R_w})^{-1}x_0 - 2C'(w))$. Using $dx_0^T(Ax_0 + B) = 0$ we get $Ax_0 - 2\sqrt{R_w}C'(w) = c(w, x_0)(Ax_0 + B)$; multiplying on the left with $v^T,\ 0 \neq v \in \ker(A)$ we get $c(w, x_0) = c(w) = -\frac{2v^T\sqrt{R_w}C'(w)}{v^TB}$; making $x_0 = 0$ we get $2\sqrt{R_w}C'(w) = -c(w)B$, so $(1 - c(w))Ax_0 = 0$ and since the quadric x_0 does not lie in $\ker(A)$, we get $c(w) = 1$ and $C(z) = (-\frac{1}{2}\int_0^z(\sqrt{R_w})^{-1}dw)B (= \frac{z}{2}e_n$ for QWC; for IQWC it is the Taylor series of $\frac{1}{2}\int_0^z(\sqrt{1-w})^{-1}dw$ at $z = 0$ with each monomial z^{k+1} replaced by $-z^{k+1}J_p^kB$, where J_p is the block of f_1 and thus a polynomial of degree p in z). Note $AC(z) + (I_n - \sqrt{R_z})B = 0 = (I_n + \sqrt{R_z})C(z) + zB$. Applying d to $Q_z(x_z) = 0$ we get $dx_z^TR_z^{-1}(Ax_z + B) = 0$, so the unit normal N_z is proportional to $\hat{N}_z := -2\partial_z x_z$. If $\mathbb{C}^n \ni x \in x_{z_1}, x_{z_2}$, then $\hat{N}_{z_j} = R_{z_j}^{-1}(Ax + B)$; using $R_z^{-1} - I_n = zAR_z^{-1},\ z_1R_{z_1}^{-1} - z_2R_{z_2}^{-1} = (z_1 - z_2)R_{z_1}^{-1}R_{z_2}^{-1}$ we get $0 = Q_{z_1}(x) - Q_{z_2}(x) = (z_1 - z_2)\hat{N}_{z_1}^T\hat{N}_{z_2}$. The polynomial equation $Q_z(x) = 0$ has degree n in z and it has multiple roots iff $0 = \partial_zQ_z(x) = |\hat{N}_z|^2$.

Ivory becomes $|V_0^1|^2 = |x_0^0 + x_0^1 - C(z)|^2 - 2(x_0^0)^T(I_n + \sqrt{R_z})x_0^1 + zC = |V_1^0|^2$; Bianchi I becomes for lengths of rulings: if $w_0^TAw_0 = w_0^T\hat{N}_0 = 0,\ w_z = \sqrt{R_z}w_0$, then $w_z^Tw_z = |w_0|^2 - zw_0^TAw_0 = |w_0|^2$; for the symmetry of the TC: $(V_0^1)^T\hat{N}_0^0 = (x_0^0)^TA\sqrt{R_z}x_0^1 - B^T(x_z^0 + x_z^1 - C(z)) + C = (V_1^0)^T\hat{N}_0^1$; for angles between segments and rulings: $(V_0^1)^Tw_0^0 + (V_1^0)^Tw_z^0 = -z(\hat{N}_0^0)^Tw_0^0 = 0$; for angles between rulings: $(w_0^0)^Tw_z^1 = (w_0^0)^T\sqrt{R_z}w_0^1 = (w_z^0)^Tw_0^1$; for angles between polar rulings: $(w_z^0)^T\hat{w}_z^0 = (w_0^0)^T\hat{w}_0^0 - z(w_0^0)^TA\hat{w}_0^0 = (w_0^0)^T\hat{w}_0^0$.

For Jacobi's result a geodesic in affine parametrization must satisfy $Q(x_0) = \dot{x}_0^T\hat{N}_0 = \ddot{x}_0^Tdx_0 = 0$, so $0 = (\dot{x}_0^T\hat{N}_0)^{\cdot} = \ddot{x}_0^T\hat{N}_0 + \dot{x}_0^T\dot{\hat{N}}_0$, or $\ddot{x}_0 = -\frac{\dot{x}_0^T\dot{\hat{N}}_0}{|\hat{N}_0|^2}\hat{N}_0$. The tangent line at $x_0(t)$ is tangent to a confocal quadric $x_{z(t)}$ iff the quadratic equation in s: $\frac{1}{2}Q_z(x_0 + s\dot{x}_0) = 0$ has a double solution, so we need its discriminant $T_0 := (\dot{x}_0^TR_z^{-1}\hat{N}_0)^2 - \dot{x}_0^TAR_z^{-1}\dot{x}_0Q_z(x_0)$ to be 0. With $a,\ b,\ c,\ d \in \mathbb{C}^n$ and $[a, b] := ba^T - ab^T,\ [[a, b], c] := (ba^T - ab^T)c$ we have $[a, b] \bullet [c, d] := \frac{1}{2}\mathrm{tr}([a, b]^T[c, d]) = a^T[b, [c, d]] = a^Tcb^Td - a^Tdb^Tc$. Using also $Q_z(x_0) = z\hat{N}_0^TR_z^{-1}\hat{N}_0$ we get $[\dot{x}_0, \hat{N}_0] \bullet [R_z^{-1}\hat{N}_0, R_z^{-1}\dot{x}_0] + |\dot{x}_0|^2\hat{N}_0^TR_z^{-1}\hat{N}_0 = 0$. After eliminating the denominator, T_0 becomes a polynomial of degree $n - 1$ in z with the coefficient of the *highest order term* (hot) being $|\dot{x}_0|^2 \neq 0$, 0 as one root and in general having distinct roots; thus Jacobi's result is proven if $\dot{z} = 0$. Dotting $T_0 = 0$ we get $\dot{T}_0 = T_1 + \dot{z}T_2 = 0$ where $T_1 := [\dot{x}_0, \dot{\hat{N}}_0] \bullet [R_z^{-1}\hat{N}_0, R_z^{-1}\dot{x}_0] + [\dot{x}_0, \hat{N}_0] \bullet [R_z^{-1}\dot{\hat{N}}_0, R_z^{-1}\dot{x}_0] +$

$2|\dot{x}_0|^2\hat{N}_0^T R_z^{-1}\dot{\hat{N}}_0 = 2(\hat{N}_0^T R_z^{-1}\dot{x}_0\dot{x}_0^T AR_z^{-1}\dot{x}_0 - \dot{x}_0^T(R_z^{-1}-I_n)\dot{x}_0\dot{x}_0^T AR_z^{-1}\hat{N}_0) = 0$ and $T_2 := \partial_z T_0 = -[\dot{x}_0,\hat{N}_0]\bullet([AR_z^{-2}\hat{N}_0, R_z^{-1}\dot{x}_0] + [R_z^{-1}\hat{N}_0, AR_z^{-2}\dot{x}_0]) - |\dot{x}_0|^2\hat{N}_0^T AR_z^{-2}\hat{N}_0 \neq 0$ in general; thus $\dot{z}=0$. We have $T_2 = 0$ precisely when z is a multiple root for the considered polynomial, but in this case if $\dot{z}\neq 0$, then similarly we have $\dot{T}_2 = T_3 + \dot{z}T_4,\ T_3 := -[\dot{x}_0,\dot{\hat{N}}_0]\bullet([AR_z^{-2}\hat{N}_0, R_z^{-1}\dot{x}_0] + [R_z^{-1}\hat{N}_0, AR_z^{-2}\dot{x}_0]) - [\dot{x}_0,\hat{N}_0]\bullet([AR_z^{-2}\dot{\hat{N}}_0, R_z^{-1}\dot{x}_0] + [R_z^{-1}\dot{\hat{N}}_0, AR_z^{-2}\dot{x}_0]) - 2|\dot{x}_0|^2\hat{N}_0^T AR_z^{-2}\hat{N}_0 =$
$= -2([\dot{x}_0,\hat{N}_0]\bullet([A^2R_z^{-2}\dot{x}_0, R_z^{-1}\dot{x}_0] + [AR_z^{-1}\dot{x}_0, AR_z^{-2}\dot{x}_0]) + |\dot{x}_0|^2\hat{N}_0^T A^2R_z^{-2}\dot{x}_0) = 0$ (use $zAR_z^{-j} = R_z^{-j} - R_z^{-j+1},\ j=1,2$), so $T_4 := \partial_z T_2 = 0$ and so on, which implies the fact that z has multiplicity $n-1$, so the geodesic is a ruling on x_0, contradicting $\dot{z}\neq 0$. Note that we have proven that the roots of $T_0, T_2, T_4, ...$ or of any linear combination thereof satisfy the same property: they remain constant along the geodesic.

For diagonal QC ($A = \Sigma_{j=1}^n a_j^{-1}e_j e_j^T$) we have $T_0 = \Sigma_{j<k}\frac{(x_0^j\dot{x}_0^k - x_0^k\dot{x}_0^j)^2}{(a_j-z)(a_k-z)} - \Sigma_j\frac{(\dot{x}_0^j)^2}{a_j-z}$; for diagonal QWC ($A = \Sigma_{j=1}^{n-1}a_j^{-1}e_j e_j^T$) we have $T_0 = \Sigma_{j<k}\frac{(x_0^j\dot{x}_0^k - x_0^k\dot{x}_0^j)^2}{(a_j-z)(a_k-z)} + \Sigma_j\frac{2x_0^j\dot{x}_0^j x_0^n - (2x_0^n - a_j)(\dot{x}_0^j)^2}{a_j-z} - |\dot{x}_0|^2$.

The part of Chasles's result involving geodesics can be easily read from the proof of Jacobi's result. Let the $n-1$ confocal quadrics be $x_{z_0:=0},\ x_{z_1},...,x_{z_{n-2}}$ and $z\in\{z_0,z_1,...,z_{n-2}\}$. Consider the line x_0+sv tangent to the $n-1$ quadrics at points $x_z := x_0 + s_z v$. We thus have $(v^T R_z^{-1}\hat{N}_0)^2 - zv^T AR_z^{-1}v\hat{N}_0^T R_z^{-1}\hat{N}_0 = 0,\ s_z = -\frac{v^T R_z^{-1}\hat{N}_0}{v^T AR_z^{-1}v},\ x_z = x_0 + s_z v = \frac{[[v,x_0],AR_z^{-1}v] - v^T R_z^{-1}Bv}{v^T AR_z^{-1}v}$ and $zv^T AR_z^{-1}v\hat{N}_0^T\hat{N}_z = z\hat{N}_0^T R_z^{-1}(A[[v,x_0],AR_z^{-1}v] + [[Av,B],R_z^{-1}v]) = v^T\hat{N}_0\hat{N}_0^T R_z^{-1}v -$
$-(v^T R_z^{-1}\hat{N}_0)^2 + zv^T AR_z^{-1}v\hat{N}_0^T R_z^{-1}\hat{N}_0 = 0$.

Therefore $N_0^T N_z = 0$ along the common tangents and by symmetry this is true for any pair of normals of the $n-1$ confocal quadrics (the choice of the initial quadric x_0 is arbitrary). The remaining part is due to Dupin and Malus (see Darboux ([13],§ 441)): a congruence of lines in $\mathbb{C}^n$ which admits $n-1$ focal sub-manifolds whose normals are orthonormal is normal. We would like to find a sub-manifold $y := x_0 + sv$ orthogonal to unit $v:\ 0 = v^T dy = v^T dx_0 + ds$. Imposing the compatibility condition $d\wedge$ we thus need $0 = dx_0^T\wedge dv$. From $dx_0 = (\Sigma_z N_z N_z^T + vv^T)dx_0$ we thus need to prove that $dx_0^T\Sigma_{z\neq z_0}N_z N_z^T\wedge dv = 0$ and $0 = dx_z^T N_z = (dx_0 + s_z dv)^T N_z$ finishes the proof.

Note that $n-1$ confocal quadrics $Q_{z_1},...,Q_{z_{n-1}}$ are seen orthogonally from any point p on any lines tangent to all $n-1$ quadrics, that is the cones through p tangent to the given quadrics intersect along the principal directions at that point; moreover all common tangent to $Q_{z_1},...,Q_{z_{n-1}}$ and passing through p are generated by reflection in the principal hyperplanes of the initial common tangent through p (Darboux ([13],§ 465)).

2.3. **Proof of Bianchi II.**

For Bianchi II consider $\mathbb{C}^n \simeq [\mathbb{C}^{nT},1]\subset\mathbb{CP}^n$ and a homography $x\mapsto H(x)$ is given by $H(x) := \frac{H'x+h_1}{h_2^T x+a},\ H := \begin{bmatrix} H' & h_1\\ h_2^T & a\end{bmatrix}\in\mathbf{GL}_{n+1}(\mathbb{C}),\ H'\in\mathbf{M}_n(\mathbb{C}),\ h_1,h_2\in\mathbb{C}^n,\ a\in\mathbb{C}$ being determined up to multiplication by a non-zero constant. Suppose H takes a family of confocal quadrics x_z given by $\begin{bmatrix} A & B\\ B^T & C\end{bmatrix}$ to another one $\tilde{x}_{\tilde{z}} := H(x_z),\ \tilde{z}(0)=0$ given by $\begin{bmatrix}\tilde{A} & \tilde{B}\\ \tilde{B}^T & \tilde{C}\end{bmatrix}$. We exclude rigid motions and homotheties (for which $H(C(\infty)) = C(\infty)$) as they do not metrically change the family x_z; as we shall see later H cannot be an affine transformation of $\mathbb{C}^n$ (that is $h_2\neq 0$) and further $\det(H\begin{bmatrix} I_n & 0\\ 0^T & 0\end{bmatrix}H^T - \lambda\begin{bmatrix} I_n & 0\\ 0^T & 0\end{bmatrix}) = \begin{vmatrix} H'H'^T - \lambda I_n & H'h_2\\ (H'h_2)^T & |h_2|^2\end{vmatrix}\neq 0$ for some $\lambda\in\mathbb{C}$ (and thus for all but $\leq n$ $\lambda\in\mathbb{C}$). Let $\tilde{H} := H^{-1}$; in the following we consider tilde quantities as counterparts of non-tilde quantities (and thus are defined to have same properties); all equalities remain valid if tilde and non-tilde quantities are exchanged.

By applying rigid motions in both the domain and range of H we make $\ker(A) = 0,\ \mathbb{C}e_1,\ \mathbb{C}f_1,\ B = 0,\ -e_1,\ -\bar{f}_1,\ b = 0,-e_1,\ -f_1,\ C = -1,0,0$ respectively for (I)Q(W)C. As $\begin{bmatrix} A & B\\ B^T & C\end{bmatrix}^{-1} =$

$\begin{bmatrix} A^{\circ -1} & b \\ b^T & C \end{bmatrix}$ we have $H \begin{bmatrix} A^{\circ -1} - zI_n & b \\ b^T & C \end{bmatrix} H^T = \lambda(z)^{-1} \begin{bmatrix} \tilde{A}^{\circ -1} - \tilde{z}(z) I_n & \tilde{b} \\ \tilde{b}^T & \tilde{C} \end{bmatrix}$; thus the singular values of $z,\ \tilde{z}$ correspond; in particular since $H^{-1}(\mathbb{CP}^{n-1}) \neq \mathbb{CP}^{n-1},\ H^{-1}(C(\infty)),\ H(C(\infty))$ are respectively singular sets of finite singular quadrics of the families $x_z,\ \tilde{x}_{\tilde{z}}$ whose corresponding eigenvalues $a_1,\ \tilde{a}_1$ thus must have geometric multiplicity 1. We need $H \begin{bmatrix} A^{\circ -1} & b \\ b^T & C \end{bmatrix} H^T = \lambda(0)^{-1} \begin{bmatrix} \tilde{A}^{\circ -1} & \tilde{b} \\ \tilde{b}^T & \tilde{C} \end{bmatrix},\ H \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix} H^T = \frac{\lambda(z)^{-1} - \lambda(0)^{-1}}{-z} \begin{bmatrix} \tilde{A}^{\circ -1} & \tilde{b} \\ \tilde{b}^T & \tilde{C} \end{bmatrix} + \frac{\lambda(z)^{-1} \tilde{z}(z)}{z} \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix}$; thus $\frac{\lambda(z)^{-1} - \lambda(0)^{-1}}{-z},\ \frac{\lambda(z)^{-1} \tilde{z}(z)}{z}$ are constants. Under the homothety $e^t\ A,\ z$ become $e^{-2t} A,\ e^{2t} z$ if x_z are QC and $e^{-t} A,\ e^t z$ if x_z are (I)QWC (when $\begin{bmatrix} A^{\circ -1} - zI_n & b \\ b^T & 0 \end{bmatrix}$ becomes $e^t \begin{bmatrix} e^t (A^{\circ -1} - zI_n) & b \\ b^T & 0 \end{bmatrix}$). If $\lambda(z)$ is constant, then $H(C(\infty)) = C(\infty)$; otherwise $\lambda(z)^{-1} = e^s (z - a_1),\ \tilde{z}(z) = a_1 \tilde{a}_1 (a_1^{-1} + (z - a_1)^{-1})$ and by applying homotheties in either the range, domain of H or both we make $\tilde{a}_1 = a_1^{-1}$; further fixing the non-zero constant upon which H depends we make $s = 0$ and $\begin{bmatrix} \tilde{A}^{\circ -1} - \tilde{a}_1 I_n & \tilde{b} \\ \tilde{b}^T & \tilde{C} \end{bmatrix} = -H \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix} H^T,\ H \begin{bmatrix} A^{\circ -1} - a_1 I_n & b \\ b^T & C \end{bmatrix} H^T = - \begin{bmatrix} I_n & 0 \\ 0^T & 0 \end{bmatrix}$. To preserve these we exclude further homotheties in either the range or domain of H.

Applying rigid motions in both the domain and range of H, we can suppose that A is of the SJ type, $\ker(A) = \mathbb{C} V_p e_1,\ B = -\bar{V}_p e_1,\ b = -V_p e_1,\ C = -\sqrt{1 - B^T b}$ and further the block of $V_p e_1$ in A is $-J_p$ (the definitions of $V_p,\ J_p$ as $V_{2k} := [f_1 ... f_k\ \bar{f}_k ... \bar{f}_1],\ k \geq 1,\ V_{2k+1} := [f_1 ... f_k\ e_{2k+1}\ \bar{f}_k ... \bar{f}_1],\ k \geq 0,\ J_p := V_p I'_{1,p-1} \bar{V}_p^T,\ p \geq 1$ can be extended to $V_0 := \emptyset,\ J_0 := \emptyset$ with the convention that matrices whose either dimension is 0 are empty and are excluded from the block decomposition, $V_0 e_1 = 0 = 0_{n,1}$ as the span of $\emptyset$). Further we require that the block of $a_1,\ \mathcal{S}(x_{a_1}) = H^{-1}(C(\infty))$ in $A^{\circ -1}$ is $a_1 I_q - J_q,\ q \geq 1$, so $A^{\circ -1} = \begin{bmatrix} -\bar{J}_p & 0 & 0 \\ 0 & a_1 I_q - J_q & 0 \\ 0 & 0 & a_1 I_r - A'^{-1} \end{bmatrix} =: (-\bar{J}_p) \oplus (a_1 I_q - J_q) \oplus (a_1 I_r - A'^{-1}),\ p, r \geq 0,\ q \geq 1,\ p+q+r = n$ and $A'^{-1} := \bigoplus_{j=2}^m ((a_1 - a_j) I_{s_j} + \bigoplus_k J_{s_{jk}}) \in \mathbf{GL}_r(\mathbb{C}),\ \Sigma_k s_{jk} = s_j > 0,\ j = 2, ..., m,\ \Sigma_j s_j = r$. To simplify the notation we drop the $\oplus$ for zero terms: for example $A J_q = (a_1 I_q - J_q)^{-1} J_q$.

Quadrics whose eigenspaces of all non-zero eigenvalues of A have dimension ≥ 2 do not enjoy the property of their confocal family being taken to another one; for example QC with $A = a^{-1} I_n$, whose only finite singular quadric (namely the isotropic cone $|x|^2 = 0$) is not projectively equivalent to $\mathbb{CP}^{n-1}$) or IQWC with $A = J_n$ (in which case there are no finite singular quadrics).

Conversely, given x_z as above, then since $\mathcal{S}(x_{a_1})$ is projectively equivalent to $C(\infty)$, as a quadric in a hyperplane of $\mathbb{CP}^n$, there exists H homography (unique up to rigid motions in the domain of H which fix $\mathcal{S}(x_{a_1})$ and rigid motions and homotheties in the range of H) such that $H(\mathcal{S}(x_{a_1})) = C(\infty)$. H takes x_z to another family of confocal quadrics $\tilde{x}_{\tilde{z}}$ (since $\tilde{x}_{\tilde{z}}$ is uniquely determined by $C(\infty)$ and $\mathcal{S}(\tilde{x}_{\tilde{a}_1}) := H(C(\infty))$).

Since $\mathcal{S}(x_{a_1}) = H^{-1}(C(\infty)) :\ |H' x + h_1|^2 = h_2^T x + a = 0$ we have $a = 0,\ h_2 = e^{c_1} V_q e_1$. If $q = 1$, then $V_q e_1 = e_{p+1},\ x_z \cap \{e_{p+1}^T x = 0\}$ is a quadric in $e_{p+1}^T x = 0$ and is taken by H into $\tilde{x}_{\tilde{z}} \cap \mathbb{CP}^{n-1}$, which thus must be a quadric and therefore $\tilde{x}_{\tilde{z}}$ are QC; if $q \geq 2$, then $x_z \cap \{(V_q e_1)^T x = 0\}$ is a degenerate quadric in $(V_q e_1)^T x = 0$ (a quadratic cone with vertex $[(V_q e_1)^T, 0] \in C(\infty)$, since the quadric x_z is tangent to $(V_q e_1)^T x = 0$ at $[(V_q e_1)^T, 0]$) and is taken by H into $\tilde{x}_{\tilde{z}} \cap \mathbb{CP}^{n-1}$, which thus must be a degenerate quadric and therefore $\tilde{x}_{\tilde{z}}$ are (I)QWC, according to $[(V_q e_1)^T, 0] \notin \mathcal{S}(x_{a_1}) (\in \mathcal{S}(x_{a_1})) \Leftrightarrow (V_q e_1)^T R_{a_1}^{\circ -1} (V_q e_1) \neq 0\ (= 0) \Leftrightarrow (V_q e_1)^T \bar{J}_q (V_q e_1) \neq 0\ (= 0) \Leftrightarrow q = 2\ (q > 2)$. Moreover vertices of importance to the characterization of quadrics transform under H as follows: $H[(V_q e_1)^T, 0] = [(V_{\tilde{p}} e_1)^T, 0]\ (= [0^T, 1])$ for $\tilde{p} > 0\ (\tilde{p} = 0),\ q = \tilde{p} + 1$ and H' splits as $H' = I_{1,n-r} H' I_{1,n-r} + H'',\ 0_{n-r,n-r} \oplus H'' \simeq H'' \in \mathbf{GL}_r(\mathbb{C}),\ \tilde{H}'' = H''^{-1}$. As $H'' H''^T = \bigoplus_{j=2}^m ((\tilde{a}_1 - \tilde{a}_j) I_{s_j} + \bigoplus_k J_{s_{jk}}),\ H''^T H'' = \bigoplus_{j=2}^m \bigoplus_k ((a_1 - a_j) I_{s_j} + J_{s_{jk}})^{-1},\ H''$ is formed by

blocks of the form $R_1(\sqrt{(a_1-a_j)^{-1}}I_{s_{jk}}+J_{s_{jk}})R_2^T$, where the rotations $R_1,\ R_2\in\mathbf{O}_{s_{jk}}(\mathbb{C})$ can be found from a diagonalization process: $R_1(2\sqrt{(a_1-a_j)^{-1}}J_{s_{jk}}+J_{s_{jk}}^2)R_1^T=R_2((\sqrt{(a_1-a_j)^{-1}}I_{s_{jk}}+J_{s_{jk}})^{-2}-(a_1-a_j)I_{s_{jk}})R_2^T=J_{s_{jk}}$. As expected, excluding the singular values $a_1,\ \infty$ of z and the corresponding singular values $\infty,\ \tilde{a}_1$ of $\tilde{z}$, the remaining singular values of $z,\ \tilde{z}$ not only correspond $(\tilde{a}_j=\tilde{a}_1+(a_j-a_1)^{-1},\ j=2,...,m)$, but also their singular quadrics are projectively equivalent (with the projective structure induced by their singular sets as previously described).

We shall now provide an analytic confirmation of these facts: we have $H'\tilde{H}'+h_1\tilde{h}_2^T=I_n,\ H'\tilde{h}_1=\tilde{H}'^Th_2=0,\ h_2^T\tilde{h}_1=1$. Multiplying the first relation on the left by $\ker(H'^T)^T$ and on the right by $\ker(\tilde{H}')$ we get $\ker(H'^T)=\mathbb{C}\tilde{h}_2,\ \ker(\tilde{H}')=\mathbb{C}h_1$; also $\tilde{a}_1I_n-\tilde{A}^{\circ-1}=H'H'^T,\ H'h_2=-\tilde{b},\ |h_2|^2=-\tilde{C};\ 0=H'H'^T\tilde{h}_2=\tilde{h}_2-\tilde{A}^{\circ-1}\tilde{h}_2$, so $\tilde{h}_2=e^{\tilde{c}_1}V_{\tilde{q}}e_1;\ -\tilde{H}'\tilde{b}=\tilde{H}'H'h_2=h_2+\tilde{C}\tilde{h}_1,\ -\tilde{H}'^T\tilde{H}'\tilde{b}=\tilde{C}\tilde{H}'^T\tilde{h}_1=0$, so $\tilde{h}_1=h_2=e_{p+1}$ for $\tilde{x}_{\tilde{z}}$ QC; $|h_1|^2bb^T=\tilde{H}'((H'\tilde{H}')^TH'\tilde{H}'-I_n)\tilde{H}'^T=(A^{\circ-1}-a_1I_n)H'^TH'(A^{\circ-1}-a_1I_n)+(A^{\circ-1}-a_1I_n)$. Multiplying it on the left by $R_{a_1}^{\circ-1}A$, on the right by $AR_{a_1}^{\circ-1}$ and using $H'^TH'b=0$ we get $R_{a_1}^{\circ-1}R_{a_1}H'^TH'R_{a_1}R_{a_1}^{\circ-1}=-R_{a_1}^{\circ-1}AR_{a_1}R_{a_1}^{\circ-1}$; further multiplying on the left by $I_n-h_2\tilde{h}_1^T$ and on the right by $I_n-\tilde{h}_1h_2^T$ we get $H'^TH'=-(I_n-h_2\tilde{h}_1^T)R_{a_1}^{\circ-1}AR_{a_1}R_{a_1}^{\circ-1}(I_n-\tilde{h}_1h_2^T)$. If x_z are (I)QWC, then $0=(I_n-h_1\tilde{h}_2^T-H'\tilde{H}')\tilde{H}'^TB=(\tilde{H}'^T+h_1b^T-a_1H')B=h_1+(\tilde{H}'^T-a_1H')B$, so $H'^Th_1=(a_1H'^TH'+h_2\tilde{h}_1^T-I_n)B,\ R_{a_1}H'^Th_1=-R_{a_1}(a_1AR_{a_1}^{\circ-1}+I_n)B=-B,\ -|h_1|^2=-|(a_1H'-\tilde{H}'^T)B|^2=a_1B^T(I_n-a_1H'^TH')B=a_1B^TR_{a_1}^{\circ-1}B;\ -\tilde{H}'^TR_{a_1}^{\circ-1}B=\tilde{H}'^TR_{a_1}^{\circ-1}R_{a_1}H'^Th_1=\tilde{H}'^TH'^Th_1$, so $h_1=-a_1B^TR_{a_1}^{\circ-1}B\tilde{h}_2-\tilde{H}'^TR_{a_1}^{\circ-1}B$.

The equalities $R_{a_1}^{\circ-1}R_{a_1}H'^TH'R_{a_1}R_{a_1}^{\circ-1}=-R_{a_1}^{\circ-1}AR_{a_1}R_{a_1}^{\circ-1},\ R_{a_1}H'^Th_1=-B,\ |h_1|^2=-C-a_1B^TR_{a_1}^{\circ-1}B$ can also be found by accounting the two equations of $H^{-1}(C(\infty))=\mathcal{S}(x_{a_1}):\ |H'x+h_1|^2=h_2^Tx=0$ and $x^T(AR_{a_1}^{\circ-1}x+2R_{a_1}^{\circ-1}B)+a_1B^TR_{a_1}^{\circ-1}B+C=0,\ x=R_{a_1}R_{a_1}^{\circ-1}x$.

Rewriting the above: $h_2=e^{c_1}V_qe_1,\ \ker(H')=\mathbb{C}\tilde{h}_1,\ \ker(H'^T)=\mathbb{C}V_{\tilde{q}}e_1,\ H'V_pe_1=Ch_1+e^{\tilde{c}_1}V_{\tilde{q}}e_1,\ H'V_qe_1=e^{-c_1}V_{\tilde{p}}e_1,\ (I_n-V_qe_1e_1^T\bar{V}_q^T)H'^Th_1=(I_p+a_1J_p)^{-1}\bar{V}_pe_1,\ |h_1|^2=(-a_1)^p,\ H'H'^T=(\tilde{a}_1I_{\tilde{p}}+\bar{J}_{\tilde{p}})\oplus J_{\tilde{q}}\oplus\tilde{A}'^{-1},\ (I_n-V_qe_1e_1^T\bar{V}_q^T)H'^TH'(I_n-\bar{V}_qe_1e_1^TV_q^T)=(I_p+a_1J_p)^{-1}J_p\oplus\bar{J}_q\oplus A',\ h_1=(-a_1)^pe^{\tilde{c}_1}V_{\tilde{q}}e_1+\tilde{H}'^T(I_p+a_1J_p)^{-1}\bar{V}_pe_1$.

We have $\ker((H'H'^T)^j)\subseteq\ker(H'^T(H'H'^T)^j)\subseteq\ker((H'H'^T)^{j+1}),\ \ker((H'^TH')^j)\subseteq\ker(H'(H'^TH')^j)\subseteq\ker((H'^TH')^{j+1}),\ (H'^T(H'H'^T)^j)^T=H'(H'^TH')^j,\ j\geq 0$, so $\dim(\ker((H'H'^T)^j))\leq\dim(\ker((H'^TH')^{j+1}))\leq\dim(\ker((H'H'^T)^{j+2})),\ j\geq 0$. For $M=M^T\in\mathbf{M}_n(\mathbb{C})$ the function $0\leq j\mapsto\dim(\ker(M^j))$ is increasing and stabilizes to the sum of the dimensions of the SJ blocks of $\ker(RMR^T)$ for $j\geq k,\ RMR^T$ being the SJ form of M and k the maximum dimension of the SJ blocks of $\ker(RMR^T)$. Since $\ker(H'^TH')=\mathbb{C}\tilde{h}_1\oplus\mathbb{C}b,\ |\tilde{h}_1|^2\neq 0$ we have $\dim(\ker((H'^TH')^j))=p+1,\ j\geq p+\delta_{p0}$; since $\dim(\ker((H'H'^T)^j))=\tilde{q},\ j\geq\tilde{q}$ we have $\tilde{q}=p+1$.

If $\tilde{x}_{\tilde{z}}$ are QC, then $\tilde{p}=0,\ q=1,\ \ker(H'^T)=\mathbb{C}V_{p+1}e_1,\ \ker(H')=\mathbb{C}e_{p+1}$, so $H'H'^T=J_{p+1}\oplus\tilde{A}'^{-1},\ H'^TH'=(I_p+a_1J_p)^{-1}J_p\oplus A'$; moreover since $H'(H'^TH')^{p+1}=(H'H'^T)^{p+1}H',\ H'$ splits as $H'=I_{1,p+1}H'I_{1,p+1}+H'',\ H''\in\mathbf{GL}_r(\mathbb{C})$. With $V_{p+1}^TH'V_p=:[0_{p,1}\ \hat{I}_pU]^T,\ U\in\mathbf{GL}_p(\mathbb{C})$ we have $U^T\hat{I}_pU=\hat{I}_p,\ UI'_p\hat{I}_pU^T=(I_p+a_1I'_p)^{-1}I'_p\hat{I}_p,\ U^Te_p=e^{\tilde{c}_1}e_p$; the second equality can be replaced by $I'_pU=(I_p+a_1I'_p)UI'_p$. With $R:=V_p\hat{I}_pU^TV_p^T+I_{p+1,n}$ we have $R\in\mathbf{O}_n(\mathbb{C}),\ J_p=R(I_p+a_1J_p)^{-1}J_pR^T$, so R appears from a diagonalization process and thus exists and is mostly unique; we can do better and actually compute U. With $e_j^TUe_k=:(-a_1)^{k-j}u_{jk}$ we have $\tilde{c}_1=0,\ U-I_p$ strictly upper triangular and

$$(I)\quad \Sigma_{s=0}^r u_{(p+1-j)-s\ (p+1-j)}u_{j+s\ j+r}=\delta_{r0},\ j,j+r=1,...,p,$$

$$(II)\quad u_{j+1\ j+r+1}+u_{j+1\ j+r}-u_{j\ j+r}=0,\ j,j+r=1,...,p-1.$$

$u'_{j\ j+r}:=\binom{\frac{p}{2}-j}{r},\ j,j+r>0$ satisfy (II) and (I): $\Sigma_{s=0}^r u'_{(p+1-j)-s\ (p+1-j)}u'_{j+s\ j+r}=\binom{\frac{p}{2}-j-1}{r}\Sigma_{s=0}^r(-1)^s\binom{r}{s}=\delta_{r0}$. Suppose $u_{j\ j+s}=u'_{j\ j+s}$ for $j,j+s=1,...,p,\ s=0,...,r-1$ and for $s=r$ and some $j=1,...,p-r$; from (II) it is true for $j,j+s=1,...,p,\ s=0,...,r$. Thus in order

to prove that $u_{jk}=u'_{jk},\ j,k=1,...,p$ we need: $u_{j\ j+s}=u'_{j\ j+s}$ for $j,j+s,r+1=1,...,p,\ s=0,...,r-1\Rightarrow u_{j\ j+r}=u'_{j\ j+r}$ for some $j=1,...,p-r$. For $p=2k$ if $r=2m-1$, then let $j=k-m+1$ in (I) to get $u_{k-m+1\ k+m}=u'_{k-m+1\ k+m}$; if $r=2m$, then let $j=k-m$ in (I) and (II) to get $u_{k-m\ k+m}=u'_{k-m\ k+m}$; for $p=2k-1$ if $r=2m-1$, then let $j=k-m$ in (I) and (II) to get $u_{k-m+1\ k+m}=u'_{k-m+1\ k+m}$; if $r=2m$, then let $j=k-m$ in (I) to get $u_{k-m\ k+m}=u'_{k-m\ k+m}$. Thus $U=U(p,a_1):=\Sigma_{j=1}^p e_je_j^T(I_p-a_1I'_p)^{\frac{p}{2}-j},\ U^{-1}=\Sigma_{j=1}^p e_je_j^T(I_p+a_1I'_p)^{\frac{p}{2}-j}$ (we have $\Sigma_{r=0}^l(-1)^{l-r}\binom{\frac{p}{2}-j}{r}\binom{\frac{p}{2}-j-r}{l-r}=\Sigma_{r=0}^l\binom{\frac{p}{2}-j}{r}\binom{j-\frac{p}{2}+l-1}{l-r}=\delta_{l0}$ as the coefficient of x^l in $(1+x)^{\frac{p}{2}-j}(1+x)^{j-\frac{p}{2}+l-1}=(1+x)^{l-1}$). Now $H'=V_{p+1}[\hat{I}_p\ 0_{p,1}]^TU^TV_p^T+H'',\ h_1=V_{p+1}(I_{p+1}-a_1I'_{p+1})^{\frac{p}{2}}e_{p+1}$, $\tilde{H}'=V_p[\hat{I}_p\Sigma_{j=1}^p a_1^j\binom{\frac{p}{2}}{j}e_j\quad U\hat{I}_p]V_{p+1}^T+H''^{-1}$. Note that $e_j^T(U\hat{I}_p)e_k=a_1^{p+1-j-k}\binom{\frac{p}{2}-k}{p+1-j-k},\ j,k=1,...,p$; for $k=0$ we get the first column $e_j^T(\hat{I}_p\Sigma_{l=1}^p a_1^l\binom{\frac{p}{2}}{l}e_l)=a_1^{p+1-j}\binom{\frac{p}{2}}{p+1-j},\ j=1,...,p$ of $V_p^T\hat{I}_p\tilde{H}'\hat{I}_{p+1}V_{p+1}$ and for $j=0$ we get the first p entries $e_k^T(I_{p+1}-a_1I'_{p+1})^{\frac{p}{2}}e_{p+1}=(-a_1)^{p+1-k}\binom{\frac{p}{2}}{p+1-k},\ k=1,...,p$ of $V_{p+1}^T\hat{I}_{p+1}h_1$.

If $x_z,\ \tilde{x}_{\tilde{z}}$ are (I)QWC, then $p,\ \tilde{p}>0,\ H'H'^TV_{\tilde{p}}=V_{\tilde{p}}(\tilde{a}_1I_{\tilde{p}}+I'^T_{\tilde{p}}),\ H'^TH'V_{\tilde{p}+1}I_{1,\tilde{p}}=V_{\tilde{p}+1}(e_1v^T+I'^T_{\tilde{p}}),\ v^T:=e_{\tilde{p}+1}^TV_{\tilde{p}+1}^TH'^TH'V_{\tilde{p}+1}I_{1,\tilde{p}}$. By induction after $j\geq 0$ we have $H'V_{\tilde{p}+1}(e_1v^T+I'^T_{\tilde{p}+1})^je_1=e^{-c_1}V_{\tilde{p}}(\tilde{a}_1I_{\tilde{p}}+I'^T_{\tilde{p}})^je_1,\ j=0,...,\tilde{p},\ V_{\tilde{p}+1}(e_1v^T+I'^T_{\tilde{p}+1})^{j+1}e_1=e^{-c_1}H'^TV_{\tilde{p}}(\tilde{a}_1I_{\tilde{p}}+I'^T_{\tilde{p}})^je_1,\ j=0,...,\tilde{p}-1$; thus $H'^TV_{\tilde{p}}=V_{\tilde{p}+1}(\bar{V}_{\tilde{p}+1}^TH'^TV_{\tilde{p}})$ and $\tilde{h}_1=V_{\tilde{p}+1}((-\tilde{a}_1)^{\tilde{p}}e^{c_1}e_1+\bar{V}_{\tilde{p}+1}^TH'^TV_{\tilde{p}}(I_{\tilde{p}}+\tilde{a}_1I'_{\tilde{p}})^{-1}e_{\tilde{p}})$, $H'^TH'=(I_p+a_1J_p)^{-1}J_p\oplus(I_{\tilde{p}+1}-h_2\tilde{h}_1^T)\bar{J}_{\tilde{p}+1}(I_{\tilde{p}+1}-\tilde{h}_1h_2^T)\oplus A'$. Since $H'(H'^TH')^{p+1}=(H'H'^T)^{p+1}H'$, we have $V_{\tilde{p}}^TH'V_p=0,\ I_rH'V_p=0,\ V_{p+1}^TH'I_r=0$; further since $H'(H'^TH')=(H'H'^T)H'$, we have $V_{p+1}^TH'V_{\tilde{p}+1}\in\mathbb{C}e_{\tilde{p}+1}\tilde{h}_1^TV_{\tilde{p}+1}$. As $V_{p+1}^TH'V_{\tilde{p}+1}e_1=e^{-c_1}V_{p+1}^TV_{\tilde{p}}e_1=0$ we have $V_{p+1}^TH'V_{\tilde{p}+1}=0$; now $V_{p+1}^TH'V_p=:[0_{p,1}\ \hat{I}_pU]^T,\ U\in\mathbf{GL}_p(\mathbb{C})$ must be as before, so $\tilde{c}_1=0,\ U=U(p,a_1)$. As $\tilde{H}'H'+\tilde{h}_1h_2^T=I_n$ we have $(V_{\tilde{p}+1}^T\tilde{H}'V_{\tilde{p}})\hat{I}_{\tilde{p}}(V_{\tilde{p}}^TH'I_r)=0,\ (I_r\tilde{H}'I_r)(I_rH'V_{\tilde{p}+1})=0,\ (I_r\tilde{H}'I_r)(I_rH'I_r)=I_r$, so $V_{\tilde{p}}^TH'I_r=0,\ I_rH'V_{\tilde{p}+1}=0$ and finally

$H'=V_{p+1}[\hat{I}_p\ 0_{p,1}]^TU(p,a_1)^TV_p^T+V_{\tilde{p}}[\hat{I}_{\tilde{p}}\Sigma_{j=1}^{\tilde{p}}\tilde{a}_1^j\binom{\frac{\tilde{p}}{2}}{j}e_j\quad U(\tilde{p},\tilde{a}_1)\hat{I}_{\tilde{p}}]V_{\tilde{p}+1}^T+H'',\ \tilde{h}_1=V_{\tilde{p}+1}(I_{\tilde{p}+1}-\tilde{a}_1I'_{\tilde{p}+1})^{\frac{\tilde{p}}{2}}e_{\tilde{p}+1}$.

A homography H does not have the property of taking a family of confocal quadrics to another iff it is an affine transformation with $H(C(\infty))\neq C(\infty)$ or $\begin{vmatrix}H'H'^T-\lambda I_n & H'h_2\\ (H'h_2)^T & |h_2|^2\end{vmatrix}=0,\ \forall\lambda\in\mathbb{C}$; in the later case $H^{-1}(C(\infty)):\ |H'x+h_1|^2=h_2^Tx+a=0$ (and thus $H(C(\infty))$) cannot appear as the singular set of a singular quadric of a family of confocal quadrics. By applying a rigid motion in the domain of H and fixing the constant upon which H depends we make $a=0,\ h_2=f_1\ (=e_1)$ according to being isotropic (or not) and bring $H^{-1}(C(\infty))$ to the form of (I)Q(W)C in $f_1^Tx=0\ (e_1^Tx=0):\ \ker(A)$ is spanned by $f_1,\ \{f_1,e_3\},\ \{f_1,f_2\}\ (e_1,\ \{e_1,e_3\},\ \{e_1,f_2\}),\ B=0,e_3,-\bar{f}_2,\ C=-1,0,0$ or a quadric of the form $f_1^Tx=x^T(Ax-2\bar{f}_1)=0,\ A=A^T,\ \ker(A)=I_{1,2}\mathbb{C}^n$. As we have seen, $H^{-1}(C(\infty))$ must be of this last type; thus $|h_1|^2=0,\ (I_n-f_1\bar{f}_1^T)H'^Th_1=-e^c\bar{f}_1,\ H'^TH'f_1=c_1f_1,\ |H'f_1|^2=0$; by applying a rotation in the range of H we make $h_1=e^cf_1$, so $H'^Tf_1=-\bar{f}_1+c_2f_1,\ f_1^TH'f_1=-1$. Since $\{R(h)|h=I_{3,n}h\in\mathbb{C}^n\}$ acts transitively on $\{x\in\mathbb{C}^n|\ |x|^2=0,\ f_1^Tx=-1\}$, by applying a rotation $R(h)$ in the range of H we further make $H'f_1=-\bar{f}_1$, so $H'^T\bar{f}_1=-c_1f_1,\ H'=(f_1\bar{f}_1^T+\bar{f}_1f_1^T+I_{3,n})H'(f_1\bar{f}_1^T+\bar{f}_1f_1^T+I_{3,n})=h_3f_1^T-\bar{f}_1\bar{f}_1^T+I_{3,n}H'',\ h_3\in\mathbb{C}^n,\ H''\in\mathbf{GL}_n(\mathbb{C}),\ I_{3,n}H''=H''I_{3,n}=H''-I_{1,2}$; further applying the translation $-h_3$ in the range of H we make $h_3=0,\ H'H'^T=I_{3,n}H''H''^T,\ \begin{vmatrix}H'H'^T-\lambda I_n & H'h_2\\ (H'h_2)^T & |h_2|^2\end{vmatrix}=\begin{vmatrix}I_n & -\lambda f_1\\ 0^T & 1\end{vmatrix}\begin{vmatrix}I_{3,n}H''H''^T-\lambda I_n & -\bar{f}_1\\ -\bar{f}_1^T & 0\end{vmatrix}\begin{vmatrix}I_n & 0\\ -\lambda f_1^T & 1\end{vmatrix}=\begin{vmatrix}I_{3,n}H''H''^T-\lambda I_{3,n} & -\bar{f}_1\\ -\bar{f}_1^T & 0\end{vmatrix}=0.$

Summing up:

Any homography $H := \begin{bmatrix} H' & h_1 \\ h_2^T & a \end{bmatrix} \in \mathbf{PGL}_n(\mathbb{C}),\ H' \in \mathbf{M}_n(\mathbb{C}),\ h_1,\ h_2 \in \mathbb{C}^n,\ a \in \mathbb{C}$, *with inverse* $(H^{-1} =)\tilde{H} := \begin{bmatrix} \tilde{H}' & \tilde{h}_1 \\ \tilde{h}_2^T & \tilde{a} \end{bmatrix}$ *can be brought, via rigid motions and homotheties in both its domain and range, to one of the two cases below:*

(I) (i) $H = \tilde{H} = I_{n+1}$*; (ii)* $a = \tilde{a} = 1,\ h_1 = \tilde{h}_1 = h_2 = \tilde{h}_2 = 0,\ \tilde{H}'^{-1} = H' \in \mathbf{GL}_n(\mathbb{C})\backslash\{\mathbb{C}I_n\}$ *SJ; (iii)* $a = \tilde{a} = 0,\ e^{-c}h_1 = h_2 = f_1,\ \tilde{h}_1 = e^c\tilde{h}_2 = \bar{f}_1,\ H' = -\bar{J}_2 + I_{3,n}H'',\ \tilde{H}' = -J_2 + I_{3,n}H''^{-1},\ c \in \mathbb{C},\ H'' \in \mathbf{GL}_n(\mathbb{C})$ *SJ,* $I_{3,n}H'' = H'' - I_{1,2}$.

(II) $a = \tilde{a} = 0$ *and* H *takes confocal (I)Q(W)C* x_z *to confocal (I)Q(W)C* $\tilde{x}_{\tilde{z}},\ \tilde{a}_1 := a_1^{-1},\ \tilde{z} := \tilde{a}_1 + (z - a_1)^{-1}$*. By symmetry all the following remain true if tilde and non-tilde quantities are exchanged: with* $U(p, a_1) := \Sigma_{j=1}^{p} e_j e_j^T (I_p - a_1 I'_p)^{\frac{p}{2}-j}$ *we have* $h_1 = V_{p+1}(I_{p+1} - a_1 I'_{p+1})^{\frac{p}{2}} e_{p+1},\ h_2 = V_{\tilde{p}+1}e_1,\ H' = V_{p+1}[\hat{I}_p\ 0_{p,1}]^T U(p, a_1)^T V_p^T + V_{\tilde{p}}[(\hat{I}_{\tilde{p}}\Sigma_{j=1}^{\tilde{p}} \tilde{a}_1^j (\frac{\frac{\tilde{p}}{2}}{j}) e_j)\ U(\tilde{p}, \tilde{a}_1)\hat{I}_{\tilde{p}}]V_{\tilde{p}+1}^T + H'',\ H'' \in \mathbf{GL}_{n-p-\tilde{p}+1}(\mathbb{C}),\ \tilde{H}'' := H''^{-1},\ H''H''^T = \bigoplus_{j=2}^{m}((\tilde{a}_1 - \tilde{a}_j)I_{s_j} + \bigoplus_k J_{s_{jk}}),\ H''^T H'' = \bigoplus_{j=2}^{m}\bigoplus_k((a_1 - a_j)I_{s_{jk}} + J_{s_{jk}})^{-1},\ \Sigma_k s_{jk} = s_j > 0,\ j = 2, ..., m,\ \Sigma_j s_j = n - p - \tilde{p} - 1,\ \tilde{a}_j = \tilde{a}_1 + (a_j - a_1)^{-1},\ j = 2, ..., m.$

Thus H'' *is formed by blocks of the form* $R_1(\sqrt{(a_1 - a_j)^{-1}}I_{s_{jk}} + J_{s_{jk}})R_2^T$*, where the rotations* $R_1,\ R_2 \in \mathbf{O}_{s_{jk}}(\mathbb{C})$ *can be found from a diagonalization process:* $J_{s_{jk}} = R_1(2\sqrt{(a_1 - a_j)^{-1}}J_{s_{jk}} + J_{s_{jk}}^2)R_1^T = R_2((\sqrt{(a_1 - a_j)^{-1}}I_{s_{jk}} + J_{s_{jk}})^{-2} - (a_1 - a_j)I_{s_{jk}})R_2^T$.

Also $A^{\circ -1} = \bar{J}_p \oplus (a_1 I_{\tilde{p}+1} - J_{\tilde{p}+1}) \oplus \bigoplus_{j=2}^{m}\bigoplus_k (a_j I_{s_{jk}} - J_{s_{jk}}),\ B = -\bar{V}_p e_1,\ C = -\delta_{p0}$.

Conversely, given a quadric in a hyperplane of $\mathbb{C}^n$*, excluding quadrics which belong, after a rigid motion, to the type* $f_1^T x = x^T(Ax - 2\bar{f}_1) = 0,\ A$ *SJ,* $\ker(A) = I_{1,2}\mathbb{C}^n$*, one can consider it to be* $\mathcal{S}(x_{a_1})$ *and complete it to a family of confocal quadrics* x_z*; then one can recover* $H,\ \tilde{H}$ *and* $\tilde{x}_{\tilde{z}}$ *as above.*

2.4. **Parametrization of confocal isotropic quadrics without center.**
Consider the equilateral QWC $Z^T(I_{1,n-1}Z - 2e_n) = 0$ and a canonical IQWC $x_0^T(Ax_0 - 2\bar{f}_1) = 0,\ \ker(A) = \mathbb{C}f_1,\ A = J_p \oplus ...$ SJ. We are looking for a linear map $L \in \mathbf{GL}_n(\mathbb{C})$ such that $x_0 = LZ$, equivalently $L^T A L = e^{2a}I_{1,n-1},\ L^T\bar{f}_1 = e^{2a}e_n$. Replacing L with $L(e^{-a}I_{1,n-1} + e^{-2a}e_n e_n^T)$ we can make $a = 0$. Thus $Le_n = f_1,\ L^T(A + \bar{f}_1\bar{f}_1^T)L = I_n$, so $L^{-1} = R^T\sqrt{A + \bar{f}_1\bar{f}_1^T},\ R^T R = I_n$ with $Re_n = \sqrt{A + \bar{f}_1\bar{f}_1^T}f_1$ (note that Re_n has, as required, length 1). Once $R \in \mathbf{O}_n(\mathbb{C})$ with the above property is found, L thus defined satisfies $L^T\bar{f}_1 = e_n$ and thus $L^T A L = I_{1,n-1}$. L with the above properties is unique up to rotations fixing e_n in its domain and a canonical choice reveals itself: $I_{1,n-1}L^{-1}\sqrt{R_z}LI_{1,n-1} = \sqrt{I_{1,n-1} - zL^T A^2 L} =: \sqrt{R'_z} - e_n e_n^T,\ A' := L^T A^2 L,\ \ker(A') = \mathbb{C}e_n \oplus \mathbb{C}L^{-1}A^{\circ -1}f_1 = \mathbb{C}e_n \oplus \mathbb{C}L^T f_1$; choose R which makes A' SJ. We have $LL^T = A^{\circ -1} + f_1 f_1^T,\ L^{-1}\sqrt{R_z}Le_n = e_n$, so $I_{1,n-1}\sqrt{R'_z} = I_{1,n-1}L^{-1}\sqrt{R_z}L = L^T A\sqrt{R_z}L$.

Consider now the ruling families parametrization on Z for $n = 2k + 1:\ u_j := \bar{f}_j^T Z,\ v_j := f_j^T Z,\ j = 1, ..., k$; thus $Z = Z(u_j, v_j) = \Sigma_{j=1}^{k}(u_j f_j + v_j\bar{f}_j + u_j v_j e_n)$; if we let $u_k = v_k$, then the $2k^{\text{th}}$ coordinate is 0 and we obtain the ruling families parametrization on Z for $n = 2k$. Let $e_n^T L^{-1}\sqrt{R_z}L =: \Sigma_{j=1}^{k}(u_j(z)f_j + v_j(z)\bar{f}_j)^T + e_n^T,\ u_{jz} := u_j + u_j(z),\ v_{jz} := v_j + v_j(z),\ Z^z = Z^z(u_j, v_j) := Z(u_{jz}, v_{jz}),\ Z^z(0) := Z^z(0_j, 0_j) = I_{1,n-1}L^T\sqrt{R_z}(L^T)^{-1}e_n + \frac{1}{2}|I_{1,n-1}L^T\sqrt{R_z}(L^T)^{-1}e_n|^2 e_n$. We have $I_{1,n-1}\sqrt{R'_z}Z^z(0) = L^T\sqrt{R_z}AA^{\circ -1}\sqrt{R_z}(L^T)^{-1}e_n = L^T(I_n - \sqrt{R_z} - zA)\bar{f}_1,\ e_n^T Z^z(0) = \frac{1}{2}\bar{f}_1^T\sqrt{R_z}A^{\circ -1}\sqrt{R_z}\bar{f}_1;\ -2\partial_z(\sqrt{R_z}A^{\circ -1}\sqrt{R_z}) = \sqrt{R_z}(R_z^{-1}AA^{\circ -1} + A^{\circ -1}AR_z^{-1})\sqrt{R_z} = 2I_n - (\sqrt{R_z})^{-1}\bar{f}_1 f_1^T - f_1\bar{f}_1^T(\sqrt{R_z})^{-1}$, so $e_n^T Z^z(0) = \bar{f}_1^T C(z)$; further $Z^z = Z + (e_n Z^T I_{1,n-1} + I_n)Z^z(0)$ and $L^{-1}x_z - \sqrt{R'_z}Z^z + 2zI_{1,n-1}\partial_z|_{z=0}Z^z(0) = L^{-1}C(z) - \sqrt{R'_z}Z^z(0) - zL^T A(L^T)^{-1}e_n = (L^{-1} - e_n\bar{f}_1^T)C(z) - L^T(I_n - \sqrt{R_z} - zA)\bar{f}_1 - zL^T A\bar{f}_1 = 0$ (use $L^{-1} = L^T A + e_n\bar{f}_1^T,\ AC(z) - I_n - \sqrt{R_z})\bar{f}_1 = 0$). Thus $x_z = L\sqrt{R'_z}Z^z + z\bar{f}_1$; with $f := e_n\ (\bar{f}_1)$ respectively for (I)QWC and $A' := A,\ L := (\sqrt{A + e_n e_n^T})^{-1}$ for QWC we have a parametrization for (I)QWC: $x_z = L\sqrt{R'_z}Z^z + z(1 - \frac{|f|^2}{2})(L^T)^{-1}e_n$. The TC becomes: $(V_0^1)^T\hat{N}_0^0 = Z_0^T I_{1,n-1}\sqrt{R'_z}Z_1 - (Z_0 +$

$Z_1)^T(I_{1,n-1}Z^z(0)+e_n)-e_n^TZ^z(0)-\frac{|f|^2}{2}z=0$, so IQWC can be considered metrically degenerated QWC.

2.5. **Totally real quadrics in $\mathbb{C}^n$.**
Note that if $V\subseteq\mathbb{C}^n$ is a real k-dimensional subspace (a-priori $0\leq k\leq 2n$) such that the Euclidean scalar product on $\mathbb{C}^n$ induces a non-degenerate real (valued) scalar product on V, then the orthogonal basis of V (formed by vectors of length ± 1) provided by Sylvester's is, by complexification, an orthogonal base of a (complex) k-dimensional subspace of $\mathbb{C}^n$, so $k\leq n$.

Consider a real n-dimensional subspace $V\subset\mathbb{C}^n$ as above and a basis $\{v_j\}_{j=1,...,n}$ of V with $v_j^Tv_k=\epsilon_j\delta_{jk},\ \epsilon_j:=\pm 1,\ j=1,...,n,\ \mathcal{E}:=\mathrm{diag}[\epsilon_1\ ...\ \epsilon_n]$. Now $R:=[v_1\ ...\ v_n]\sqrt{\mathcal{E}}\in\mathbf{O}_n(\mathbb{C})$ and $R^TV=\sqrt{\mathcal{E}}\mathbb{R}^n$ satisfies $\overline{x}=\mathcal{E}x,\ \forall x\in R^TV$.

We are interested in the quadrics which intersect a totally real subspace $\sqrt{\mathcal{E}}\mathbb{R}^n$ along an $(n-1)$-dimensional totally real quadric and bringing such quadrics to the canonical form (a form depending on as few as possible constants) by means of rigid motions preserving $\sqrt{\mathcal{E}}\mathbb{R}^n$. The rigid motions preserving $\sqrt{\mathcal{E}}\mathbb{R}^n$ are $(R,t),\ R\in\mathbf{O}_n(\mathbb{C}),\ \bar{R}=\mathcal{E}R\mathcal{E},\ t\in\sqrt{\mathcal{E}}\mathbb{R}^n$; the homotheties preserving $\sqrt{\mathcal{E}}\mathbb{R}^n$ are $e^a,\ a\in\mathbb{R}$.

However in order to take advantage of the SJ form it is more convenient to intersect a quadric in canonical form with a totally real affine space $(R_0,t_0)\sqrt{\mathcal{E}}\mathbb{R}^n,\ (R_0,t_0)\in\mathbf{O}_n(\mathbb{C})\ltimes\mathbb{C}^n$ and put the condition that the sub-manifold thus obtained has dimension $n-1$. Note that (R_0,t_0) is uniquely determined modulo compositions on the right with rigid motions preserving $\sqrt{\mathcal{E}}\mathbb{R}^n$, so $R:=\bar{R}_0\mathcal{E}R_0^T,\ t:=t_0-R^T\bar{t}_0$ are uniquely determined. Note $R\in\mathbf{O}_n(\mathbb{C}),\ \bar{R}^T=R,\ \det R=\det\mathcal{E}$; conversely given R as above one can find R_0 as above (which is uniquely determined modulo multiplication on the right with rotations which preserve $\sqrt{\mathcal{E}}\mathbb{R}^n$). This is a standard dimensional argument: since the map $\mathbf{O}_n(\mathbb{C})\ni R_0\mapsto\bar{R}_0\mathcal{E}R_0^T\in\{R\in\mathbf{O}_n(\mathbb{C})|\bar{R}^T=R,\ \det R=\det\mathcal{E}\}$ is continuous with connected codomain of real dimension $\frac{n(n-1)}{2}$ and kernel of the same real dimension, it is onto. Similarly because of the real decomposition $\mathbb{C}^n\ni v=\frac{v+\mathcal{E}\bar{v}}{2}+\frac{v-\mathcal{E}\bar{v}}{2}\in\sqrt{\mathcal{E}}\mathbb{R}^n\oplus i\sqrt{\mathcal{E}}\mathbb{R}^n$, from t with $\bar{t}=-Rt$ one can obtain t_0 as above (which is unique modulo $R_0\sqrt{\mathcal{E}}\mathbb{R}^n$). Thus we need $\begin{bmatrix}R_0&t_0\\0&1\end{bmatrix}^T\begin{bmatrix}A&B\\B^T&C\end{bmatrix}\begin{bmatrix}R_0&t_0\\0&1\end{bmatrix}=e^{2c}\begin{bmatrix}\mathcal{E}&0\\0&1\end{bmatrix}\begin{bmatrix}\bar{R}_0&\bar{t}_0\\0&1\end{bmatrix}^T\begin{bmatrix}\bar{A}&\bar{B}\\\bar{B}^T&\bar{C}\end{bmatrix}\begin{bmatrix}\bar{R}_0&\bar{t}_0\\0&1\end{bmatrix}\begin{bmatrix}\mathcal{E}&0\\0&1\end{bmatrix}$; if such a condition is not satisfied, then applying the rigid motion $(R_0,t_0)^{-1}$ and using the real decomposition $\mathbb{C}^n=\sqrt{\mathcal{E}}\mathbb{R}^n\oplus i\sqrt{\mathcal{E}}\mathbb{R}^n$ wrt $\sqrt{\mathcal{E}}\mathbb{R}^n$ the considered sub-manifold becomes the intersection of two (possibly empty or degenerate) quadrics in $\sqrt{\mathcal{E}}\mathbb{R}^n$ and thus it has dimension at most $n-2$.

Applying the compatibility condition conjugation to the above relation we get $c\in i\mathbb{R}$; multiplying $\begin{bmatrix}A&B\\B^T&C\end{bmatrix}$ with e^{-c} we make $c=0$ (note however that by doing this B and C will change from their usual values for quadrics in canonical form).

Now $\bar{A}=RAR^T,\ \bar{B}=R(At+B),\ \bar{C}=t^T(B+R^T\bar{B})+C$; since $\bar{J}_p$ is obtained from J_p by applying reflections in the $e_{2j}^Tx=0$ hyperplanes (which take f_j to $\bar{f}_j$), multiplying R on the left with the product r of these reflections we obtain the rotation rR preserving the SJ form of a matrix and with a determinant and real condition. Thus the set of eigenvalues of A for SJ blocks of the same dimension is closed under conjugation, so eigenvalues of A, besides being either real or complex conjugate have corresponding SJ blocks of the same dimension. Moreover R exchanges the subspaces of $\mathbb{C}^n$ on which the SJ blocks of the eigenvalues a_j and $\bar{a}_j$ are supported, although it may not preserve the subspaces corresponding to the SJ blocks of same dimension, so R admits the corresponding block diagonal decomposition. If the corresponding blocks of $\mathcal{E}$ and of R have the same determinant, then one can choose R_0 with the same block decomposition, following that the blocks on which the determinants differ will be grouped by two to get a block for R_0; the simplest choice of R_0 will be the one giving so far the form closest to the canonical form; further simplifications will appear below. For the blocks $\bigoplus_{k=1}^{s_j}J_{p_k}$ corresponding to eigenvalue $a_j\in\mathbb{R}$ we have the corresponding block R_j of R with $R_j(\bigoplus_{k=1}^{s_j}J_{p_k})R_j^T=\bigoplus_{k=1}^{s_j}\bar{J}_{p_k}$ (which must still

satisfy the reality condition $\bar{R}_j^T = R_j$). For complex conjugate eigenvalues $a_j, \bar{a}_j \in \mathbb{C} \setminus \mathbb{R}$ we have the corresponding SJ block $(a_j I_l + \bigoplus_{k=1}^{s_j} J_{p_k}) \bigoplus (\bar{a}_j I_l + \bigoplus_{k=1}^{s_j} J_{p_k})$ of dimension $2l$; by considering $R(A - a_j I_n)^k R^T,\ k \geq 1$ we get $R_j = \begin{bmatrix} 0 & Q_j \\ \bar{Q}_j^T & 0 \end{bmatrix},\ Q_j \in \mathbf{O}_l(\mathbb{C})$ so we have reduced the problem to finding $Q_l \in \mathbf{O}_l(\mathbb{C})$ with $Q_l(\bigoplus_{k=1}^{s_j} J_{p_k})Q_l^T = \bigoplus_{k=1}^{s_j} J_{p_k}$ (without a reality condition).

Now we can already infer that most quadrics do not contain totally real quadrics and that totally real quadrics for which $\mathrm{spec}(A) \neq \{0\}$ belong to totally real confocal families obtained for $z \in \mathbb{R}$ (the necessity follows from the requirement on the eigenvalues of A; the sufficiency is clear from the definition of the confocal family). Note also that only for diagonal QC with $A = -\mathcal{E}|A|$ we have empty totally real quadrics, but this can be corrected by considering pairs $(\mathcal{E}, -\mathcal{E})$ instead of only $\mathcal{E}$. This pairing is natural, since multiplication by i exchanges both $\mathcal{E}$ to $-\mathcal{E}$ and the signature of the induced linear element of the totally quadric. According to Sylvester's this linear element has signature $(\frac{\mathrm{tr}(\mathcal{E}+|\mathcal{E}|)}{2}, \frac{\mathrm{tr}(\mathcal{E}-|\mathcal{E}|)}{2} - 1)$ along the regions where the normal is time-like and $(\frac{\mathrm{tr}(\mathcal{E}+|\mathcal{E}|)}{2} - 1, \frac{\mathrm{tr}(\mathcal{E}-|\mathcal{E}|)}{2})$ along the regions where the normal is space-like.

For QC we have $t = 0,\ C = \pm 1$; by a real homothety one can make $C = -1$.

For QWC we have $Re_n = \epsilon e_n,\ \epsilon := \pm 1,\ B = -e^{-c} e_n,\ e^{2c} = \epsilon,\ t = 0$.

For IQWC if the SJ block of f_1 is J_p, then R admits a corresponding block diagonal decomposition and on the first p coordinates R is $\pm$ the product of the reflections which take J_p to $\bar{J}_p$, so $Rf_1 = \epsilon \bar{f}_1,\ R\bar{f}_1 = \epsilon f_1,\ \epsilon := \pm 1,\ B = -e^{-c}\bar{f}_1,\ e^{2c} = \epsilon,\ t = 0$.

Therefore in all cases one can choose $t_0 := 0$ and the ambient totally real affine space passes through 0.

In order to get a classification of totally real quadrics and their confocal families we need to exclude cases obtained as above and which coincide. Since by complexification of two rigidly applicable totally real quadrics we obtain two rigidly applicable (complex) quadrics, two such totally real quadrics must belong to the same complex metric type; moreover the rigid motion by means of which the above rigidly applicability correspondence is realized must be a rigid motion which preserves the canonical form and conversely a rigid motion of the later type takes totally real quadrics to totally real quadrics. Such a rigid motion cannot preserve a totally real affine n-dimensional space, since a quadric intersects such a space in at most a totally real quadric. The group of rigid motions preserving the canonical form of a quadric acts transitively on each type of totally real quadrics in that quadric. Thus the last step in finding the canonical form of a totally real quadric is revealed from rotations preserving the SJ form: among all such rotations R' choose one which makes $\bar{R}' R R'^T$ as simple as possible and then continue with the program described above to find the simplest R_0.

Thus totally real quadrics naturally appear in canonical form not in totally real subspaces $\sqrt{\mathcal{E}}\mathbb{R}^n$, but in totally real subspaces $R_0\sqrt{\mathcal{E}}\mathbb{R}^n$. If one takes this to be the definition of the canonical form of totally real quadrics, then with the exception of finding the signature $\mathcal{E}$ one needs no work from a metric point of view beyond finding the canonical form of the corresponding (complex) quadric and checking the conditions on the eigenvalues and their corresponding SJ blocks.

3. Algebraic preparatives for quadrics in $\mathbb{C}^3$

3.1. **Canonical confocal quadrics and homographies of Bianchi II.**

In what follows we shall restrict our attention to quadrics in $\mathbb{C}^3$; we shall choose the ruling families parametrization invariant under the Ivory affinity. Intersections of isotropic rulings in $\mathbb{C}^3$ give all umbilics (because the first and second fundamental forms are proportional); also because of the preservation of lengths of segments on rulings under the Ivory affinity umbilics are preserved by the Ivory affinity. Excluding (pseudo-)spheres, the remaining four points of intersections of isotropic rulings (counting multiplicities) are situated on the circle at ∞, $C(\infty)$ (Bézout). When the four points are distinct we get general diagonal Q(W)C: $x_z \cap C(\infty) = x_z \pitchfork C(\infty)$; for higher multiplicities the quadrics are tangent to $C(\infty)$: $x_z \cap C(\infty) = (x_z \cap^T C(\infty)) \cup (x_z \pitchfork C(\infty))$. If

there are two points with double multiplicities, then we get diagonal Q(W)C and of revolution; in all other cases we get Darboux quadrics ($x_z \cap^T C(\infty)$ consists of one point); they cannot be realized as quadrics in $\mathbb{R}^3$, but if we intersect them with certain totally real spaces, then we get positive definite linear element so we get real deformations of these quadrics. If $x_z \cap \mathbb{CP}^2$ is non-degenerate, then x_z are QC; otherwise $x_z \cap \mathbb{CP}^2$ is a cone whose vertex belonging or not to $C(\infty)$ decides that x_z are IQWC or QWC.

Let $X \subset \mathbb{C}^3$ be the unit sphere $|X|^2 = 1$ and $u := \frac{e_3^T X+1}{f_1^T X},\ v := \frac{e_3^T X-1}{f_1^T X},\ u \neq v \in \mathbb{C} \cup \infty$ be the parametrization of X by its ruling families: $X = X(u,v) = \frac{-uvf_1+2\bar{f}_1+(u+v)e_3}{u-v}$. X has asymptotic isotropic cone $u = v = \tau:\ \infty Y(\tau),\ Y(\tau) := -\tau^2 f_1 + 2\bar{f}_1 + 2\tau e_3$. Consider $\sigma_1 : (u,v) \mapsto (iu, iv)$ and the involutions $\sigma_1^2,\ \sigma_2, \sigma_3, \sigma_4, \sigma_5 : (u,v) \mapsto (-u,-v),\ (v,u),\ (2u^{-1}, 2v^{-1}),\ (\bar{u}, \bar{v}),\ (\frac{2-\sqrt{2}u}{\sqrt{2}+u}, \frac{2-\sqrt{2}v}{\sqrt{2}+v})$; note the relations $\sigma_1\sigma_3 = \sigma_3\sigma_1^{-1},\ \sigma_1\sigma_4 = \sigma_4\sigma_1^{-1},\ \sigma_1\sigma_5 = \sigma_1^2(\sigma_5\sigma_1)^2,\ \sigma_3\sigma_5 = \sigma_5\sigma_1^2$. The group generated by reflections in the planes $e_1^T x, e_2^T x, e_3^T x = 0$ corresponds to the group generated by $\sigma_2\sigma_3,\ \sigma_1^2\sigma_2\sigma_3,\ \sigma_1^2\sigma_2$ and the group of permutations of coordinates generated by $e_1 \leftrightarrow e_2,\ e_2 \leftrightarrow e_3,\ e_3 \leftrightarrow e_1$ corresponds to the group generated by $\sigma_1\sigma_2\sigma_3,\ \sigma_1\sigma_2\sigma_3\sigma_5\sigma_1,\ \sigma_2\sigma_5$. Let $Z \subset \mathbb{C}^3$ be the equilateral paraboloid $Z^T(I_{1,2}Z - 2e_3) = 0$ and $u := \bar{f}_1^T Z,\ v := f_1^T Z,\ u,\ v \in \mathbb{C}$ be a parametrization of Z by its ruling families: $Z = Z(u,v) = uf_1 + v\bar{f}_1 + uve_3$. We have:

$$f_1 \times \bar{f}_1 = ie_3,\ \bar{f}_1 \times e_3 = i\bar{f}_1,\ e_3 \times f_1 = if_1;\ Y(u) = \frac{u}{2}(Y'(u) + \overline{Y'(-2\bar{u}^{-1})}) = -\frac{u^2}{2}\overline{Y(-2\bar{u}^{-1})};$$

$$X(u,v) = \frac{1}{u-v}Y(u) - \frac{1}{2}Y'(u) = \frac{1}{u-v}Y(v) + \frac{1}{2}Y'(v);\ X \circ \sigma_2 = -X,\ \bar{X} = X \circ (\sigma_1^2\sigma_2\sigma_3\sigma_4);$$

$$Z \circ \sigma_2 = (e_2^T x \mapsto -e_2^T x) \circ Z,\ \bar{Z} = Z \circ (\sigma_2\sigma_4);\ Y(u)^T Y(v) = -2(u-v)^2,$$

$$Y(u) \times Y'(u) = 2iY(u),\ Y(u) \times Y(v) = -2i(u-v)^2 X(u,v);\ |dX|^2 = \frac{4du \odot dv}{(u-v)^2},$$

$$X_{uv} = -\frac{2}{(u-v)^2}X;\ I_3 = -\frac{Y(u)Y(v)^T + Y(v)Y(u)^T}{2(u-v)^2} + X(u,v)X(u,v)^T. \tag{3.1}$$

(we use the notation $\phi \odot \varphi := \frac{\phi\otimes\varphi+\varphi\otimes\phi}{2}$ for ϕ, φ $(0,1)$ tensors). Note also:

$$(Ax) \times (Ay) = (\det A)(A^T)^{-1}(x \times y),\ A \in \mathbf{GL}_3(\mathbb{C}),\ x, y \in \mathbb{C}^3 \tag{3.2}$$

of which we shall make consistent use (on the standard basis it states that entries of $(\det A)(A^T)^{-1}$ are co-factors of A; the formula remains valid for $A \in \mathbf{M}_3(\mathbb{C})$, if one replaces $(\det A)(A^T)^{-1}$ with the adjugate of A or for the cross product of $n-1$ vectors in $\mathbb{C}^n$).

Under the rotation $R \in \mathbf{O}_3(\mathbb{C}),\ \det R = 1$, the rulings of the unit sphere X transform as: $(u,v) \mapsto (\frac{e_3^T RY(u)}{f_1^T RY(u)}, \frac{e_3^T RY(v)}{f_1^T RY(v)}) = (\frac{au+b}{cu+d}, \frac{av+b}{cv+d})$, where $R = \frac{d^2}{2(ad-bc)}Y(\frac{b}{d})f_1^T - \frac{c^2}{ad-bc}Y(\frac{a}{c})\bar{f}_1^T + X(\frac{a}{c}, \frac{b}{d})e_3^T = \frac{a^2}{2(ad-bc)}f_1 Y(-\frac{b}{a})^T - \frac{c^2}{ad-bc}\bar{f}_1 Y(-\frac{d}{c})^T - e_3 X(\frac{b}{a}, \frac{d}{c})^T$; if $\det R = -1$ compose with σ_2. For Z R must satisfy $Re_3 = e_3$; if $\det R = 1$, then $R = e^{ic}f_1\bar{f}_1^T + e^{-ic}\bar{f}_1 f_1^T + e_3 e_3^T$ and $(u,v) \mapsto (e^{ic}u, e^{-ic}v)$; otherwise compose with σ_2. We have the unit pseudo-sphere $|iX|^2 = -1,\ iX \subset \mathbb{R}^2 \times i\mathbb{R} \Leftrightarrow X \subset (i\mathbb{R})^2 \times \mathbb{R} \Leftrightarrow u\bar{v} = 2;\ |d(iX)|^2 = \frac{8dv\odot d\bar{v}}{(2-|v|^2)^2}$ and the real unit sphere $X \subset \mathbb{R}^3,\ |X|^2 = 1 \Leftrightarrow u\bar{v} = -2;\ |dX|^2 = \frac{8dv\odot d\bar{v}}{(2+|v|^2)^2}$. The usual linear element of the (pseudo-)sphere $(\frac{4dv\odot d\bar{v}}{(1-|v|^2)^2})\ \frac{4dv\odot d\bar{v}}{(1+|v|^2)^2}$ is obtained by replacing v with $\sqrt{2}v$, or equivalently for $f_1 := e_1 - ie_2,\ Y(v) := -v^2 f_1 + \bar{f}_1 + 2ve_3$, but we prefer to normalize $f_1^T \bar{f}_1 = 1$ since the computations involving SJ matrices would otherwise acquire as cumbersome constants powers of 2. The real equilateral paraboloid of revolution $Z \subset \mathbb{R}^3$ is obtained for $u = \bar{v}$; the real equilateral hyperbolic paraboloid $\mathrm{diag}[1\ \ i\ \ 1]Z \subset \mathbb{R}^3$ is obtained for $u = \bar{u},\ v = \bar{v}$.

With $\epsilon(z,a) := \frac{\sqrt{1-za^{-1}}(\sqrt{a^{-1}})^{-1}}{\sqrt{a-z}} = \pm 1$ the confocal families of diagonal Q(W)C and of the Darboux quadrics, the ruling families parametrization invariant under the Ivory affinity, isotropic

rulings, umbilics and ruling parameters (if any) which give $x_z \cap^T C(\infty)$ can be put as: QC:(3.3)-(3.5), QWC:(3.6)-(3.7), IQWC:(3.8)-(3.9):

$$x_z^T A R_z^{-1} x_z - 1 = 0,\ A^{-1} := \Sigma_{j=1}^3 a_j e_j e_j^T;\ x_z(u,v) := \sqrt{R_z}(\sqrt{A})^{-1} X(u,v);\ |\sqrt{R_z}(\sqrt{A})^{-1} Y(v)|^2 =$$
$$0 \Leftrightarrow \frac{v}{\sqrt{2}} = \pm\sqrt{\frac{a_1-a_3}{a_1-a_2}} \pm i\sqrt{\frac{a_2-a_3}{a_2-a_1}}\ (= 0, \infty \text{ if } a_1 = a_2);\ \sqrt{R_z}(\sqrt{A})^{-1}(\pm\sqrt{\frac{a_2-a_1}{a_2-a_3}} e_2 \pm$$
$$\sqrt{\frac{a_3-a_1}{a_3-a_2}} e_3),\ \sqrt{R_z}(\sqrt{A})^{-1}(\pm\sqrt{\frac{a_1-a_2}{a_1-a_3}} e_1 \pm \sqrt{\frac{a_3-a_2}{a_3-a_1}} e_3),\ \sqrt{R_z}(\sqrt{A})^{-1}(\pm\sqrt{\frac{a_1-a_3}{a_1-a_2}} e_1 \pm$$
$$\sqrt{\frac{a_2-a_3}{a_2-a_1}} e_2)\ (\pm\sqrt{R_z}(\sqrt{A})^{-1} e_3\ if\ a_1 = a_2);\ \text{none}\ (x_z(0,0),\ x_z(\infty,\infty) \text{ if } a_1 = a_2). \tag{3.3}$$

$$x_z^T A R_z^{-1} x_z - 1 = 0,\ A^{-1} := (a_1 I_2 + J_2) \oplus a_2 I_1;\ x_z(u,v) := \sqrt{R_z}(\sqrt{A})^{-1} X(u,v) =$$
$$(\frac{\sqrt{a_1 - z}}{\epsilon(z,a_1)}(I_2 + \frac{J_2}{2(a_1-z)}) \oplus \frac{\sqrt{a_2-z}}{\epsilon(z,a_2)} I_1) X(u,v);\ |\sqrt{R_z}(\sqrt{A})^{-1} Y(v)|^2 = 0 \Leftrightarrow \frac{v}{\sqrt{2}} = \infty,$$
$$\pm\frac{1}{\sqrt{a_1-a_2}};\ \pm\sqrt{R_z}(\sqrt{A})^{-1}(\frac{f_1}{\sqrt{a_1-a_2}} \pm e_3),\ \pm\sqrt{R_z}(\sqrt{A})^{-1}\frac{f_1 + 2(a_1-a_2)\bar{f}_1}{2\sqrt{a_1-a_2}}$$
$$(\text{none if } a_1 = a_2);\ x_z(\infty,\infty). \tag{3.4}$$

$$x_z^T A R_z^{-1} x_z - 1 = 0,\ A^{-1} := a_1 I_3 + J_3;\ x_z(u,v) := \sqrt{R_z}(\sqrt{A})^{-1} X(u,v) = \frac{\sqrt{a_1 - z}}{\epsilon(z,a_1)}(I_3 +$$
$$\frac{J_3}{2(a_1-z)} - \frac{J_3^2}{8(a_1-z)^2}) X(u,v);\ |\sqrt{R_z}(\sqrt{A})^{-1} Y(v)|^2 = 0 \Leftrightarrow v = \infty,\ 0;\ \pm\sqrt{R_z}(\sqrt{A})^{-1} e_3;$$
$$x_z(\infty,\infty). \tag{3.5}$$

$$x_z^T (A R_z^{-1} x_z - 2e_3) + z = 0,\ A^{\circ -1} := a_1 I_1 \oplus a_2 I_1;\ x_z(u,v) := \sqrt{R_z} L Z(u,v) + \frac{z}{2} e_3,$$
$$L := (\sqrt{A \oplus I_1})^{-1};\ |x_{zu}(u, \pm\sqrt{\frac{a_2-a_1}{2}})|^2 = |x_{zv}(\pm\sqrt{\frac{a_2-a_1}{2}}, v)|^2 = 0;\ \sqrt{R_z} L(\pm\sqrt{a_2-a_1} e_1 +$$
$$\frac{a_2 - a_1 + z}{2} e_3),\ \sqrt{R_z} L(\pm\sqrt{a_1-a_2} e_2 + \frac{a_1 - a_2 + z}{2} e_3);\ \text{none}\ (x_z(0,\infty),\ x_z(\infty,0) \text{ if}$$
$$a_1 = a_2). \tag{3.6}$$

$$x_z^T (A R_z^{-1} x_z - 2e_3) + z = 0,\ A^{\circ -1} := a_1 I_2 + J_2;\ x_z(u,v) := \sqrt{R_z} L Z(u,v) + \frac{z}{2} e_3,$$
$$L := (\sqrt{A \oplus I_1})^{-1},\ \sqrt{R_z} L = \frac{\sqrt{a_1 - z}}{\epsilon(z,a_1)}(I_2 + \frac{J_2}{2(a_1-z)}) \oplus I_1;\ |x_{zu}(u,0)|^2 =$$
$$|x_{zv}(\pm i, v)|^2 = 0;\ \pm i\sqrt{a_1 - z} f_1 + \frac{z}{2} e_3;\ x_z(\infty,0). \tag{3.7}$$

$$(x_z - z\bar{f}_1)^T (A R_z^{-1}(x_z - z\bar{f}_1) - 2\bar{f}_1) = 0,\ A^{\circ -1} := \bar{J}_2 \oplus a_1 I_1;\ x_z(u,v) :=$$
$$L\sqrt{I_3 - z L^T A^2 L} Z(u_z, v_z) + z\bar{f}_1,\ L := f_1 e_3^T + \bar{f}_1 e_1^T - \sqrt{a_1} e_3 e_2^T,$$
$$u(z) = v(z) := -\frac{z}{2\sqrt{2}};\ |x_{zu}(u,\infty)|^2 = |x_{zv}(\infty, v)|^2 = |x_{zu}(u, \frac{a_1}{2\sqrt{2}})|^2 =$$
$$|x_{zv}(\frac{a_1}{2\sqrt{2}}, v)|^2 = 0;\ \frac{(a_1 - z)^2}{8} f_1 + \frac{a_1 + z}{2} \bar{f}_1;\ x_z(\infty,\infty). \tag{3.8}$$

$$(x_z - z\bar{f}_1)^T(AR_z^{-1}(x_z - z\bar{f}_1) - 2\bar{f}_1) = 0,\ A^{\circ -1} := \bar{J}_3;\ x_z(u,v) :=$$
$$L\sqrt{I_3 - zL^TA^2L}Z(u_z, v_z) + z\bar{f}_1,\ L := J_3 + \bar{J}_3^2,\ u(z) := -\frac{z}{2},$$

$$v(z) := -\frac{z^2}{8};\ |x_{zu}(u,\infty)|^2 = |x_{zv}(\infty,v)|^2 = |x_{zu}(u,0)|^2 = 0;\ \text{none};\ x_z(\infty,\infty). \tag{3.9}$$

Let $f := e_3$ ($:= \bar{f}_1$) respectively for (I)QWC, $u(z) = v(z) = 0$ for QWC, $u_z := u + u(z),\ v_z := v + v(z)$. Note:

$$L^{-1} = L^TA + e_3f^T,\ L^Tf = e_3,\ L^{-1}\sqrt{R_z}L = \sqrt{R_z'} + e_3(u(z)f_1 + v(z)\bar{f}_1)^T,$$
$$A' := L^TA^2L = A \text{ for QWC}, = a_1^{-1}e_2e_2^T \text{ for IQWC (3.8)}, = \bar{J}_3^2 \text{ for IQWC (3.9)};$$
$$(L^TL)^{-1}e_3 = e_3 \text{ for QWC}, = e_1 \text{ for IQWC (3.8)}, = f_1 \text{ for IQWC (3.9)};$$
$$L^{-1}x_z(u,v) = \sqrt{R_z'}Z(u_z,v_z) + z(1 - \frac{|f|^2}{2})(L^TL)^{-1}e_3,\ L^T\hat{N}_0 = I_{1,2}Z - e_3. \tag{3.10}$$

Note that the last relation of (3.1) has a metric projective symmetric equivalent for any quadric:

$$x_{zu}x_{zv}^T + x_{zv}x_{zu}^T - \frac{x_{zuv}x_z^T + x_zx_{zuv}^T}{2} = -\frac{2}{\mathcal{B}}(zI_3 - A^{\circ -1}) \text{ for QC},$$
$$x_{zu}x_{zv}^T + x_{zv}x_{zu}^T - (x_{zuv}x_z^T + x_zx_{zuv}^T) = -\frac{2}{\mathcal{B}}(zI_3 - A^{\circ -1}) \text{ for (I)QWC},$$
$$\mathcal{B} := (u-v)^2 \text{ for QC}, := 2 \text{ for (I)QWC}. \tag{3.11}$$

For QC it is just the last relation of (3.1) after using the next to last relation of (3.1) and multiplying on both left and right with $(\sqrt{R_z})^{-1}\sqrt{A}$. For (I)QWC after multiplying on the left with L^{-1} and on the right with $(L^T)^{-1}$ it becomes $(\sqrt{R_z'}f_1 + v_ze_3)(\sqrt{R_z'}\bar{f}_1 + u_ze_3)^T + (\sqrt{R_z'}\bar{f}_1 + u_ze_3)(\sqrt{R_z'}f_1 + v_ze_3)^T - e_3(u_z\sqrt{R_z'}f_1 + v_z\sqrt{R_z'}\bar{f}_1 + u_zv_ze_3 + z(1 - \frac{|f|^2}{2})(L^TL)^{-1}e_3)^T - (u_z\sqrt{R_z'}f_1 + v_z\sqrt{R_z'}\bar{f}_1 + u_zv_ze_3 + z(1 - \frac{|f|^2}{2})(L^TL)^{-1}e_3)e_3^T = -z(L^TL)^{-1} + L^{-1}A^{\circ -1}(L^T)^{-1}$; this boils down to $-L^TA^2L = (1 - \frac{|f|^2}{2})((L^TL)^{-1}e_3e_3^T + e_3e_3^T(L^TL)^{-1}) - (L^TL)^{-1} (= -I_{1,2}(L^TL)^{-1}I_{1,2}),\ I_{1,2} = L^{-1}A^{\circ -1}(L^T)^{-1}$ which are straightforward. Note that (3.11) admits easy generalizations to quadrics in $\mathbb{C}^n$; for the (I)QWC case the relevant parametrization and necessary identities have already been introduced; for the QC case the only important issue is finding the equivalent of the last relation of (3.1). Again just as for (I)QWC a parametrization of the unit sphere $X_{2n+1} \subset \mathbb{C}^{2n+1}$ comes first, following that $X_{2n} := X_{2n+1} \cap \{x \in \mathbb{C}^{2n+1} | e_{2n+1}^Tx = 0\}$. With $X(u,v)* : \mathbb{C}^{2n-1} \to \mathbb{C}^{2n+1},\ X * V := [\frac{2-uv}{\sqrt{2}(u-v)}\ i\frac{2+uv}{\sqrt{2}(u-v)}\ \frac{u+v}{u-v}V^T]^T$ we have by induction $X_{2n+1}(u^1,v^1,...,u^n,v^n) = X(u^n,v^n) * X_{2n-1}(u^1,v^1,...,u^{n-1},v^{n-1})$.

Note that all quadrics $\{x_z\}_{z\in\mathbb{C}}$ confocal to the given one x_0 and all quadrics homothetic to x_0 are of the x_0 type. Since modulo rigid motions there is a single IQWC (3.9), all quadrics x_z confocal to the IQWC (3.9) x_0 or homothetic to x_0 are rigidly applicable to x_0. This can be confirmed analytically as follows: the rigid motion $(I_3 - \frac{z}{2}(f_1e_3^T - e_3f_1^T) - \frac{z^2}{8}e_3e_3^T, \frac{z^3}{16}f_1 - \frac{3z}{2}\bar{f}_1 - \frac{3z^2}{8}e_3) \in \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ brings $x_z(u,v)$ to the canonical form $x_0(u-z,v)$; also the rigid motion $(R(c),0) \in \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ brings $e^{2ic}x_0(u,v)$ to the canonical form $x_0(e^{ic}u, e^{2ic}v)$.

$x_z \cap^T C(\infty)$ can be seen after a homography (homographies preserve tangency and its order); we consider only $z = 0$. Under the homography $\mathbb{C}^3 \ni x := [x^1\ x^2\ x^3\ 1] \mapsto [\frac{f_1^Tx}{f_1^Tx}\ \frac{e_3^Tx}{f_1^Tx}\ \frac{1}{f_1^Tx}\ 1]$, $C(\infty)$ becomes $c(t) := -2[t^2\ t\ 0]$ and $[f_1^T,\ 0]$ becomes 0. The curves $(u^{-1}, v) := (O(t^2), -\sqrt{a_1}t + O(t^2))$, $(u^{-1}, v^{-1}) := (-i\sqrt{\frac{2}{a_1}}t + O(t^2), i\sqrt{\frac{2}{a_1}}t + O(t^2))$ respectively on the transformed (3.7), (3.8) meet $c(t)$ tangentially at $t = 0$; the curves $(u^{-1}, v^{-1}) := (t - \frac{3}{a_1}t^2 + O(t^3), t - \frac{3}{a_1}t^2 + O(t^3))$, $(u^{-1}, v^{-1}) := (-2t + O(t^3), -2t^2 + O(t^3))$ respectively on the transformed (3.5), (3.9) osculate $c(t)$ at $t = 0$ up to order 2; the curves $(u^{-1}, v^{-1}) := (t - \frac{t^3}{a_1} + O(t^4), t - \frac{t^3}{a_1} + O(t^4))$ on the transformed (3.4) for $a_2 = a_1$ osculate $c(t)$ at $t = 0$ up to order 3.

According to Eisenhart ([19],§ 151), Goursat integrated the equations for the deformations of the paraboloids (3.7) and (3.9); also Darboux reduced in [15] the deformations of the Darboux quadrics to the deformations of the sphere.

For QC (3.3) if $a_j,\ j=1,2,3$ are distinct, then x_z meets $C(\infty),\ \mathcal{S}(x_{a_j}),\ j=1,2,3$ only transversally; if $a_1=a_2\neq a_3$, then x_z meets $C(\infty),\ \mathcal{S}(x_{a_3})$ only tangentially at $[f_1^T,0],\ [\bar{f}_1^T,0]$ and $\mathcal{S}(x_{a_1})$ only transversally at $\pm\sqrt{a_3-z}e_3$. For QC (3.4) if $a_1\neq a_2$, then x_z meets $C(\infty),\ \mathcal{S}(x_{a_2})$ tangentially at $[f_1^T,0]$ and transversally at two more points each; further x_z meets $\mathcal{S}(x_{a_1})$ only transversally; if $a_1=a_2$, then x_z meets $C(\infty),\ \mathcal{S}(x_{a_1})$ only tangentially at $[f_1^T,0]$. For QC (3.5) x_z meets $C(\infty),\ \mathcal{S}(x_{a_1})$ tangentially at $[f_1^T,0]$ and transversally at one respectively two more points. For QWC (3.6) if $a_1\neq a_2$, then x_z meets $C(\infty)$ only transversally; further it meets $\mathcal{S}(x_{a_j}),\ j=1,2$ tangentially at $[e_3^T,0]$ and transversally at two more points each; if $a_1=a_2$, then x_z meets $C(\infty)$ only tangentially at $[f_1^T,0],\ [\bar{f}_1^T,0]$ and $\mathcal{S}(x_{a_1})$ only transversally at $\frac{z}{2}e_3,\ [e_3^T,0]$. For QWC (3.7) x_z meets $C(\infty),\ \mathcal{S}(x_{a_1})$ tangentially at $[f_1^T,0]$, respectively $[e_3^T,0]$ and transversally at two more points each. For QC (3.8) x_z meets $C(\infty),\ \mathcal{S}(x_{a_1})$ tangentially at $[f_1^T,0]$ and transversally at two respectively one more points. For QC (3.9) x_z meets $C(\infty)$ tangentially at $[f_1^T,0]$ and transversally at one more point. The points of intersection with finite singular quadrics are situated in $\mathbb{CP}^2$ iff they are tangency points.

For $n=3$ Bianchi has a beautiful geometric proof of Bianchi II (see also Darboux ([13],§ 603)): all coordinates on a surface whose definition involves only projective invariants (tangency of different orders; for example tangent planes of a surface, osculating planes of a curve) are preserved by homographies. In particular asymptotes and conjugate systems are preserved; since on developables the asymptotes coincide, homographies take developables to developables. There is a developable circumscribed to two given arbitrary curves and the isotropic developable with generating curve $c(u)$ is the developable $c(u)+vY(f(u))$ circumscribed (tangent) to $c,\ C(\infty)$, so $c'(u)^TY(f(u))=0$; thus it is an algebraic surface of degree four if c is a conic. Now nonsingular quadrics x_z intersect $\mathcal{S}(x_{a_1})$ at four points (Bézout) and the isotropic rulings of the developable through those points are all the isotropic rulings on x_z; by inspection all quadrics are uniquely determined by their isotropic rulings (most quadrics have three distinct isotropic rulings of one ruling family; as a quadric is uniquely determined by three rulings of a ruling family the statement follows in these cases). As a quadric varies in its confocal family its umbilics describe the conics which are the singular sets of singular quadrics of the family and its isotropic rulings isotropic developables generated by these conics. Thus H takes the isotropic developable of the conic $\mathcal{S}(x_{a_1})$ to the isotropic developable of the conic $\mathcal{S}(\tilde{x}_{\tilde{a}_1}):=H(C(\infty))$ and $\tilde{x}_{\tilde{z}}$ is uniquely recovered. For example for QC (3.3) if $\frac{v}{\sqrt{2}}=\epsilon_1(\sqrt{\frac{a_1-a_3}{a_1-a_2}}+i\epsilon_2\sqrt{\frac{a_2-a_3}{a_2-a_1}}),\ c(z):=\sqrt{R_z}(\sqrt{A})^{-1}\epsilon_3(\sqrt{\frac{a_1-a_3}{a_1-a_2}}e_1+\ \epsilon_4\sqrt{\frac{a_2-a_3}{a_2-a_1}}e_2),\ \epsilon_j=\pm1,\ j=1,...,4$, then $c'(z)^T\sqrt{R_z}(\sqrt{A})^{-1}Y(v)=i\epsilon_3\sqrt{2}\sqrt{\frac{a_1-a_3}{a_1-a_2}}\sqrt{\frac{a_2-a_3}{a_2-a_1}}(\sqrt{\frac{a_1-a_3}{a_1-a_2}}+i\epsilon_2\sqrt{\frac{a_2-a_3}{a_2-a_1}})(\epsilon_2-\epsilon_4)=0$ for $\epsilon_2=\epsilon_4$.

Let $H\in\mathbf{PGL}_4(\mathbb{C})$ be a homography of $\mathbb{C}^3\ni x\simeq[x^T,1]\in\mathbb{CP}^3$, with $\tilde{x}:=\frac{I_{1,3}Hx}{e_4^THx}$. We shall always have $\tilde{x}_{\tilde{z}}(\tilde{u},\tilde{v}):=H(x_z(u,v)),\ H:=\begin{bmatrix}H' & h_1\\ h_2^T & 0\end{bmatrix},\ \tilde{H}:=H^{-1}=\begin{bmatrix}\tilde{H}' & \tilde{h}_1\\ \tilde{h}_2^T & 0\end{bmatrix}$; the rulings $\tilde{u},\ \tilde{v}$ may change with z (this corresponds to symmetries of $\tilde{x}_{\tilde{z}}$ wrt the principal planes).

If $H'=\Sigma_{j=2}^3\frac{1}{\sqrt{a_1-a_j}}e_je_j^T,\ \tilde{H}'=\Sigma_{j=2}^3\sqrt{a_1-a_j}e_je_j^T,\ h_1=h_2=\tilde{h}_1=\tilde{h}_2=e_1$, then $x_z(u,v)$, $\tilde{x}_{\tilde{z}}(\tilde{u},\tilde{v})$ are QC of type (3.3), $\tilde{a}_1:=a_1^{-1},\ \tilde{a}_j:=\tilde{a}_1+(a_j-a_1)^{-1},\ j=2,3,\ \tilde{z}:=\tilde{a}_1+(z-a_1)^{-1}$; H takes $x_z\cap\{e_1^Tx=0\}=\{\sqrt{R_z}(\sqrt{A})^{-1}X(2v^{-1},v)\}$ to $\tilde{x}_{\tilde{z}}\cap\mathbb{CP}^2=\{[(\sqrt{\tilde{R}_{\tilde{z}}}(\sqrt{\tilde{A}})^{-1}Y(\tilde{v}))^T,0]\}$, so $(\tilde{u},\tilde{v})=(i\epsilon_3\frac{2-i\epsilon_2\sqrt{2}u}{\sqrt{2}+i\epsilon_2u},-i\epsilon_3\frac{2+i\epsilon_2\sqrt{2}v}{\sqrt{2}-i\epsilon_2v}),\ \epsilon_j:=\frac{i\epsilon(\tilde{z},\tilde{a}_1)\epsilon(z,a_j)\sqrt{a_j-z}\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(\tilde{z},\tilde{a}_j)\sqrt{a_1-a_j}\sqrt{\tilde{a}_j-\tilde{z}}},\ j=2,3$. If $a_2\neq a_3$, then $H(x_z\pitchfork C(\infty))=\tilde{x}_{\tilde{z}}\pitchfork\mathcal{S}(\tilde{x}_{\tilde{a}_1}),\ H(x_z\pitchfork\mathcal{S}(x_{a_1}))=\tilde{x}_{\tilde{z}}\pitchfork C(\infty),\ H(x_z\pitchfork\mathcal{S}(x_{a_j}))=\tilde{x}_{\tilde{z}}\pitchfork\mathcal{S}(\tilde{x}_{\tilde{a}_j}),\ j=2,3$; if $a_2=a_3$, then $H(x_z\cap^T C(\infty))=\tilde{x}_{\tilde{z}}\cap^T C(\infty),\ H(x_z\pitchfork\mathcal{S}(x_{a_2}))=\tilde{x}_{\tilde{z}}\pitchfork\mathcal{S}(\tilde{x}_{\tilde{a}_2})$. Under the

homothety $\frac{1}{\sqrt{(a_1-a_2)(a_1-a_3)}}$ and $x^2 \leftrightarrow x^3$ in the range of H, $\tilde{a}_j$ become $a_j - w$, $j = 1, 3$, $w := a_1 - \frac{(a_1-a_2)(a_1-a_3)}{a_1}$, so the new H takes the family of confocal QC (3.3) into itself.

If $H' = i\sqrt{\tilde{a}_2 - \tilde{a}_1}((a_2 - a_1)f_1\bar{f}_1^T + \frac{1}{2}f_1f_1^T - \frac{1}{a_2-a_1}\bar{f}_1f_1^T)$, $\tilde{H}' = \frac{-i}{\sqrt{\tilde{a}_2-\tilde{a}_1}}(\frac{1}{a_2-a_1}f_1\bar{f}_1^T + \frac{1}{2}f_1f_1^T - (a_2 - a_1)\bar{f}_1f_1^T)$, $h_1 = h_2 = \tilde{h}_1 = \tilde{h}_2 = e_3$, then $x_z(u, v)$, $\tilde{x}_{\tilde{z}}(\tilde{u}, \tilde{v})$ are QC of the type (3.4), $\tilde{a}_2 := a_2^{-1}$, $\tilde{a}_1 := \tilde{a}_2 + (a_1 - a_2)^{-1}$, $\tilde{z} := \tilde{a}_2 + (z - a_2)^{-1}$; H takes the cone $x_z \cap \{f_1^T x = 0\} = \{\sqrt{R_z}(\sqrt{A})^{-1}X(\infty, v)\} \cup \{\sqrt{R_z}(\sqrt{A})^{-1}X(u, \infty)\}$ (with vertex $x_z \cap^T C(\infty)$) to the cone $\tilde{x}_{\tilde{z}} \cap \{f_1^T x = 0\} = \{\sqrt{\tilde{R}_{\tilde{z}}}(\sqrt{\tilde{A}})^{-1}X(\infty, \tilde{v})\} \cup \{\sqrt{\tilde{R}_{\tilde{z}}}(\sqrt{\tilde{A}})^{-1}X(\tilde{u}, \infty)\}$ with same vertex, so $(\tilde{u}, \tilde{v}) = (-\epsilon_1 u, \epsilon_1 v)$, $\epsilon_1 := \frac{i\epsilon(z,a_1)\epsilon(z,a_2)\epsilon(\tilde{z},\tilde{a}_1)\sqrt{a_1-z}}{\sqrt{\tilde{a}_2-\tilde{a}_1}\sqrt{a_2-z}\sqrt{\tilde{a}_1-\tilde{z}}}$ if $\frac{\sqrt{a_2-z}\sqrt{\tilde{a}_2-\tilde{z}}}{\epsilon(z,a_2)\epsilon(\tilde{z},\tilde{a}_2)} = 1$ and composed with σ_2 otherwise; also $H(x_z \cap^T C(\infty)) = \tilde{x}_{\tilde{z}} \cap^T \mathcal{S}(\tilde{x}_{\tilde{a}_2})$, $H(x_z \pitchfork C(\infty)) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_2})$, $H(x_z \cap^T \mathcal{S}(x_{a_2})) = \tilde{x}_{\tilde{z}} \cap^T C(\infty)$, $H(x_z \pitchfork \mathcal{S}(x_{a_2})) = \tilde{x}_{\tilde{z}} \pitchfork C(\infty)$, $H(x_z \pitchfork \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_1})$.

If $H' = ie_3(\frac{\tilde{a}_1}{2}f_1 - \bar{f}_1)^T + \sqrt{\tilde{a}_1 - \tilde{a}_2}e_2e_3^T$, $\tilde{H}' = if_1e_3^T + \frac{1}{\sqrt{\tilde{a}_1-\tilde{a}_2}}e_3e_2^T$, $h_1 = \tilde{h}_2 = e_1$, $h_2 = if_1$, $\tilde{h}_1 = -i(\bar{f}_1 + \frac{a_1}{2}f_1)$, then $x_z(u, v)$ are QC of type (3.4), $\tilde{x}_{\tilde{z}}(\tilde{u}, \tilde{v})$ are QWC of type (3.6): $\tilde{a}_1 := a_1^{-1}$, $\tilde{a}_2 := \tilde{a}_1 + (a_2 - a_1)^{-1}$, $\tilde{z} := \tilde{a}_1 + (z - a_1)^{-1}$; H takes the cone $x_z \cap \{f_1^T x = 0\} = \{\sqrt{R_z}(\sqrt{A})^{-1}X(\infty, v)\} \cup \{\sqrt{R_z}(\sqrt{A})^{-1}X(u, \infty)\}$ (with vertex $x_z \cap^T C(\infty) \notin \mathcal{S}(x_{a_1})$) to the cone $\tilde{x}_{\tilde{z}} \cap \mathbb{CP}^2 = \{[\tilde{x}_{\tilde{z}}(\infty, \tilde{v})^T, 0]\} \cup \{[\tilde{x}_{\tilde{z}}(\tilde{u}, \infty)^T, 0]\} = \{[(\sqrt{\tilde{R}_{\tilde{z}}}L\bar{f}_1 + 2\tilde{v}e_3)^T, 0]\} \cup \{[(\sqrt{\tilde{R}_{\tilde{z}}}L\bar{f}_1 + 2\tilde{u}e_3)^T, 0]\}$ with vertex $[e_3^T, 0] \notin C(\infty)$, so $(\tilde{u}, \tilde{v}) = (\frac{-i\epsilon_1}{\sqrt{2}}u, \frac{i\epsilon_1}{\sqrt{2}}v)$, $\epsilon_1 := \frac{\sqrt{a_1-z}\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(z,a_1)\epsilon(\tilde{z},\tilde{a}_1)}$ if $\frac{-i\epsilon(z,a_2)\epsilon(\tilde{z},\tilde{a}_1)\sqrt{\tilde{a}_2-\tilde{z}}}{\sqrt{\tilde{a}_1-\tilde{a}_2}\sqrt{\tilde{a}_1-\tilde{z}}\sqrt{a_2-z}} = \epsilon(\tilde{z}, \tilde{a}_2)$ and composed with σ_2 otherwise; also $H(x_z \cap^T C(\infty)) = \tilde{x}_{\tilde{z}} \cap^T \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \pitchfork C(\infty)) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \pitchfork \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \pitchfork C(\infty)$, $H(x_z \cap^T \mathcal{S}(x_{a_2})) = \tilde{x}_{\tilde{z}} \cap^T \mathcal{S}(\tilde{x}_{\tilde{a}_2})$, $H(x_z \pitchfork \mathcal{S}(x_{a_2})) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_2})$.

If $H' = i(f_1\bar{f}_1^T + \bar{f}_1(-\tilde{a}_1f_1 + e_3)^T)$, $\tilde{H}' = -i(f_1\bar{f}_1^T + e_3f_1^T)$, $h_1 = \tilde{h}_2 = e_3$, $\tilde{h}_1 = i(\bar{f}_1 + \tilde{a}_1e_3)$, $h_2 = -if_1$, then $x_z(u, v)$ are QC of type (3.5), $\tilde{x}_{\tilde{z}}(\tilde{u}, \tilde{v})$ are IQWC of type (3.8): $\tilde{a}_1 := a_1^{-1}$, $\tilde{z} := \tilde{a}_1 + (z - a_1)^{-1}$; H takes the cone $x_z \cap \{f_1^T x = 0\} = \{\sqrt{R_z}(\sqrt{A})^{-1}X(\infty, v)\} \cup \{\sqrt{R_z}(\sqrt{A})^{-1}X(u, \infty)\}$ with vertex $x_z \cap^T C(\infty) \in \mathcal{S}(x_{a_1})$ to the cone $\tilde{x}_{\tilde{z}} \cap \mathbb{CP}^2 = \{[\tilde{x}_{\tilde{z}}(\infty, \tilde{v})^T, 0]\} \cup \{[\tilde{x}_{\tilde{z}}(\tilde{u}, \infty)^T, 0]\} = \{[(\bar{f}_1 + (\sqrt{2}\tilde{v} - \frac{\tilde{z}}{2})f_1 + i\frac{\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(\tilde{z},\tilde{a}_1)}e_3)^T, 0]\} \cup \{[(\bar{f}_1 + (\sqrt{2}\tilde{u} - \frac{\tilde{z}}{2})f_1 - i\frac{\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(\tilde{z},\tilde{a}_1)}e_3)^T, 0]\}$ with vertex $\tilde{x}_{\tilde{z}} \cap^T C(\infty) \in C(\infty)$, so $(\tilde{u}, \tilde{v}) = (\frac{1}{\sqrt{2}}(\frac{\tilde{a}_1}{2} - u), \frac{1}{\sqrt{2}}(\frac{\tilde{a}_1}{2} - v))$ if $\frac{\sqrt{a_1-z}\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(z,a_1)\epsilon(\tilde{z},\tilde{a}_1)} = 1$ or composed with σ_2 otherwise; also $H(x_z \cap^T C(\infty)) = \tilde{x}_{\tilde{z}} \cap^T \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \pitchfork C(\infty)) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \cap^T \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \cap^T C(\infty)$, $H(x_z \pitchfork \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \pitchfork C(\infty)$.

If $H' = i(f_1e_3^T + e_3(\frac{\tilde{a}_1}{2}f_1 - \bar{f}_1)^T)$, $\tilde{h}_1 = -i(\frac{\tilde{a}_1}{2}f_1 + \bar{f}_1)$, $h_2 = if_1$ and for $\tilde{H}'$, h_1, $\tilde{h}_2$ replace $\tilde{a}_1$ with a_1, then $x_z(u, v)$, $\tilde{x}_{\tilde{z}}(\tilde{u}, \tilde{v})$ are QWC of type (3.7), $\tilde{a}_1 := a_1^{-1}$, $\tilde{z} := \tilde{a}_1 + (z - a_1)^{-1}$; H takes the cone $x_z \cap \{f_1^T x = 0\} = \{\frac{\sqrt{a_1-z}}{\epsilon(z,a_1)}uf_1 + \frac{z}{2}e_3\} \cup \{[(\frac{\sqrt{a_1-z}}{\epsilon(z,a_1)}f_1 + ve_3)^T, 0]\}$ with vertex $x_z \cap^T C(\infty) \notin \mathcal{S}(x_{a_1})$ to the cone $\tilde{x}_{\tilde{z}} \cap \mathbb{CP}^2 = \{[(\frac{\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(\tilde{z},\tilde{a}_1)}(\bar{f}_1 + \frac{f_1}{2(\tilde{a}_1-\tilde{z})}) + \tilde{u}e_3)^T, 0]\} \cup \{[(\frac{\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(\tilde{z},\tilde{a}_1)}f_1 + \tilde{v}e_3)^T, 0]\}$ with vertex $[e_3^T, 0] \notin C(\infty)$, so $(\tilde{u}, \tilde{v}) = (\epsilon_1 u, \frac{-\epsilon_1}{v})$, $\epsilon_1 := \frac{\sqrt{a_1-z}\sqrt{\tilde{a}_1-\tilde{z}}}{\epsilon(z,a_1)\epsilon(\tilde{z},\tilde{a}_1)}$; also $H(x_z \cap^T C(\infty)) = \tilde{x}_{\tilde{z}} \cap^T \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \pitchfork C(\infty)) = \tilde{x}_{\tilde{z}} \pitchfork \mathcal{S}(\tilde{x}_{\tilde{a}_1})$, $H(x_z \cap^T \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \cap^T C(\infty)$, $H(x_z \pitchfork \mathcal{S}(x_{a_1})) = \tilde{x}_{\tilde{z}} \pitchfork C(\infty)$.

Homographies with $H' = -\bar{J}_2 \oplus e^a I_1$, $\tilde{H}' = -J_2 \oplus e^{-a} I_1$, $e^{-c}h_1 = h_2 = f_1$, $\tilde{h}_1 = e^c\tilde{h}_2 = \bar{f}_1$ or $H = \begin{bmatrix} H' & 0 \\ 0^T & 1 \end{bmatrix}$, $e^c I_3 \neq H' \in \mathbf{GL}_3(\mathbb{C})$ SJ do not have the property of taking a family of confocal quadrics to another one and up to rigid motions and homotheties in both the range and domain of H all homographies $H \in \mathbf{PGL}_4(\mathbb{C})$ can be brought to one of the cases above or to the identity. Quadrics whose confocal family cannot be taken to another one by homographies are the (pseudo-)spheres, QC (3.4) and QWC (3.6) for $a_1 = a_2$, IQWC (3.9).

3.2. **Totally real confocal quadrics in $\mathbb{C}^3$ and the corresponding homographies of Bianchi II.**

With $\mathcal{A} := i \det(\sqrt{A})^{-1}$ for QC, $:= i \det L$ for (I)QWC ($= \sqrt{a_1}$ for IQWC (3.8), $= i$ for IQWC (3.9)) we have the Gauß curvature $K(x_0) = \frac{-1}{\mathcal{A}^2|\hat{N}_0|^4}$ of x_0.

We are interested in surfaces x_0^0 in the quadric x_0 having real linear element (that is $|dx_0^0|^2 = |d\bar{x}_0^0|^2$). By Sylvester's such surfaces must have real dimension 2, so they are obtained by imposing two functionally independent real relations between the four real parameters of x_0. We shall see in § 4.2 that the only such surfaces in spheres are the totally real (pseudo-)spheres. The totally real unit spheres $X(u,v)$ are obtained as follows: $X \subset \mathbb{R}^3$ has signature $(2,0)$ for $v = -2\bar{u}^{-1},\ u \in \mathbb{C}$; $X \subset (i\mathbb{R})^2 \times \mathbb{R}$ has signature $(0,2)$ for $v = 2\bar{u}^{-1},\ u \in \mathbb{C}$; $X \subset \mathbb{R}^2 \times i\mathbb{R}$ has signature $(1,1)$ for $(u,v) = (\sqrt{2}e^{is}, \sqrt{2}e^{it}),\ s,t \in \mathbb{R}$; the totally real unit pseudo-spheres $iX(u,v)$ are obtained through multiplication by i, when the components of both the signature of X and of the ambient totally real space are exchanged.

For the remaining quadrics $K(x_0^0) = \overline{K(x_0^0)}$, or equivalently

$$(I)\ \ \mathcal{A}|\hat{N}_0^0|^2 = \epsilon\overline{\mathcal{A}|\hat{N}_0^0|^2},\ \epsilon := \pm 1$$

provides a non-vacuous real relation between the four real parameters of x_0. Keeping account of $\hat{N}_0^0 = Ax_0^0 + B$ and applying d to (I) we get $(dx_0^0)^T(\mathcal{A}A\hat{N}_0^0) = \epsilon\overline{(dx_0^0)^T(\mathcal{A}A\hat{N}_0^0)}$. Now Sylvester's allows us to find an adapted moving frame $R[\sqrt{\epsilon_1}e_1\ \ \sqrt{\epsilon_2}e_2\ \ e_3],\ \epsilon_j := \pm 1,\ j = 1,2$ chosen so that $\mathrm{diag}[\epsilon_1\ \ \epsilon_2]$ is the signature of $|dx_0^0|^2,\ R \subset \mathbf{O}_3(\mathbb{C}),\ R^{-1}dx_0^0 = \sqrt{\epsilon_1}\psi_1 e_1 + \sqrt{\epsilon_2}\psi_2 e_2$ and $\psi_j,\ j = 1,2$ are real (valued) independent 1-forms. If $R^{-1}\mathcal{A}A\hat{N}_0^0 =: \sqrt{\epsilon_1}v_1e_1 + \sqrt{\epsilon_2}v_2e_2 + v_3e_3,\ v_j \in \mathbb{C}$, then we get $\epsilon_1(v_1 - \epsilon\bar{v}_1)\psi_1 + \epsilon_2(v_2 - \epsilon\bar{v}_2)\psi_2 = 0$, so $v_j = \epsilon\bar{v}_j,\ v_j^2 = \bar{v}_j^2,\ j = 1,2$. This implies that the tangential component $\frac{(\hat{N}_0^0 \times (\mathcal{A}A\hat{N}_0^0)) \times \hat{N}_0^0}{|\hat{N}_0^0|^2}$ of the vector $\mathcal{A}A\hat{N}_0^0$ has real length; using also (I) we obtain:

$$(II)\ \ \mathcal{A}^3|V_2|^2 = \epsilon\overline{\mathcal{A}^3|V_2|^2},\ V_2 := \hat{N}_0^0 \times A\hat{N}_0^0.$$

Similarly to the step (I) $\Rightarrow$ (II), one can apply d to (II) and get $\mathcal{A}^7|V_3|^2 = \epsilon\overline{\mathcal{A}^7|V_3|^2},\ V_3 := \hat{N}_0^0 \times A(A(V_2 \times \hat{N}_0^0) - V_2 \times A\hat{N}_0^0)$; in fact one can get a whole hierarchy of consequences, but (I)&(II) will be sufficient for our purposes, because they are functionally independent for all quadrics which are not spheres. We still need to satisfy

$$(III)\ \ |dx_0^0|^2 - |d\bar{x}_0^0|^2 = 0.$$

Since we already have the ruling families parametrization (u,v) on x_0, it is more convenient to work with the parametrization $(u,v,\bar{u},\bar{v})$ or closely related ones $(U,V,\bar{U},\bar{V})$, $U,\ V$ being simple rational functions of u and v with $\frac{dU \wedge dV}{du \wedge dv} \neq 0$ (all functions below will be rational). (I)&(II) will be equivalent to two real relations $f_1(V,\bar{V}) = f_2(U,\bar{U}) = 0$ or will imply $f(U,V,\bar{V}) = 0$ (and thus its conjugate). If one can solve for $U = F(V,\bar{V})$, then this and its conjugate will be functionally independent and thus will imply (I)&(II). However, in this case (III) further imposes two real conditions, namely the condition that $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is degenerate as a symmetric bilinear form in $dV,\ d\bar{V}$ and the condition that the coefficient of $dV \odot d\bar{V}$ in $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is 0; these conditions must not impose a real condition between V and $\bar{V}$. For most cases this program becomes tedious, so other simplifications (usually considering only the hot's in $V,\ \bar{V}$) may be required.

One can try to find another functional relationship between $U,V,\bar{U},\bar{V}$, which is easier to obtain than the third functional relationship of the hierarchy described above or than the method described above; hopefully the condition that (I)&(II) and this new functional relationship are functionally dependent will provide other useful information. An example of such a functional relationship is as follows: (I) gives one of the a-priori independent variables $U,V,\bar{U},\bar{V}$ (for example $\bar{U}$) as a function of the other three variables; replacing this into (III) further provides a functional relationship between the remaining three variables, namely the condition that $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is degenerate as a symmetric bilinear form in the differentials of the three remaining variables; this condition is valid since the real 2-dimensionality of x_0^0 implies that $|dx_0^0|^2 - |d\bar{x}_0^0|^2$, as a symmetric bilinear form in the differentials of the three remaining variables, admits two linearly independent orthogonal isotropic vectors, namely $\partial_s,\ \partial_t$ for s,t real parametrization of x_0^0. Although finding this functional

relationship is more difficult than finding the ones imposed by (I)&(II), a nice feature appears: this functional relationship always factors as a product of $|\hat{N}_0^0|^2 \neq 0$ and another term being 0, because if $|\hat{N}_0^0|^2 = 0$, then x_0^0 is a complex holomorphic curve with linear element extended to isotropic one on x_0^0 (it is real 2-dimensional, but not real valued), so it has degenerate real 3-dimensional linear element. Unfortunately in all cases except IQWC (3.9) this functional relationship is the consequence $f(U, V, \bar{V}) = 0$ of (I)&(II) (for IQWC (3.9) it is the consequence $f_1(u, \bar{u}) = 0$ of (I)&(II) if we begin with $\bar{v}$ as a function of $u,\ \bar{u},\ v$), so it is useful only as a confirmation of the already completed computations.

We shall prove, after a study case by case, that all surfaces in quadrics and having real linear element are totally real, so most quadrics do not contain such surfaces; a simpler unifying argument using rolling will appear in § 6.6.

For QC we have $\mathcal{A} = i \det(\sqrt{A})^{-1},\ x_0^0 = (\sqrt{A})^{-1}X,\ \hat{N}_0^0 = \sqrt{A}X,\ |\hat{N}_0^0|^2 = X^T AX,\ |V_2|^2 = \det A(X \times AX)^T A^{-1}(X \times AX),\ |dx_0^0|^2 = dX^T A^{-1} dX$.

For (I)QWC we have $\mathcal{A} = i \det L,\ x_0^0 = LZ,\ \hat{N}_0^0 = (L^T)^{-1}(I_{1,2}Z - e_3),\ |\hat{N}_0^0|^2 = (I_{1,2}Z - e_3)^T (L^T L)^{-1}(I_{1,2}Z - e_3),\ |V_2|^2 = (\det L)^{-2}((I_{1,2}Z - e_3) \times (A'Z - (L^T L)^{-1}e_3 + |f|^2 e_3))^T (L^T L)((I_{1,2}Z - e_3) \times (A'Z - (L^T L)^{-1}e_3 + |f|^2 e_3)),\ |dx_0^0|^2 = dZ^T (L^T L) dZ,\ L^T L = A^{\circ -1} + e_3 e_3^T$ for QWC, $= e_3 e_1^T + e_1 e_3^T + a_1 e_2 e_2^T$ for IQWC (3.8), $= J_3^2 + \bar{J}_3$ for IQWC (3.9), $(L^T L)^{-1} = A + e_3 e_3^T$ for QWC, $= e_3 e_1^T + e_1 e_3^T + a_1^{-1} e_2 e_2^T$ for IQWC (3.8), $= \bar{J}_3^2 + J_3$ for IQWC (3.9).

Since the most metrically degenerate a complex metric type of a quadric is, the less totally real types it contains and also keeping account of the homographies of Bianchi II, the discussion will take place as follows: IQWC (3.9), IQWC (3.8) & QC (3.5), QWC (3.7), QWC (3.6) & QC (3.4), QC (3.3).

For $\epsilon, \epsilon_1 := \pm 1$ note $\frac{\sqrt{-\epsilon}}{\sqrt{\epsilon}} = i\epsilon,\ \frac{\sqrt{-\epsilon_1\sqrt{-\epsilon}}}{\sqrt{\epsilon_1\sqrt{-\epsilon}}} = -i\epsilon\epsilon_1$, etc, of which we shall make consistent use below.

For IQWC (3.9) (I)&(II) become $u^2 - 2v = -\epsilon(\bar{u}^2 - 2\bar{v}),\ 3u^2 + 2v = -\epsilon(3\bar{u}^2 + 2\bar{v})$, or $u^2 = -\epsilon\bar{u}^2,\ v = -\epsilon\bar{v}$. Keeping account of $|dx_0^0|^2 = 2du \odot d(uv) + dv^2$, (III) will be satisfied.

Thus we have four quadrics x_0^0 in $x_0 = x_0(u, v) = uvf_1 + u\bar{f}_1 + ve_3$, given by $(u, v) = (\sqrt{\epsilon_1\sqrt{-\epsilon}}s, \sqrt{-\epsilon}t),\ \epsilon_1 := \pm 1,\ s, t \in \mathbb{R}$. Since ϵ_1 in $|dx_0^0|^2$ can be absorbed by $(s, t) \mapsto (\epsilon_1 s, \epsilon_1 t)$, there are two types of real linear element of IQWC(3.9) from a real point of view (namely $-\epsilon(2ds \odot d(st) + dt^2)$); from a complex point of view there is only one, since further $-\epsilon$ can be absorbed by $(s, t) \mapsto (\sqrt[4]{-\epsilon}s, \sqrt{-\epsilon}t)$. All symmetries of the linear element are realized by rigid motions composed when necessary with a multiplication by i: $iR(\frac{\pi}{4})x_0(u, v) = x_0(\sqrt{i}u, iv)$; ϵ_1 appears in the linear element of x_0 because of the rotation $(iR(\frac{\pi}{4}))^2 = -R(\frac{\pi}{2})$ which generates the group of rigid motions preserving the canonical form of $x_0 :\ -R(\frac{\pi}{2})x_0(u, v) = x_0(-iu, -v)$. Note that the IQWC (3.9) x_0 is the only quadric such that ix_0 differs from x_0 by a rigid motion; for all other quadrics ix_0 will be a quadric of the same type, but which is different up to rigid motions from x_0. The signature of $|dx_0^0|^2$ is $(1, 1)$ for $2\epsilon_1 t < s^2$ and $\mathrm{diag}[-\epsilon\ \ -\epsilon]$ for the other inequality; the change of signature of $|dx_0^0|^2$ occurs along the curves where the normal directions are isotropic.

With $\mathcal{E} := \mathrm{diag}[1\ \ -1\ \ -\epsilon],\ R_0 := \frac{f_1 \bar{f}_1^T}{\sqrt{\epsilon_1\sqrt{-\epsilon}}} + \sqrt{\epsilon_1\sqrt{-\epsilon}}\bar{f}_1 f_1^T + e_3 e_3^T \in \mathbf{O}_3(\mathbb{R})$ we have $\mathcal{E}f_1 = \bar{f}_1$, so $\mathcal{E}\bar{f}_1 = f_1$; further $\overline{R_0^T x_0^0} = \mathcal{E}R_0^T x_0^0$ and thus all four surfaces x_0^0 are totally real.

Note that the rigid motion $(I_3 - \frac{z}{2}(f_1 e_3^T - e_3 f_1^T) - \frac{z^2}{8}e_3 e_3^T, \frac{z^3}{16}f_1 - \frac{3z}{2}\bar{f}_1 - \frac{3z^2}{8}e_3) \in \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ brings $x_z(u, v)$ to the canonical form $x_0((u - z), v)$, so x_z^0 has real linear element iff $(u - z, v)$ are as (u, v) above; in particular both $x_0^0,\ x_z^0$ have real linear element iff $z \in \sqrt{\epsilon_1\sqrt{-\epsilon}}\mathbb{R}$, so each of the four surfaces comes with its totally real confocal family. Since $iR(\frac{\pi}{4})x_z(u, v) = x_{\sqrt{i}z}(\sqrt{i}u, iv)$, multiplication by i and the change of ϵ_1 has same effect on the totally real confocal family as that on x_0^0; further since $\overline{R_0^T x_z^0} = \mathcal{E}R_0^T x_z^0$ we conclude that x_0^0 and x_z^0 stay in the same totally real space. Note that the Ivory affinity preserves each totally real confocal family and the collection of these

four totally real confocal families meets each of $\{x_{(\sqrt{i})^j z}\}_{j=0,...7},\ z \in \mathbb{R}^*$ fixed just in one of their four totally real surfaces.

Consider the ruling families parametrization $x_z(\hat{u},\hat{v}) := \frac{z}{2}(-\hat{u}\hat{v}f_1 + (1-\frac{\hat{u}}{z})2\bar{f}_1 + (\hat{u}+\hat{v})e_3)$; note that it differs from the usual ruling families parametrization on x_z invariant under the Ivory affinity (it is obtained by applying to the later the transformation $(u,v) \mapsto (\frac{z}{2}-u, \frac{2}{z}(v-\frac{z^2}{8}))$); for totally real surface x_0^0 and $z \in \sqrt{\epsilon_1\sqrt{-\epsilon}}\mathbb{R}$ we have $(\hat{u},\hat{v}) = (\sqrt{\epsilon_1\sqrt{-\epsilon}}\hat{s}, \frac{\sqrt{-\epsilon}}{\sqrt{\epsilon_1\sqrt{-\epsilon}}}\hat{t}),\ \hat{s},\ \hat{t} \in \mathbb{R}$. Thus $[x_z^0(\hat{u},\hat{v})^T, 1] = [\frac{z}{|z|}(-\hat{u}\hat{v}f_1^T + (1-\frac{\hat{u}}{z})2\bar{f}_1^T + (\hat{u}+\hat{v})e_3^T), \frac{2}{|z|}]$; as $z \to \infty$ in its real line and $\hat{u},\hat{v}$ are fixed we get $[x_\infty^0(\hat{u},\hat{v})^T, 0] = [\sqrt{\epsilon_1\sqrt{-\epsilon}}(-\hat{u}\hat{v}f_1^T + 2\bar{f}_1^T + (\hat{u}+\hat{v})e_3^T), 0] = [(\epsilon\epsilon_1\hat{s}\hat{t}f_1^T + 2\bar{f}_1^T + \sqrt{-\epsilon}(\epsilon_1\hat{s}+\hat{t})e_3^T)R_0^T, 0]$, which is a double cover of the concave region of the plane $[(\mathbb{R}\times i\mathbb{R}\times\sqrt{-\epsilon}\mathbb{R})R_0^T, 0]$ at ∞ of the totally real space $R_0(\mathbb{R}\times i\mathbb{R}\times\sqrt{-\epsilon}\mathbb{R})$. The change of ϵ in the linear element because of multiplication by i thus becomes evident at the infinite singular quadric, where also the change of ϵ_1 by the rotation $-R(\frac{\pi}{2})$ can be seen. The change of ϵ occurs because all real lines $\lambda\mathbb{R}\cup\{\infty\} \subset \mathbb{C}\cup\{\infty\},\ \lambda \in \mathbb{C}\setminus\{0\}$ meet at ∞ and 0; when $z=\infty$ and $\hat{u},\hat{v}$ are finite in their corresponding real lines, u,v are infinite in their corresponding real lines (although $du = -d\hat{u}$ and dv for $d\hat{v}=0$ may be finite). In this case $\hat{u},\hat{v}$ change when changing their real lines, but u,v don't change, so the change in the linear element is continuous from a projective point of view. Note that the above considered lines meet also at 0, which can be considered a locus of change of linear element ($u=v=du=dv=0$), but 0 is not the natural choice, being a point instead of a surface naturally appearing as a limiting case of a family of surfaces.

In the diagram below continuous lines are quadrics of the respective confocal family and with real rulings; to get all four totally real confocal families one must apply to the diagram below a rotation $-R(\frac{\pi}{2})$ and superimpose the two.



For IQWC (3.8), with $U := \frac{u+v}{\sqrt{2}},\ V := \frac{u-v}{\sqrt{2}},\ a := \sqrt{a_1}$ we have $\mathcal{A} = a,\ |dx_0^0|^2 = -a^2dV^2 + dU\odot d(U^2-V^2)$. (I)&(II) become $2aU + \frac{V^2}{a} = \epsilon(2\bar{a}\bar{U} + \frac{\bar{V}^2}{\bar{a}}),\ 2V^2(\frac{U}{a^3}+\frac{1}{a}) - a^3 = \epsilon(2\bar{V}^2(\frac{\bar{U}}{\bar{a}^3}+\frac{1}{\bar{a}}) - \bar{a}^3)$, which imply $(\frac{V^2}{a^4} - \frac{\bar{V}^2}{\bar{a}^4})(2aU + a^3(2-\frac{\bar{V}^2}{\bar{a}^4})) - (a^3-\epsilon\bar{a}^3)(1-\frac{\bar{V}^2}{\bar{a}^4})^2 = 0$. If $a^3 - \epsilon\bar{a}^3 \neq 0$, then one can solve for U as a rational function of $V,\ \bar{V}$; however in this case the condition that the coefficient of $dV\odot d\bar{V}$ in $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is 0 cannot be realized without imposing a real relation between V and $\bar{V}$. This can be seen by considering the hot's in $V,\ \bar{V}$: with $\lambda := \frac{V^2}{a^4} - \frac{\bar{V}^2}{\bar{a}^4}$ we have $\lambda U = \frac{V^2\bar{V}^2}{2|a|^4\bar{a}^2} - \frac{\epsilon\bar{V}^4}{2|a|^2\bar{a}^4} + \text{lot},\ \lambda^2 dU = (\frac{\epsilon\bar{a}^3-a^3}{|a|^{10}\bar{a}^3}V\bar{V}^4 + \text{lot})dV + (\frac{V^4\bar{V}}{|a|^8a^2} - \frac{2\epsilon V^2\bar{V}^3}{|a|^{10}} + \frac{\epsilon\bar{V}^5}{|a|^2\bar{a}^8} + \text{lot})d\bar{V}$, so $\lambda^5((2U\frac{\partial U}{\partial V} - V)\frac{\partial U}{\partial V} - (2\bar{U}\frac{\partial\bar{U}}{\partial V} - \bar{V})\frac{\partial\bar{U}}{\partial V})$ is a polynomial of degree 14 in $V,\ \bar{V}$, with the coefficient of $V\bar{V}^{13}$ being a nonzero multiple of $\epsilon\bar{a}^3 - a^3$ (it is obtained by considering only $\lambda^5 U\frac{\partial U}{\partial V}\frac{\partial U}{\partial V}$). Thus $\epsilon\bar{a}^3 - a^3 = 0$ and (III) further imposes $\frac{V^2}{a^4} - \frac{\bar{V}^2}{\bar{a}^4} = 0$; using again (I) we get $\bar{V} = \epsilon_1\frac{\bar{a}^2}{a^2}V,\ \epsilon_1 := \pm 1,\ \bar{U} = \frac{\epsilon a}{\bar{a}}U$ and (III) is satisfied.

With $\mu := \frac{\epsilon\bar{a}}{a} = \frac{\epsilon|a|^2}{a^2} = \frac{\epsilon|a_1|}{a_1}$ we have $\mu^3 = 1$, so there are 6 types of IQWC (3.8) $x_0 = x_0(u,v) = uvf_1 + \frac{u+v}{\sqrt{2}}\bar{f}_1 + i\sqrt{\epsilon\mu^2|a_1|}\frac{u-v}{\sqrt{2}}e_3$ containing surfaces x_0^0 with real linear element; each contains two surfaces x_0^0, given by $(u,v) = (\sqrt{\mu}s, \sqrt{\mu}t),\ s,t \in \mathbb{R}$ and $v = \mu\bar{u},\ u \in \mathbb{C}$. In the first case the sign $(\sqrt{\mu})^3$ in $|dx_0^0|^2$ can be absorbed by $(s,t) \mapsto (-s,-t)$; in the second case μ in $|dx_0^0|^2$ can be absorbed by $u \mapsto \mu^{-1}u$, so we have four types of real linear element of IQWC (3.8) from a real point of view (namely $-\epsilon|a_1|\frac{(d(s-t))^2}{2} + \sqrt{2}d(st)\odot d(s+t)$ and $-\epsilon|a_1|\frac{(d(u-\bar{u}))^2}{2} + \sqrt{2}d|u|^2\odot d(u+\bar{u})$); from a complex point of view we have only two types, since further ϵ can be absorbed by $a_1 \mapsto -a_1$. We have $-R(\pi)x_0(u,v) = x_0(v,u)$; the rigid motions preserving the canonical form account for the

choice of μ as a cubic root of unity: $R(\frac{2\pi k}{3})x_0(u,v) = (-R(\pi))^j \hat{x}_0(-\sqrt{\hat{\mu}}u, -\sqrt{\hat{\mu}}v),\ \sqrt{\hat{\mu}}\bar{u} = \hat{\mu}\overline{\sqrt{\hat{\mu}}u}$, where $k = 1, 2,\ \mu := 1,\ \hat{\mu} := e^{\frac{2\pi i k}{3}},\ \sqrt{\hat{a}_1} := (-1)^{j+1}(\sqrt{\hat{\mu}})^{-1}\sqrt{a_1},\ \hat{\epsilon} := \epsilon$; j is chosen so that $\sqrt{\hat{a}_1}$ is a square root. Multiplication by i changes ϵ while preserving μ; it is enough to choose $\mu := 1,\ a_1 > 0 :\ iR(-\frac{\pi}{2})x_0(u,v) = \hat{x}_0(-u,-v))$, where $\hat{\mu} := 1,\ \hat{a}_1 := -a_1$.

In the first case $|dx_0^0|^2$ has signature $(1,1)$ for $2\sqrt{2}\epsilon|a_1|(\sqrt{\mu})^3(s+t) > -(s-t)^2$, respectively $\mathrm{diag}[-\epsilon\ -\epsilon]$ for the other inequality; in the second case $|dx_0^0|^2$ has signature $(1,1)$ for $2\sqrt{2}\epsilon|a_1|(\mu u + \bar{\mu}\bar{u}) < -(\mu u - \bar{\mu}\bar{u})^2$, respectively $\mathrm{diag}[\epsilon\ \epsilon]$ for the other inequality. We can make in what follows $\mu = 1,\ a_1 = \epsilon|a_1|$. We have $\bar{x}_0^0 = \mathcal{E}x_0^0$, where in the first case $\mathcal{E} := \mathrm{diag}[1\ -1\ -\epsilon]$ and in the second case $\mathcal{E} := \mathrm{diag}[1\ -1\ \epsilon]$; thus x_0^0 are totally real.

Since $x_z(u,v) - z\bar{f}_1 = (-R(\pi))^j \hat{x}_0(u - \frac{z}{2\sqrt{2}}, v - \frac{z}{2\sqrt{2}}),\ \sqrt{\hat{a}_1} := (-1)^j\sqrt{a_1}\sqrt{1 - za_1^{-1}}$ is the canonical form of x_z, x_z^0 is totally real iff $\hat{a}_1 \in \mathbb{R}$; in particular both $x_0^0,\ x_z^0$ are totally real iff $z \in \mathbb{R}$; in both cases $\hat{\mu} = \mu = 1$, but $\hat{\epsilon}$ may be different than ϵ (precisely when $1 - za_1^{-1} < 0$). In both cases since $\bar{x}_z^0 = \hat{\mathcal{E}}x_z^0$ we conclude that x_z^0 stays in the same totally real space as that of x_0^0 if $1 - za_1^{-1} > 0$ and stays in a totally real space which differs from that of x_0^0 by a multiplication with i in the third coordinate if $1 - za_1^{-1} < 0$; the Ivory affinity preserves each totally real confocal family. For $z = a_1$ we have the singular totally real quadric $x_{a_1}^0 = \mathbb{R} \times i\mathbb{R} = \{m_1 f_1 + m_2 \bar{f}_1 | m_1,\ m_2 \in \mathbb{R}\}$; the concave domain given by the inequality $(m_2 - a_1)^2 - 2m_1 > 0$ is realized as a singular totally real quadric with linear element of the first case type $(x_{a_1}^0(s,t) = stf_1 + \frac{s+t}{\sqrt{2}}\bar{f}_1 + a_1\bar{f}_1,\ s,t \in \mathbb{R})$ and the convex domain given by the other inequality is realized as a singular totally real quadric with linear element of the second case type $(x_{a_1}^0(u,\bar{u}) = |u|^2 f_1 + \frac{u+\bar{u}}{\sqrt{2}}\bar{f}_1 + a_1\bar{f}_1,\ u \in \mathbb{C})$; both domains meet in the singular part $\mathcal{S}(x_{a_1}^0)(m_2) := \frac{(m_2-a_1)^2}{2}f_1 + m_2\bar{f}_1$ for $s = t = u = \bar{u} := \pm\frac{m_2 - a_1}{\sqrt{2}}$. The concavity (convexity) can be easily seen because of the parametrization by two real (complex conjugate) parameters s, t $(u,\ \bar{u})$. Note again that the change of μ due to the group generated by $R(\frac{2\pi}{3})$ or the multiplication by i has same effect on the confocal family as that on x_0^0; it is enough to consider $\mu := 1,\ z \in \mathbb{R}$, when $\hat{z} = -\sqrt{\hat{\mu}}z$.

The homography of Bianchi II previously considered and its composition with a homothety i takes the totally real IQWC (3.8) $x_0^0(u,v)$ to the QC (3.5) $\tilde{x}_0^0(u,v) := \pm\sqrt{\epsilon_2}(\sqrt{a_1(I_3 + a_1 J_3)^{-1}})^{-1}X(\frac{a_1}{2} - \sqrt{2}u, \frac{a_1}{2} - \sqrt{2}v),\ \epsilon_2 := \pm 1$, so it has the linear element of the image $|d\tilde{x}_0^0|^2 = \frac{2\epsilon_2}{(u-v)^4}(a_1(d(u-v))^2 - 2\sqrt{2}(vdu - udv) \odot d(u-v) + 2a_1^{-1}(u-v)^2 du \odot dv)$, which is real only for $\mu := 1$; for $\mu \neq 1$ it is real after an imaginary homothety. We have $z \in \mathbb{R} \Rightarrow \tilde{z} \in \mathbb{R}$, so these homographies take totally real confocal quadrics to totally real confocal quadrics; as we shall see below all totally real QC (3.5) appear as above. The discussion of the infinite singular quadric of IQWC (3.8) (where either ϵ or the two cases change) is unnecessary at this point, since it will correspond via the above considered homographies to the discussion of the finite singular quadric for QC (3.5); also at that time the discussion of the corresponding infinite singular quadric will not be necessary, it corresponding to the already completed discussion of the finite singular quadric for IQWC (3.8). In the diagram below $a_1 = |a_1|$, (dis)continuous lines represent the quadrics of the respective confocal families having (imaginary) real rulings; the spectral parameter z $(-z)$ increases with the arrow.



For QC (3.5) with $a := (\sqrt{a_1^{-1}})^{-1},\ V := a^4(u-v)^2,\ U := 2\frac{1-a^2(u+v)}{V} + \frac{1}{2}$ we have $\mathcal{A} = ia^3$, $|dx_0^0|^2 = \frac{a^2}{V}((\frac{d(UV)}{2} - \frac{dV}{4} + \frac{dV}{V})^2 - \frac{U}{2V}dV^2)$. (I)&(II) become $aU = -\epsilon\bar{a}\bar{U},\ a((U+\frac{2}{V}-\frac{1}{2})^2 - 2\frac{U}{V}) = -\epsilon\bar{a}((\bar{U}+\frac{2}{V}-\frac{1}{2})^2 - 2\frac{\bar{U}}{V})$, which imply $f(U,V,\bar{V}) := (1+\frac{\epsilon a}{\bar{a}})(\frac{\epsilon a}{\bar{a}}U^2 + 2(\frac{1}{V}-\frac{1}{\bar{V}})U + (\frac{2}{V}-\frac{1}{2})^2) - 2(\frac{1}{V}-\frac{1}{\bar{V}})(U+\frac{2}{V}+\frac{2}{\bar{V}}-1) = 0$. If $1+\frac{\epsilon a}{\bar{a}} \neq 0$, then $\frac{\partial f}{\partial U} \neq 0$ and one can solve for dU from $df = 0$; the condition that the coefficient of $dV \odot d\bar{V}$ in $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is 0 becomes the cubic equation $\frac{\epsilon a}{\bar{a}}(U\frac{\partial f}{\partial U}(\frac{\partial f}{\partial V} - \frac{\partial f}{\partial \bar{V}}) + (V-\bar{V})\frac{\partial f}{\partial V}\frac{\partial f}{\partial \bar{V}} - \frac{\partial f}{\partial U}\frac{\partial f}{\partial \bar{V}}(\frac{2}{V}-\frac{1}{2})) - \frac{\partial f}{\partial U}\frac{\partial f}{\partial \bar{V}}(\frac{2}{\bar{V}}-\frac{1}{2}) = 0$ in U, which together with the quadratic equation $f(U,V,\bar{V}) = 0$ in U imposes a real relation between V and $\bar{V}$.

Thus $\bar{a} + \epsilon a = 0$, when (III) implies $\bar{V} = V$ (the condition $\bar{V} \neq V$ imposes $U = 1 - \frac{2}{V} - \frac{2}{\bar{V}} = \bar{U}$ and a functional relationship between V and $\bar{V}$); finally $a_1 = -\epsilon|a_1|,\ (u,v) = (s,t),\ s,t \in \mathbb{R}$ or $v = \bar{u},\ u \in \mathbb{C}$.

We have two types of QC (3.5) $x_0 = x_0(u,v) = \frac{\sqrt{|a_1|}}{\sqrt{-\epsilon}(u-v)}(-(u+\frac{\epsilon}{2|a_1|})(v+\frac{\epsilon}{2|a_1|})f_1 + 2\bar{f}_1 + (u+v-\frac{\epsilon}{|a_1|})e_3)$ containing surfaces with real linear element; each contains two such surfaces x_0^0. Thus we have four types of real linear element of QC (3.5) from a real point of view (namely $8\frac{(tds-sdt)\odot d(s-t)}{(s-t)^4} - 4\epsilon|a_1|\frac{ds\odot dt}{(s-t)^2},\ 8\frac{(\bar{u}du-ud\bar{u})\odot d(u-\bar{u})}{(u-\bar{u})^4} - 4\epsilon|a_1|\frac{du\odot d\bar{u}}{(u-\bar{u})^2}$); from a complex point of view we have only two, since ϵ can be absorbed by $a_1 \mapsto -a_1$. We have $ix_0^0(u,v) = -\epsilon(I_3 - 2e_3e_3^T)\hat{x}_0^0(-u,-v)$, $\hat{a}_1 := -a_1$, so $\hat{\epsilon} := -\epsilon$.

In the first case $|dx_0^0|^2$ has signature $(1,1)$ for $4(1+\epsilon|a_1|(s+t)) > -|a_1|^2(s-t)^2$, respectively $\mathrm{diag}[-\epsilon\ -\epsilon]$ for the other inequality; in the second case $|dx_0^0|^2$ has signature $(1,1)$ for $4(1+\epsilon|a_1|(u+\bar{u})) < -|a_1|^2(u-\bar{u})^2$, respectively $\mathrm{diag}[\epsilon\ \epsilon]$ for the other inequality.

With $\epsilon' := \mathrm{diag}[-\epsilon\ \epsilon\ -\epsilon]$ in the first case or $:= \mathrm{diag}[\epsilon\ -\epsilon\ \epsilon]$ in the second case we have $\bar{x}_0^0 = \epsilon' x_0^0$, so x_0^0 are totally real.

Now x_z contains totally real surfaces iff $a_1 - z \in \mathbb{R}$; in particular both $x_0^0,\ x_z^0$ are totally real iff $a_1, z \in \mathbb{R}$, in which case $x_z^0(u,v) = \hat{x}_0^0(u,v)$ for $1 - za_1^{-1} > 0$ and $= \epsilon\hat{x}_0^0(u,v)$ for the other inequality, where $\hat{a}_1 := a_1 - z$. In both cases, since $\bar{x}_z^0 = \hat{\epsilon}' x_z^0$, we conclude that x_z^0 stays in the same totally real space as that of x_0^0 if $1 - za_1^{-1} > 0$ and stays in a totally real space which differs from that of x_0^0 by a multiplication with i for the other inequality; the Ivory affinity preserves each confocal family.

For QWC (3.7) with $a := \sqrt{a_1^{-1}},\ U := uv,\ V := v^2$ we have $\mathcal{A} = ia^{-2},\ |dx_0^0|^2 = (d(uv))^2 + dv^2 + \frac{2}{a^2}du \odot dv = dU^2 + \frac{dV^2}{4V} + \frac{1}{a^2}dV \odot (\frac{dU}{V} - \frac{UdV}{2V^2})$. (I)&(II) become $a^2(2a^2U - a^4V + 1) = -\epsilon\bar{a}^2(2\bar{a}^2\bar{U} - \bar{a}^4\bar{V} + 1),\ a^{12}(2U - a^2V(3+a^4V)) = -\epsilon\bar{a}^{12}(2\bar{U} - \bar{a}^2\bar{V}(3+\bar{a}^4\bar{V}))$, which imply $2a^4(a^8 - \bar{a}^8)U - a^{14}V(3+a^4V) - \epsilon\bar{a}^{14}\bar{V}(3+\bar{a}^4\bar{V}) + a^2\bar{a}^8(a^4V-1) + \epsilon\bar{a}^2a^8(\bar{a}^4\bar{V}-1) = 0$. If $a^8 - \bar{a}^8 \neq 0$, then (III) cannot be satisfied without imposing at least a real condition between V and $\bar{V}$; thus $a^8 = \bar{a}^8$, from which also follows $a^2(a^4V+1)^2 + \epsilon\bar{a}^2(\bar{a}^4\bar{V}+1)^2 = 0$ and (I)&(II) are equivalent, when x_0^0 exists, to

$$(I)'\quad a^2(2a^2U - a^4V + 1) = -\epsilon\bar{a}^2(2\bar{a}^2\bar{U} - \bar{a}^4\bar{V} + 1),$$
$$(II)'\quad a^2(a^4V+1)^2 = -\epsilon\bar{a}^2(\bar{a}^4\bar{V}+1)^2.$$

Now $|dx_0^0|^2 = (d(uv))^2 + dv^2 + \frac{2}{a^2}du \odot dv = dU^2 + \frac{dV^2}{4V} + \frac{1}{a^2}dV \odot (\frac{dU}{V} - \frac{UdV}{2V^2})$. Applying d to (I)'&(II)' we get $-\epsilon d\bar{V} = \frac{a^6(a^4V+1)}{\bar{a}^6(\bar{a}^4\bar{V}+1)}dV,\ -\epsilon d\bar{U} = \frac{a^4}{\bar{a}^4}dU + \frac{a^6(\bar{a}^4\bar{V} - a^4V)}{2\bar{a}^4(\bar{a}^4\bar{V}+1)}dV$, so (III) becomes the condition that a symmetric bilinear form in $dU,\ dV$ is identically 0. In particular the condition that the coefficient of $dU \odot dV$ is 0 gives $(a^4V - \bar{a}^4\bar{V})((a^4V-1)(\bar{a}^4\bar{V}-1) - 2) = 0$, which together with (II)' implies $a^4V = \bar{a}^4\bar{V}$, so finally $\bar{a}_1 = -\epsilon a_1, \bar{v} = -\epsilon\epsilon_1 v,\ \bar{u} = \epsilon_1 u,\ \epsilon_1 := \pm 1$ and (III) is satisfied.

Thus we have two types of QWC (3.7) $x_0 = x_0(u,v) = (\sqrt{a_1^{-1}})^{-1}((u+\frac{v}{2a_1})f_1 + v\bar{f}_1) + uve_3$ containing surfaces x_0^0 with real element (given by the choice of ϵ in $a_1 = (\sqrt{-\epsilon})^{-1}\epsilon_2|a_1|,\ \epsilon_2 := \pm 1$); each contains two surfaces x_0^0 (given by the choice of ϵ_1 in $(u,v) = (\sqrt{\epsilon_1}s, \sqrt{-\epsilon}\sqrt{\epsilon_1}t),\ s,t \in \mathbb{R}$). The transformation $u \mapsto -u$ changes ϵ_2 in $|dx_0^0|^2$, so we have from a real point of view four different

types of real linear element of QWC (3.7) (namely $|dx_0^0|^2 = -\epsilon((d(st))^2 + \epsilon_1 dt^2 + 2|a_1|ds \odot dt)$); from a complex point of view we have only one type, since $(s,t) \mapsto (is,-it)$ changes ϵ_1 and $(a_1,s,t) \mapsto (ia_1,is,-it)$ changes ϵ. All symmetries of the linear element are realized by rigid motions composed when necessary with a multiplication by i or with a simple complex affine transformation: $R(\pi)x_0(u,v) = x_0(-u,-v)$, $(-i\epsilon\epsilon_2 f_1\bar{f}_1^T + i\epsilon\epsilon_2\bar{f}_1 f_1^T - e_3e_3^T)x_0(u,v) = \hat{x}_0(-u,v)$, where $\hat{a}_1 := -a_1$, $i(\frac{\sqrt{\epsilon_2\sqrt{\epsilon}}}{\sqrt{\epsilon_2\sqrt{-\epsilon}}}f_1\bar{f}_1^T + \frac{\sqrt{\epsilon_2\sqrt{-\epsilon}}}{\sqrt{\epsilon_2\sqrt{\epsilon}}}\bar{f}_1 f_1^T + \epsilon e_3e_3^T)x_0(u,v) = \hat{x}_0(\epsilon u, iv)$, where $\hat{a}_1 := i\epsilon a_1$, $\mathrm{diag}[i \;\; i \;\; -1](\frac{\sqrt{\epsilon\epsilon_2\sqrt{-\epsilon}}}{\sqrt{\epsilon_2\sqrt{-\epsilon}}}f_1\bar{f}_1^T + \frac{\sqrt{\epsilon_2\sqrt{-\epsilon}}}{\sqrt{\epsilon\epsilon_2\sqrt{-\epsilon}}}\bar{f}_1 f_1^T + \epsilon e_3e_3^T)x_0(u,v) = \hat{x}_0(i\epsilon u, iv)$, where $\hat{a}_1 := \epsilon a_1$.

Now $|dx_0^0|^2$ has signature $(1,1)$ for $\epsilon_1 t^2 - 2\epsilon_1\epsilon_2 st|a_1| - |a_1|^2 < 0$ and $\mathrm{diag}[-\epsilon \;\; -\epsilon]$ for the other inequality. With $\epsilon' := \mathrm{diag}[1 \;\; -1 \;\; -\epsilon]$, $R_0 := \sqrt{\frac{\epsilon_1\epsilon_2}{\sqrt{-\epsilon}}}f_1\bar{f}_1^T + (\sqrt{\frac{\epsilon_1\epsilon_2}{\sqrt{-\epsilon}}})^{-1}\bar{f}_1 f_1^T + e_3e_3^T$ we have $\overline{R_0^T x_0^0} = \epsilon' R_0^T x_0^0$, so all x_0^0 are totally real.

Since $x_z(u,v) - \frac{z}{2}e_3 = R(\pi)^j\hat{x}_0(u,v)$, where $\hat{a}_1 := a_1 - z$, $j := 0,1$ is the canonical form of x_z, x_z^0 is totally real iff $a_1 - z$ is as a_1 above; in particular both x_0^0, x_z^0 are totally real iff $\bar{z} = -\epsilon z$. In this case $j = 0$ for $\hat{\epsilon}_2 = \epsilon_2$ or $\epsilon\epsilon_2 = 1$ and $j = 1$ otherwise; since $\overline{R_0^T x_z^0} = \epsilon' R_0^T x_z^0$ for $\hat{\epsilon}_2 = \epsilon_2$, it is natural to require that the whole confocal family should stay in the same totally real space $R_0(\mathrm{R}) \times i\mathbb{R} \times (\sqrt{-\epsilon}\mathbb{R})$); this will suffice to conclude $\hat{\epsilon}_1 = \epsilon_1\epsilon_2\hat{\epsilon}_2$. The rigid motions considered above and the multiplication by i have the same effect on the totally real confocal family x_z^0 as that on x_0^0 (ϵ_2 is not replaced with $\hat{\epsilon}_2$ in the definition of these rigid motions); however the affine transformation considered above cannot have the same effect (as we know that the only affine transformations which take confocal quadrics to confocal quadrics are homotheties and rigid motions): it involves a translation $-\epsilon z e_3$ on the rhs term and ϵ_2 is replaced with $\hat{\epsilon}_2$. Note that the Ivory affinity does not preserve the totally real confocal family x_z^0; for $\epsilon_2\hat{\epsilon}_2 = -1$ it mixes the cases $\epsilon_2 := 1$ and $\epsilon_2 := -1$, but according to the above affine transformation (or variants thereof). When z varies with $\bar{z} = -\epsilon z$ and $\hat{a}_1$ changes sign, $\hat{\epsilon}_1$ changes. Again this can be explained if one considers the singular quadric x_{a_1}.

Consider the ruling families parametrization $x_z^0(\hat{u},\hat{v}) := (\hat{u} + \frac{\hat{v}}{2})f_1 + (a_1 - z)\hat{v}\bar{f}_1 + (\hat{u}\hat{v} + \frac{z}{2})e_3$; note that it differs from the usual ruling families parametrization on x_z^0 invariant under the Ivory affinity (it is obtained by applying to the later the transformation $(u,v) \mapsto (\sqrt{1 - za_1^{-1}}(\sqrt{a_1^{-1}})^{-1}u, \frac{(\sqrt{a_1^{-1}})^{-1}}{a_1\sqrt{1-za_1^{-1}}}v)$, which, as $z \to a_1$, contracts u and expands v), but in a way such that $\hat{u}$, $\hat{v}$ vary in real lines, if $\bar{z} = -\epsilon z$ and u, v vary in their corresponding real lines. Moreover, the real lines of $\hat{u}$, $\hat{v}$ are independent of the sign of $1 - za_1^{-1}$, since ϵ_1 changes sign with $1 - za_1^{-1}$. Multiplying by i and applying further rigid motions, if necessary, it is enough to consider only the case $a_1 > 0$, $\epsilon_1 := 1$, when we have $(\hat{u},\hat{v}) = (\hat{s},\hat{t})$, $\hat{s},\hat{t} \in \mathbb{R}$. In this case $x_{a_1}^0(\hat{s},\hat{t}) = (\hat{s} + \frac{\hat{t}}{2})f_1 + (\hat{s}\hat{t} + \frac{a_1}{2})e_3$ appears as a singular case for both types of totally real linear element $|dx_z^0|^2$ with $1 - za_1^{-1} > 0$ and $1 - za_1^{-1} < 0$, it has degenerate (isotropic) linear element $(d(\hat{s}\hat{t}))^2$, covers twice the concave domain $\{m_1f_1 + m_2e_3 | m_1,m_2 \in \mathbb{R},\ m_1^2 - 2m_2 + a_1 > 0\}$ and meets $\mathcal{S}(x_{a_1}) = \{m_1f_1 + m_2e_3 | m_1,m_2 \in \mathbb{C},\ m_1^2 - 2m_2 + a_1 = 0\}$ for $\hat{s} = \frac{\hat{t}}{2}$. Although u,v change their real lines for $z \neq a_1$, they do not change it for $z = a_1$, when $u = \infty$, $v = 0$, $uv = \hat{u}\hat{v}$, $du = \infty$ if $d\hat{u} \neq 0$ and $dv = 0$ if $d\hat{v} \neq \infty$.

The homography of Bianchi II previously considered already takes totally real confocal quadrics QWC (3.7) to same type of quadrics: $\tilde{x}_0^0(u,v) = \sqrt{a_1^{-1}}(uf_1 - \frac{1}{v}\bar{f}_1) - \frac{1}{2\sqrt{a_1^{-1}v}}f_1 - \frac{u}{v}e_3$; it and its composition with a homothety i are the only such homographies. We thus have $\tilde{a}_1 = a_1^{-1}$, $\tilde{u} = u$, $\tilde{v} = -v^{-1}$; for $(\sqrt{a_1})^{-1} = -\sqrt{a_1^{-1}}$ or multiplication by i further rigid motions may be required.

Again the discussion of the infinite singular quadric of QWC (3.7) (where ϵ may change) is not necessary, since it corresponds via the above considered homographies to the already completed discussion of the finite singular quadric of QWC (3.7).

For QWC (3.6) with $a := (\sqrt{a_1^{-1}})^{-1},\ b := -i(\sqrt{a_2^{-1}})^{-1},\ \mathcal{U} := \frac{u+v}{\sqrt{2}},\ \mathcal{V} := \frac{u-v}{\sqrt{2}}$ we have $x_0^0 = a\mathcal{U}e_1 + b\mathcal{V}e_2 + \frac{\mathcal{U}^2-\mathcal{V}^2}{2}e_3,\ \hat{N}_0^0 = \frac{\mathcal{U}}{a}e_1 - \frac{\mathcal{V}}{b}e_2 - e_3,\ A\hat{N}_0^0 = \frac{\mathcal{U}}{a^3}e_1 + \frac{\mathcal{V}}{b^3}e_2,\ V_2 = \frac{\mathcal{V}}{b^3}e_1 - \frac{\mathcal{U}}{a^3}e_2 + \frac{a^2+b^2}{a^3b^3}\mathcal{U}\mathcal{V}e_3,\ \mathcal{A} = -ab.$ With $(U,V) := (\mathcal{U}^2,\mathcal{V}^2)$ (I)&(II) become $\frac{b}{a}U + \frac{a}{b}V + ab = \epsilon(\frac{\bar{b}}{\bar{a}}\bar{U} + \frac{\bar{a}}{\bar{b}}\bar{V} + \bar{a}\bar{b}),\ \frac{b^3}{a^3}U + \frac{a^3}{b^3}V + \frac{(a^2+b^2)^2}{a^3b^3}UV = \epsilon(\frac{\bar{b}^3}{\bar{a}^3}\bar{U} + \frac{\bar{a}^3}{\bar{b}^3}\bar{V} + \frac{(\bar{a}^2+\bar{b}^2)^2}{\bar{a}^3\bar{b}^3}\bar{U}\bar{V})$, which imply $\frac{b}{a}U(\frac{b^2}{a^2} - \frac{\bar{b}^2}{\bar{a}^2} + \frac{(a^2+b^2)^2}{a^3b^3}\frac{a}{b}V - \frac{(\bar{a}^2+\bar{b}^2)^2}{\bar{a}^3\bar{b}^3}\frac{\bar{a}}{\bar{b}}\bar{V}) + \frac{a^3}{b^3}V - \epsilon\frac{\bar{a}^3}{\bar{b}^3}\bar{V} - (\frac{\bar{b}^2}{\bar{a}^2} + \frac{(\bar{a}^2+\bar{b}^2)^2}{\bar{a}^3\bar{b}^3}\frac{\bar{a}}{\bar{b}}\bar{V})(\frac{a}{b}V + ab - \epsilon(\frac{\bar{a}}{\bar{b}}\bar{V} + \bar{a}\bar{b})) = 0.$ If $a^2+b^2 = 0$, then also $ab = \epsilon\bar{a}\bar{b}$; since both a^{-1} and $-ib^{-1}$ are square roots, we get $b = -ia$ and $a^2 = -\epsilon\bar{a}^2$; now (I) becomes $\bar{U} - \bar{V} = -\epsilon(U-V)$, or $\bar{u}\bar{v} = -\epsilon uv$. Keeping account of $|dx_0^0|^2 = \frac{1}{4}(a^2\frac{dU^2}{U} + b^2\frac{dV^2}{V} + (dU-dV)^2)$, (III) factors as $d\log\frac{\bar{v}}{v} \odot d\log\frac{\bar{v}}{u} = 0$, so we have two types of QWC (3.6) with $a_1 = a_2$ containing surfaces x_0^0 with real linear element (given by ϵ in $a_1 = \epsilon_2(\sqrt{-\epsilon})^{-1}|a_1|,\ \epsilon_2 := \pm 1$); each type contains three surfaces x_0^0, according to $(u,v) = (\sqrt{-\epsilon}e^{ic}s, e^{-ic}t),\ s,t \in \mathbb{R}$ or $v = e^{-2c}\epsilon_1\sqrt{-\epsilon}\bar{u},\ u \in \mathbb{C},\ \epsilon_1 := \pm 1$; in both cases $c \in \mathbb{R}$ is a constant. In the first case $s \mapsto -s$ changes ϵ_2 in $|dx_0^0|^2$; in the second case c can be absorbed by $u \mapsto e^{-c}u$, so we have 6 types of real linear element of QWC (3.6) with $a_1 = a_2$ (namely $2|a_1|ds \odot dt - \epsilon(d(st))^2$ or $2\epsilon_1\epsilon_2|a_1|du \odot d\bar{u} - \epsilon(d|u|^2)^2$); from a complex point of view we have only two types, since the transformation $(a_1,u) \mapsto (i\epsilon a_1, i\epsilon u)$ changes ϵ in the first case or $\epsilon,\ \epsilon_1$ in the second one and $a_1 \mapsto -a_1$ changes ϵ_2 in the second case. All symmetries of the linear element are realized by rigid motions composed when necessary with a multiplication by i or a simple complex affine transformation: $R(-c)$ in the first case or $R(ic)$ in the second one makes $c = 0,\ (I_3 - 2e_2e_2^T)x_0(u,v) = x_0(v,u),\ (\frac{\sqrt{-\epsilon_2\sqrt{-\epsilon}}}{\sqrt{\epsilon_2\sqrt{-\epsilon}}}f_1\bar{f}_1^T + \frac{\sqrt{\epsilon_2\sqrt{-\epsilon}}}{\sqrt{-\epsilon_2\sqrt{-\epsilon}}}\bar{f}_1f_1^T - e_3e_3^T)x_0(u,v) = \hat{x}_0(-u,v)$, where $\hat{a}_1 := -a_1$, so it changes ϵ_2 in the first case and both $\epsilon_1,\ \epsilon_2$ in the second case; $ix_0(u,v) = (-i\frac{\sqrt{\epsilon_2\sqrt{-\epsilon}}}{\sqrt{\epsilon_2\sqrt{\epsilon}}}f_1\bar{f}_1^T + i\frac{\sqrt{\epsilon_2\sqrt{\epsilon}}}{\sqrt{\epsilon_2\sqrt{-\epsilon}}}\bar{f}_1f_1^T + \epsilon e_3e_3^T)\hat{x}_0(i\epsilon u, v)$, where $\hat{a}_1 := i\epsilon a_1$ so multiplication by i changes, up to a rigid motion, ϵ in the first case and $\epsilon,\ \epsilon_1$ in the second one; also $\mathrm{diag}[i\ i\ 1]x_0^0(u,v) = \hat{x}_0^0(\epsilon\epsilon_2 u, \epsilon\epsilon_2 v),\ \hat{a}_1 := -a_1$, so it changes ϵ_2.

In the first case $|dx_0^0|^2$ has signature $(1,1)$ for $2\epsilon\epsilon_2 st < |a_1|$ (respectively $\mathrm{diag}[-\epsilon\ \ -\epsilon]$ for the other inequality); in the second case we have signature $(1,1)$ for $2\epsilon\epsilon_1\epsilon_2 e^{-2c}|u|^2 > |a_1|$ (note that we need in this case $\epsilon\epsilon_1\epsilon_2 = 1$), respectively $\mathrm{diag}[\epsilon_1\epsilon_2\ \ \epsilon_1\epsilon_2]$ for the other inequality. In the first case with $R_0 := \sqrt{\epsilon_2\sqrt{-\epsilon}}f_1\bar{f}_1^T + \frac{1}{\sqrt{\epsilon_2\sqrt{-\epsilon}}}\bar{f}_1f_1^T + e_3e_3^T,\ \epsilon' := \mathrm{diag}[1\ \ -1\ \ -\epsilon]$ we have $\overline{R_0^Tx_0^0} = \epsilon' R_0^Tx_0^0$; in the second case with $\epsilon' := \mathrm{diag}[\epsilon_1\epsilon_2\ \ \epsilon_1\epsilon_2\ \ -\epsilon]$ we have $\bar{x}_0^0 = \epsilon' x_0^0$, so all surfaces x_0^0 are totally real.

Since $x_z(u,v) - \frac{z}{2}e_3 = \hat{x}_0(u,v),\ \hat{a}_1 := a_1 - z$ is the canonical form of x_z, x_z^0 is totally real iff $\hat{a}$ is as a above; in particular both $x_0^0,\ x_z^0$ are totally real iff $z \in \sqrt{-\epsilon}\mathbb{R}$ (although ϵ_2 may change).

If $a^2+b^2 \neq 0$, then with $\lambda := \frac{(a^2+b^2)^2}{a^3b^3}$ the consequence of (I)&(II) can be written as $(\lambda\frac{b}{a}U - \bar{\lambda}\frac{\bar{a}}{\bar{b}}\bar{V} + \frac{a^2}{b^2} - \frac{\bar{b}^2}{\bar{a}^2})(\frac{b^2}{a^2} - \frac{\bar{b}^2}{\bar{a}^2} + \lambda\frac{a}{b}V - \bar{\lambda}\frac{\bar{a}}{\bar{b}}\bar{V}) - (\bar{\lambda} - \epsilon\lambda)\frac{(\bar{V}(\bar{a}^2+\bar{b}^2)+\bar{b}^4)^2}{\bar{a}\bar{b}^5} = 0$. If $\bar{\lambda} \neq \epsilon\lambda$, then one can solve the previous for U as a rational function of $V,\ \bar{V}$, but (III) cannot be satisfied without imposing at least a real relation between V and $\bar{V}$. This can be seen by looking at the hot's in $U,\ V,\ \bar{V}$; with $\mu := \lambda\frac{a}{b}V - \bar{\lambda}\frac{\bar{a}}{\bar{b}}\bar{V}$ we have: $\lambda\frac{b}{a}U\mu = \bar{\lambda}\frac{\bar{a}}{\bar{b}}\bar{V}(\mu + (\bar{\lambda} - \epsilon\lambda)\frac{\bar{a}}{\bar{b}}\bar{V}) + \mathrm{lot},\ \lambda\frac{b}{a}\mu^2dU = \bar{\lambda}\frac{\bar{a}}{\bar{b}}(\mu d(\bar{V}(\mu + (\bar{\lambda} - \epsilon\lambda)\frac{\bar{a}}{\bar{b}}\bar{V})) - \bar{V}(\mu + (\bar{\lambda} - \epsilon\lambda)\frac{\bar{a}}{\bar{b}}\bar{V})d\mu) + \mathrm{lot}$. If we ignore the lot's, the hot of the numerator of the coefficient of $dV \odot d\bar{V}$ in $|dx_0^0|^2 - |d\bar{x}_0^0|^2$ is a homogeneous polynomial of degree 4 in V and $\bar{V}$, namely $(\mu^2\frac{\partial U}{\partial V})(\mu^2\frac{\partial U}{\partial V}) - \mu^2(\mu^2\frac{\partial U}{\partial V}) - (\mu^2\frac{\partial\bar{U}}{\partial V})(\mu^2\frac{\partial\bar{U}}{\partial V}) + \mu^2(\mu^2\frac{\partial\bar{U}}{\partial V})$; the coefficient of $\bar{V}^4$ is $-\epsilon(a^2+b^2)(\bar{\lambda} - \epsilon\lambda)\frac{a\bar{a}^5\bar{\lambda}^3}{b^3\bar{b}^5} \neq 0$, so (III) imposes a functional relationship between V and $\bar{V}$. Thus $\bar{\lambda} = \epsilon\lambda$, when we have either $\frac{b}{a}U - \epsilon\frac{\bar{a}}{\bar{b}}\bar{V} + \frac{1}{\lambda}(\frac{a^2}{b^2} - \frac{\bar{b}^2}{\bar{a}^2}) = 0$ or $\frac{b^2}{a^2} - \frac{\bar{b}^2}{\bar{a}^2} + \lambda(\frac{a}{b}V - \epsilon\frac{\bar{a}}{\bar{b}}\bar{V}) = 0$. In the later case (I)&(II) are equivalent to

$$(I)'\quad \frac{a}{b}V - \epsilon\frac{\bar{a}}{\bar{b}}\bar{V} = \frac{1}{\lambda}(\frac{\bar{b}^2}{\bar{a}^2} - \frac{b^2}{a^2}),$$

$$(II)' \quad \frac{b}{a}U - \epsilon\frac{\bar b}{\bar a}\bar U = \frac{1}{\lambda}(\frac{\bar a^2}{\bar b^2} - \frac{a^2}{b^2}).$$

Applying d to (I)'&(II)' we get $\frac{a}{b}dV - \epsilon\frac{\bar a}{\bar b}d\bar V = \frac{b}{a}dU - \epsilon\frac{\bar b}{\bar a}d\bar U = 0$; now (III) imposes $\frac{a}{\bar a} = \epsilon\frac{\bar b}{b} = \pm 1$; finally we get $a_1 = \epsilon_1|a_1|,\ a_2 = -\epsilon\epsilon_1|a_2|,\ \epsilon_1 := \pm 1,\ (u,v) = (\sqrt{\epsilon_2}s, \sqrt{\epsilon_2}t),\ s,t \in \mathbb{R}$ or $v = \epsilon_2\bar u,\ u \in \mathbb{C},\ \epsilon_2 := \pm 1$.

In the former case $\frac{b}{a}U - \epsilon\frac{\bar a}{b}\bar V + \frac{1}{\lambda}(\frac{a^2}{b^2} - \frac{\bar b^2}{\bar a^2}) = 0$ and its conjugate are equivalent to (I)&(II). Differentiating these we get $\frac{b}{a}dU - \epsilon\frac{\bar a}{b}d\bar V = \frac{\bar b}{\bar a}d\bar U - \epsilon\frac{a}{b}dV = 0$; now (III) imposes $\bar a^2 = \epsilon b^2$; finally we get $a_2 = -\epsilon\bar a_1,\ a_1 \in \mathbb{C}\setminus\{0\}$ (note that $a_1 \in \mathbb{R}\setminus\{0\}$ is allowed), $(u,v) = (\sqrt{\epsilon_1\sqrt{\epsilon}}s, \sqrt{-\epsilon_1\sqrt{\epsilon}}t),\ s,t \in \mathbb{R},\ \epsilon_1 := \pm 1$.

For QC (3.4) with $a := (\sqrt{a_1^{-1}})^{-1},\ b := (\sqrt{a_2^{-1}})^{-1},\ V := (u-v)^{-2},\ U := (u+v)^2V$ (I)&(II) become $b(U(1-\frac{a^2}{b^2})-1+\frac{4V}{a^2}) = -\epsilon\bar b(\bar U(1-\frac{\bar a^2}{b^2})-1+\frac{4\bar V}{\bar a^2}),\ b(U(1-\frac{a^2}{b^2})((1-\frac{a^2}{b^2})(U-1)+(3-\frac{a^2}{b^2})\frac{4V}{a^2})+\frac{b^2}{a^2}(\frac{4V}{a^2})^2) = -\epsilon\bar b(\bar U(1-\frac{\bar a^2}{b^2})((1-\frac{\bar a^2}{b^2})(\bar U-1)+(3-\frac{\bar a^2}{b^2})\frac{4\bar V}{\bar a^2})+\frac{\bar b^2}{\bar a^2}(\frac{4\bar V}{\bar a^2})^2)$; with $\mathcal{V} := \frac{4V}{a^2},\ \mathcal{E} := 1-\frac{a^2}{b^2},\ \mathcal{U} := U\mathcal{E}-1+\mathcal{V}$ these imply $(1+\frac{\epsilon b}{b})(\frac{\epsilon b}{b}\mathcal{U}^2+(\mathcal{E}(\mathcal{V}-1)-\overline{\mathcal{E}}(\overline{\mathcal{V}}-1))\mathcal{U}+1-\mathcal{E}((1-\mathcal{V})^2+\mathcal{V}^2))-(\mathcal{E}(\mathcal{V}-1)-\overline{\mathcal{E}}(\overline{\mathcal{V}}-1))\mathcal{U}+\mathcal{E}((1-\mathcal{V})^2+\mathcal{V}^2)-\overline{\mathcal{E}}((1-\overline{\mathcal{V}})^2+\overline{\mathcal{V}}^2) = 0$; further $|dx_0^0|^2 = \frac{b^2}{4U}dU^2 - \frac{a^2}{2V}dU\odot dV + \frac{4V-a^2(1-U)}{4V^2}dV^2$.

If $a^2 = b^2$, then (I)&(II) imply $\bar a = -\epsilon a,\ \bar V = V$; now (III) becomes $\frac{V}{U}(d\frac{U}{V})^2 - \frac{\bar V}{\bar U}(d\frac{\bar U}{\bar V})^2 = 0$, or $(du+dv)^2 - (d\bar u + d\bar v)^2 = 0$ and finally $a_1 = a_2 = -\epsilon|a_1|,\ (u,v) = (\sqrt{\epsilon_1}(s+ic), \sqrt{\epsilon_1}(t+ic)),\ s,t \in \mathbb{R}$ or $v = \epsilon_1\bar u - i\sqrt{\epsilon_1}c,\ u \in \mathbb{C},\ c \in \mathbb{R}$ constant, $\epsilon_1 := \pm 1$.

If $a^2 \neq b^2$, then we have $|dx_0^0|^2 = \frac{a^2(d\mathcal{U}-d\mathcal{V})^2}{4\mathcal{E}(1-\mathcal{E})(\mathcal{U}+1-\mathcal{V})} - \frac{a^2}{2\mathcal{E}\mathcal{V}}d(\mathcal{U}-\mathcal{V})\odot d\mathcal{V} + \frac{a^2}{4\mathcal{V}^2}(\mathcal{V}-1+\frac{\mathcal{U}+1-\mathcal{V}}{\mathcal{E}})d\mathcal{V}^2$.

3.3. **Two algebraic consequences of the tangency configuration.**
Choose a surface x_0^0 in the quadric x_0 (that is $(u_0,v_0) = (u_0(s,t), v_0(s,t))$, where s,t are independent variables (real or complex, but not necessarily both of the same type) such that $u_0,\ v_0$ are still independent; thus $T_{x_0^0}x_0^0 = T_{x_0^0}x_0$). $T_{x_0^0}x_0$ cuts x_z along a conic; for every point x_z^1 on this conic we have the two facets at x_z^1 spanned by $V_0^1 := x_z^1 - x_0^0$ and one of the rulings of x_z at x_z^1. The two distributions of facets $\mathcal{D}^1,\ \mathcal{D}'^1$ are integrable, their leaves being the ruling families on x_z (we shall see in § 5.1 that the complete integrability depends only on the linear element of x_0^0). Choose $m_0^1 := \mathcal{B}_1x_{zu_1}^1 \times V_0^1$ a normal field of the distribution $\mathcal{D}^1$ with leaves $x_{zu_1}^1$ (and similarly $m_0'^1 := \mathcal{B}_1x_{zv_1}^1 \times V_0^1$ by considering the other ruling family on x_z^1). Note that $m_0^1 = -i\det(\sqrt{R_z}(\sqrt{A})^{-1})(\sqrt{R_z})^{-1}\sqrt{A}Y(v_1) - x_0^0 \times \sqrt{R_z}(\sqrt{A})^{-1}Y(v_1)$ for QC, $= 2x_{zu_1}^1 \times (x_z^1(0,v_1) - x_0^0)$ for (I)QWC ($m_0'^1 = -i\det(\sqrt{R_z}(\sqrt{A})^{-1})(\sqrt{R_z})^{-1}\sqrt{A}Y(u_1) + x_0^0 \times \sqrt{R_z}(\sqrt{A})^{-1}Y(u_1)$ for QC, $= 2x_{zv_1}^1 \times (x_z^1(u_1,0) - x_0^1)$ for (I)QWC) depends quadratically on v_1 (u_1), fact which will make the differential system of the B transformation a Ricatti equation.

The next two algebraic results (essentially due to Bianchi; see (122)) play a fundamental rôle in the theory of deformations of quadrics:

If $x_z^1 \in T_{x_0^0}x_0$, then:

I The change in the linear element from x_z^1 to x_0^1 is four times the product of the orthogonal projections of the differentials of the rulings of x_z^1 on the normal of x_0 at x_0^0.

II The facets at x_z^1 spanned by V_0^1 and one of the rulings of x_z^1 reflect in $T_{x_0^0}x_0$; therefore the distributions $\mathcal{D}^1,\ \mathcal{D}'^1$ reflect in Tx_0^0.

This can be put as: if $(V_0^1)^T\hat N_0^0 = 0$, then:

$$(3.12)\ 4(x_{zu_1}^1)^TN_0^0(N_0^0)^Tx_{zv_1}^1du_1dv_1 = |dx_z^1|^2 - |dx_0^1|^2,\ (x_{zv_1}^1)^T(I_3 - 2N_0^0(N_0^0)^T)m_0^1 = 0.$$

Since $|dx_z|^2 - |dx_0|^2 = -zdx_0^TAdx_0 = -z|dX|^2$ for QC, $= -z|I_{1,2}dZ|^2$ for (I)QWC, these boil down to: $\mathcal{B}_1(x_{zu_1}^1)^T\hat N_0^0(\hat N_0^0)^Tx_{zv_1}^1 = -z|\hat N_0^0|^2,\ (x_{zv_1}^1 \times x_{zu_1}^1)^T\hat N_0^0(\hat N_0^0)^TV_0^1 = (\hat N_0^0)^T(x_{zu_1}^1(x_{zv_1}^1)^T + x_{zv_1}^T(x_{zu_1}^1)^T)(V_0^1 \times \hat N_0^0)$.

The relation

$$(3.13) \qquad (x_{zv_1}^1(x_{zu_1}^1)^T + x_{zu_1}^1(x_{zv_1}^1)^T)\hat N_0^0 = x_{zu_1v_1}^1(V_0^1)^T\hat N_0^0 - \frac{2}{\mathcal{B}_1}(V_0^1 + z\hat N_0^0)$$

follows immediately from (3.11).

Multiplying (3.13) with $(\hat{N}_0^0)^T$ on the left we get the fact that the first relation of (3.12) for (I)QWC is equivalent to the TC $(V_0^1)^T\hat{N}_0^0=0$; for QC it is equivalent to the TC $(V_0^1)^T\hat{N}_0^0=0$ or to $X_0^T\sqrt{R_z}X_1=-1$, which can be brought to the TC by $u_0\leftrightarrow v_0$. Multiplying (3.13) with $(V_0^1\times\hat{N}_0^0)^T$ on the left we get the fact that the second relation of (3.12) is equivalent to the TC $(V_0^1)^T\hat{N}_0^0=0$ or to $0=(\hat{N}_0^0)^T(V_{0u_1}^1\times V_0^1)_{v_1}=(\hat{N}_0^0)^T(\frac{m_0^1}{\mathcal{B}_1})_{v_1}$. If $(V_0^1)^T\hat{N}_0^0\neq 0$ and the facets reflect in $T_{x_0^0}x_0$, then from $(V_0^1)^Tm_0'^1=0$ we get $(\hat{N}_0^0)^Tm_0^1=0$; thus $m_0^1,\ m_0'^1$ are multiples of $\hat{N}_z^1$ and $(\hat{N}_0^0)^T\hat{N}_z^1=0$; essentially by Chasles's result we get $(V_0^1)^T\hat{N}_0^0=0$, a contradiction. Therefore the two algebraic consequences of the TC are actually equivalent to the TC.

Note:

$$\begin{aligned}(V_0^1)^T\hat{N}_0^0&=X_0^T\sqrt{R_z}X_1-1\text{ for QC},\\ &=Z_0^TI_{1,2}\sqrt{R_z'}Z_1-f^T(x_z^0+x_z^1-C(z))=Z_0^TI_{1,2}\sqrt{R_z'}Z_1-\\ -(Z_0+Z_1)^T(u(z)f_1+v(z)\bar{f}_1+e_3)-u(z)v(z)-\frac{|f|^2}{2}z&\text{ for (I)QWC.}\end{aligned}\tag{3.14}$$

If we let $u_1=v_1=0$ in (3.14) for (I)QWC we get $(\hat{N}_0^0)^TC(z)+f^Tx_z^0=(\hat{N}_0^0)^Tx_0^0(=u_0v_0)$.

Consider now the TC $(V_0^1)^T\hat{N}_0^0=0$; one can choose v_1 arbitrarily and define

$$u_1:=v_1+\frac{Y(v_1)^T\sqrt{R_z}X_0}{1-\frac{1}{2}Y'(v_1)^T\sqrt{R_z}X_0}\text{ for QC},\ u_1:=\frac{v_1(\hat{N}_0^0)^Tx_{zv}(0,0)-f^Tx_z^0}{v_1-(\hat{N}_0^0)^Tx_{zu}(0,0)}\text{ for (I)QWC;}\tag{3.15}$$

it is a ratio of two functions separately linear in u_0,v_0,v_1 and thus a homography is established between u_0,v_0,u_1,v_1.

If we let $u_1=v_1=0$ in (3.13) for (I)QWC and multiply with $(\hat{N}_0^0)^T$ on the left we get:

$$\frac{z}{2}|\hat{N}_0^0|^2=f^Tx_z^0-x_{zu}(0,0)^T\hat{N}_0^0(\hat{N}_0^0)^Tx_{zv}(0,0);\tag{3.16}$$

note also for (I)QWC

$$(\hat{N}_0^0)^T(u_1x_{zu}(0,0)+v_1x_{zv}(0,0))=(\hat{N}_0^0)^TV_0^1+f^Tx_z^0+u_1v_1,\tag{3.17}$$

which is equivalent to (3.15) in the TC case. The algebraic relation

$$(N_0^0)^T(2zm_0^1+m_0^1\times m_{0v_1}^1)=0\tag{3.18}$$

will appear as the complete integrability condition of the Ricatti equation subjacent to the B transformation. Using (3.12) this becomes: $0=\frac{z(m_0^1)^Tx_{zv_1}^1}{(N_0^0)^Tx_{zv_1}^1}-\mathcal{B}_1(x_{zu_1}^1)^TN_0^0(V_0^1)^Tm_{0v_1}^1=$
$=\frac{z(V_0^1)^T(\mathcal{B}_1x_{zv_1}^1\times x_{zu_1}^1+(\mathcal{B}_1x_{zu_1}^1\times V_0^1)_{v_1})}{(N_0^0)^Tx_{zv_1}^1}$, which is straightforward. Replacing (m_0^1,v_1) with $(m_0'^1,u_1)$ we get a similar relation.

3.4. **The Ivory affinity provides a rigid motion.**

From the fact that the Ivory affinity preserves lengths and angles of segments between (and rulings on) confocal quadrics, we shall find that if $x_0^0(u_0,v_0),\ x_0^1(u_1,v_1)$ are on the quadric x_0 and choosing rulings $w_0^j,\ w_z^j$ at each of the points $x_0^j,\ x_z^j$, then there is an rigid motion (R_0^1,t_0^1) of $\mathbb{C}^3$ which takes x_0^0 to x_z^0, x_z^1 to x_0^1, w_0^0 to w_z^0 and w_z^1 to w_0^1:

$$(R_0^1,t_0^1)x_0^0=x_z^0,\ (R_0^1,t_0^1)x_z^1=x_0^1,\ R_0^1w_0^0=w_z^0,\ R_0^1w_z^1=w_0^1.\tag{3.19}$$

The first two conditions can be replaced by $R_0^1V_0^1=-V_1^0,\ V_0^1:=x_z^1-x_0^0,\ V_1^0:=x_z^0-x_0^1$.



Such an R_0^1 can be found because: $[(x_z^1-x_0^0)\ \ w_0^0\ \ w_z^1]^T[(x_z^1-x_0^0)\ \ w_0^0\ \ w_z^1] =$
$= \begin{bmatrix} |x_z^1-x_0^0|^2 & (w_0^0)^T(x_z^1-x_0^0) & (w_z^1)^T(x_z^1-x_0^0) \\ (w_0^0)^T(x_z^1-x_0^0) & |w_0^0|^2 & (w_0^0)^Tw_z^1 \\ (w_z^1)^T(x_z^1-x_0^0) & (w_0^0)^Tw_z^1 & |w_z^1|^2 \end{bmatrix} = [(x_0^1-x_z^0)\ \ w_z^0\ \ w_0^1]^T[(x_0^1-x_z^0)\ \ w_z^0\ \ w_0^1].$
To fix the ideas we shall henceforth make a choice: $w_0^j := x_{0u_j}^j,\ j=0,1$; the rotations (in the case of the TC $(V_0^1)^T\hat N_0^0=0$) when changing the ruling family on x_0^1, on x_0^0 or on both are obtained, according to the second relation of (3.12), by first reflecting in $T_{x_0^0}x_0$ or further in $T_{x_0^1}x_0$ or both; conversely, the second relation of (3.12) is a consequence of the existence of the RMPIA. Because of this $x_{0v_1}^1 = R_0^1(I_3-2N_0^0(N_0^0)^T)x_{zv_1}^1$; multiplying this on the left with $(x_{0u_1}^1)^T$ we get the first relation of (3.12). Note that (R_0^1,t_0^1) depends only on (v_0,v_1): with $F := [\mathcal{B}_0x_{0u_0}^0 \quad \mathcal{B}_1x_{zu_1}^1 \quad (x_0^0(0,v_0)-x_z^1(0,v_1))]$ we have:

$$R_0^1 = ((0\leftrightarrow z)\circ F)F^{-1};\ t_0^1 = x_z^0(0,v_0) - R_0^1x_0^0(0,v_0). \tag{3.20}$$

Let $\Delta^- = \Delta^-(z,v_0,v_1) := -\mathcal{B}_0(m_0^1)^Tx_{0u_0}^0,\ \Delta^+ = \Delta^+(z,u_0,v_1) := \mathcal{B}_0(m_0^1)^Tx_{0v_0}^0$; similarly we can consider $\Delta'^{\pm}$ by changing the ruling family on x_z^1 (replace (m_0^1,v_1) with $(m_0'^1,u_1)$).

We have $\frac{-\Delta^-}{4\mathcal{A}} = \frac14 Y(v_0)^T(\det\sqrt{R_z}(\sqrt{R_z})^{-1}+\sqrt{R_z})Y(v_1)$ for QC, $= -i((f_1+v_0e_3)\times(\sqrt{R_z'}f_1+v_{z1}e_3))^T(v_{z1}\sqrt{R_z'}\bar f_1 - v_0\bar f_1 + z(1-\frac{|f|^2}{2})(L^TL)^{-1}e_3) = (1+\det\sqrt{R_z'})v_0v_1 - \bar f_1^T\sqrt{R_z'}f_1((v_0+v_1+v(z))^2 - 2v_0v_1) - zf_1^T(L^TL)^{-1}e_3(v_0+v_1+v(z)) + z\frac{|f|^2}{2}f_1^T\sqrt{R_z'}f_1 = v_0v_1(1+\det\sqrt{R_z'}) - (v_0^2+v_1^2)\bar f_1^T\sqrt{R_z'}f_1 + \frac z2 f_1^T\sqrt{R_z'}f_1$ for QWC, $= -\frac{1+\det\sqrt{R_z'}}{2}(v_0-v_1)^2 + v(z)(1-\det\sqrt{R_z'})(v_0+v_1) + \frac{v(z)^2}{2}(3-\det\sqrt{R_z'})$ for IQWC (3.8), $= -(v_0-v_1)^2 - 2v(z)(v_0+v_1) - v(z)^2$ for IQWC (3.9). We have $\det F = \Delta^- = (v_0\leftrightarrow v_1)\circ\Delta^- = \det((v_0\leftrightarrow v_1)\circ F) = \det((0\leftrightarrow z)\circ F)$, so $\det R_0^1 = 1$.

Differentiating $(V_0^1)^T\hat N_0^0 = (V_1^0)^T\hat N_0^1$ wrt u_0 we get:

$$(V_0^1)^T\hat N_{0u_0}^0 = (x_{zu_0}^0)^T\hat N_0^1. \tag{3.21}$$

Consider now the TC $(V_0^1)^T\hat N_0^0 = 0$; because $R_0^1\frac{m_0^1}{|m_0^1|} = \pm\frac{\hat N_0^1}{|\hat N_0^1|}$ we have $R_0^1V_0^1 = \mathcal{A}\hat N_0^1\times R_0^1\hat N_0^0$ (multiply it on the left with $(R_0^1(x_{0u_0}^0\times x_0^0))^T$, use (3.21), $x_{0u}\times x_0 = \mathcal{A}\hat N_{0u},\ \hat N_0^Tx_0=1$ for QC and $x_0^T\hat N_0 = uv,\ x_{0u}\times x_0 = -v\mathcal{A}(\hat N_0 - u\hat N_{0u})$ for(I)QWC). Thus:

$$R_0^1V_0^1 = \mathcal{A}\hat N_0^1\times R_0^1\hat N_0^0,\ R_0^1m_0^1 = \mathcal{A}\mathcal{B}_1\hat N_0^1(\hat N_0^0)^Tx_{zu_1}^1. \tag{3.22}$$

If we change the ruling family on x_0^0 or on x_z^1 or on both, then the sign of the rhs changes in the second or the first or both relations of (3.22). Using (3.12) and (3.22) we thus have:

$$\begin{aligned} \Delta^- &= -\mathcal{A}\mathcal{B}_0\mathcal{B}_1(x_{zu_0}^0)^T\hat N_0^1(\hat N_0^0)^Tx_{zu_1}^1, \\ \Delta^+ &= -\mathcal{A}\mathcal{B}_0\mathcal{B}_1(x_{zv_0}^0)^T\hat N_0^1(\hat N_0^0)^Tx_{zu_1}^1, \\ \Delta^+\Delta^- &= -z\mathcal{A}^2\mathcal{B}_0\mathcal{B}_1^2|\hat N_0^1|^2((x_{zu_1}^1)^T\hat N_0^0)^2. \end{aligned} \tag{3.23}$$

By changing the ruling family on x_z^1 we get similar formulae for $\Delta'^{\pm}$ (the sign of the rhs does not change), $\Delta'^+\Delta'^-,\ \Delta^-\Delta'^-,\ \Delta^+\Delta'^+$. With $\mathcal{N} := \frac{1}{\mathcal{A}\mathcal{B}|\hat N_0|^2}$ the first relation of (3.12) and (3.23) imply

$$\Delta^-\Delta'^+ = \frac{z^2}{\mathcal{N}_0\mathcal{N}_1} = \Delta^+\Delta'^-; \tag{3.24}$$

conversely $\Delta^-\Delta'^+ - \Delta^+\Delta'^- = 0$ is equivalent to $0 = \frac{(m_0^1)^T x_{0v_0}^0 (m_0'^1)^T x_{0u_0}^0 - (m_0^1)^T x_{0u_0}^0 (m_0'^1)^T x_{0v_0}^0}{(N_0^0)^T (x_{0u_0}^0 \times x_{0v_0}^0)} = (m_0'^1 \times m_0^1)^T N_0^0 = -\mathcal{B}_1^2 (N_z^1)^T (x_{zu_1}^1 \times x_{zv_1}^1)(V_0^1)^T N_z^1 (V_0^1)^T N_0^0$, so it is equivalent either to the TC or to $(V_0^1)^T N_z^1 = 0$.

Suppose $\{m_j\}_{j=1,2,3}$ are functions of the independent variables $\{y_k\}_{k=1,...,4}$ (that is $dy_1 \wedge ... \wedge dy_4 \neq 0,\ dm_j \wedge dy_1 \wedge ... \wedge dy_4 = 0$). If we can restrict y_4 to $dy_1 \wedge dy_2 \wedge dy_3 \neq 0,\ dy_4 = \sum_{j=1}^3 m_j dy_j$, then imposing the compatibility condition $d\wedge$ to the last relation we get

$$\partial_{y_4} \log \frac{m_{j+1}}{m_j} + \frac{1}{m_j}\partial_{y_j} \log m_{j+1} - \frac{1}{m_{j+1}}\partial_{y_{j+1}} \log m_j = 0,\ j =\circlearrowleft_1^3 . \tag{3.25}$$

Applying d to the TC $(V_0^1)^T \hat{N}_0^0 = 0$, using the first relation of (3.12), (3.21) and (3.23) we get

$$du_1 = -\frac{\mathcal{N}_0}{z}(\Delta'^- du_0 + \Delta'^+ dv_0) - \frac{\Delta'^+}{\Delta^+} dv_1. \tag{3.26}$$

Applying (3.25) to $(y_1, y_2, y_3, y_4, m_1, m_2, m_3) := (u_0, v_0, v_1, u_1, -\frac{\mathcal{N}_0\Delta'^-}{z}, -\frac{\mathcal{N}_0\Delta'^+}{z}, -\frac{\Delta'^+}{\Delta^+}),\ j = 1, 3$ and to $(y_1, y_2, y_3, y_4, m_1, m_2, m_3) := (v_0, u_0, v_1, u_1, -\frac{z}{\mathcal{N}_1\Delta^-}, -\frac{\mathcal{N}_0\Delta'^-}{z}, -\frac{\Delta'^-}{\Delta^-}),\ j = 1$ we get the algebraic relations

$$\begin{aligned} \frac{\mathcal{N}_0}{z}\partial_{u_1} \log \frac{\Delta'^+}{\Delta'^-} = \frac{1}{\Delta'^-}\partial_{u_0} \log(\mathcal{N}_0\Delta'^+) - \frac{1}{\Delta'^+}\partial_{v_0} \log(\mathcal{N}_0\Delta'^-) = \frac{1}{\Delta'^-}\partial_{u_0} \log \frac{\Delta'^+}{\Delta^+} = \\ \frac{\mathcal{N}_0}{z}\partial_{u_1} \log(\mathcal{N}_1\Delta'^+) - \frac{1}{\Delta'^+}\partial_{v_0} \log(\mathcal{N}_0\Delta'^-). \end{aligned} \tag{3.27}$$

Note that while explicit derivatives $\partial_{\cdot}$ obey usual calculus rules, one must avoid the use of consequences of the TC inside the $\partial_{\cdot}$ sign; for example $\partial_{u_0} \log \frac{\Delta^-}{\Delta'^-} = 0 \neq \partial_{u_0} \log \frac{\Delta^+}{\Delta'^+}$. Since the first relation of (3.27) does not depend on v_1, it is valid for independent u_0, v_0, u_1.

Note $((u_0, v_0) \leftrightarrow (u_1, v_1)) \circ (\Delta^-, \Delta^+, \Delta'^-, \Delta'^+) = (\Delta^-, \Delta'^-, \Delta^+, \Delta'^+)$; using (3.24) the relation (3.26) (and thus (3.27)) has the symmetries $(u_0, v_0, \Delta^+, \Delta'^-, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \Delta'^-, \Delta^+, \mathcal{N}_1)$, $(u_0, \Delta^+, \Delta'^+) \leftrightarrow (v_0, \Delta^-, \Delta'^-),\ (u_1, \Delta'^-, \Delta'^+) \leftrightarrow (v_1, \Delta^-, \Delta^+)$. Under these symmetries and compositions thereof, there are no other algebraic identities besides (3.27) and its immediate consequences (like $\frac{1}{\Delta'^-}\partial_{u_0} \log \frac{\Delta'^+}{\Delta^+} = -\frac{1}{\Delta'^+}\partial_{v_0} \log \frac{\Delta'^-}{\Delta^-}$) which appear from (3.26) and its equivalent versions.

The TC has also other symmetries: for QC $u_0 \leftrightarrow v_0$ ($u_1 \leftrightarrow v_1$) and $\sqrt{R_z} \leftrightarrow -\sqrt{R_z}$ (which reflect $x_0^0(x_0^1)$ in the origin and restore the TC: $\sqrt{R_z}X(u, v) = -\sqrt{R_z}X(v, u)$); for QWC (3.6) and IQWC (3.8) $u_0 \leftrightarrow v_0,\ (u_1 \leftrightarrow v_1)$ and $\sqrt{R_z'} \leftrightarrow r_{e_2}\sqrt{R_z'} = \sqrt{R_z'}r_{e_2},\ r_{e_2} := f_1 f_1^T + \bar{f}_1 \bar{f}_1^T + e_3 e_3^T = I_3 - 2e_2 e_2^T$ (which reflect $x_0^0\ (x_0^1)$ in the plane $e_2^T x = 0$ for QWC (3.6) or in the plane $e_3^T x = 0$ for IQWC (3.8) and restore the TC: $\sqrt{R_z'}I_{1,2}Z(u, v) = r_{e_2}\sqrt{R_z'}I_{1,2}Z(v, u)$).

For QC $\Delta^+ = ((u_0, \sqrt{R_z}) \leftrightarrow (v_0, -\sqrt{R_z})) \circ \Delta^-,\ \Delta'^+ = ((u_0, u_1) \leftrightarrow (v_0, v_1)) \circ \Delta^-$; for QWC (3.6) and IQWC (3.8) $\Delta^+ = ((u_0, \sqrt{R_z'}) \leftrightarrow (v_0, r_{e_2}\sqrt{R_z'})) \circ \Delta^-,\ \Delta'^+ = ((u_0, u_1) \leftrightarrow (v_0, v_1)) \circ \Delta^-$. For QWC (3.7) and IQWC (3.9) the change $\sqrt{R_z'} \leftrightarrow \sqrt{R_z'}r_{e_2}$ for changing the ruling family on x_0^1 or $f_1 \leftrightarrow r_{e_2}f_1 = \bar{f}_1$ for changing the ruling family on x_0^0 still has some effect similar to QWC (3.6) and IQWC (3.8), but since $\sqrt{R_z'}r_{e_2} \neq r_{e_2}\sqrt{R_z'}$ new formulae appear: $\frac{-\Delta^+}{4\mathcal{A}} = v_1(u_0(1 - \det\sqrt{R_z'}) - v_1 \bar{f}_1^T \sqrt{R_z'} \bar{f}_1) + \frac{z}{2}\bar{f}_1^T \sqrt{R_z'} f_1,\ \frac{-\Delta'^+}{4\mathcal{A}} = u_0 u_1(1 + \det\sqrt{R_z'}) - (u_0^2 + u_1^2)\bar{f}_1^T \sqrt{R_z'} f_1 + \frac{z}{2}\bar{f}_1^T \sqrt{R_z'} \bar{f}_1$ for QWC (3.7), $\frac{-\Delta^+}{4\mathcal{A}} = \frac{z}{2}u_0(u_0 - z) - zv_1 + \frac{z^3}{8},\ \frac{-\Delta'^+}{4\mathcal{A}} = \frac{z^2}{4} - (u_0 - u_1)^2$ for IQWC (3.9).

3.5. **Second iteration of the tangency configuration.**

Consider the SITC: $x_{z_1}^0,\ x_{z_2}^3 \in T_{x_0^1}x_0$. In the pencil of planes containing $x_{z_2}^0,\ x_{z_1}^3$ choose a certain one tangent to x_0 at x_0^2 (in general there are two choices; for the other choice we must consider the other ruling families on $x_0^2,\ x_0^3$). To have a consistent notation we shall order $0 < z_1 < z_2$ and the position of j, k in $V_k^j,\ (R_k^j, t_k^j)$ is increasing wrt this order: for example $V_0^3 := x_{z_2}^3 - x_{z_1}^0 \in T_{x_0^1}x_0$ since there is no other incidence relationship involving the upper indices $0, 3$ and lower indices $0, z_1$

or $0, z_2$; $R_3^0(x^0_{z_2u_0}, x^3_{z_1u_3}, V_3^0) = (x^0_{z_1u_0}, x^3_{z_2u_3}, -V_0^3)$, $(R_3^0)^{-1} = R_0^3$. Note:

$$V_0^3 = V_1^3 - V_1^0,\ V_3^0 = V_2^0 - V_2^3,\ V_1^2 = V_0^2 - V_0^1,\ V_2^1 = V_3^1 - V_3^2,\ (V_0^1)^T\hat{N}_0^0 = (V_0^2)^T\hat{N}_0^0 =$$
$$(V_1^0)^T\hat{N}_0^1 = (V_1^3)^T\hat{N}_0^1 = (V_2^0)^T\hat{N}_0^2 = (V_2^3)^T\hat{N}_0^2 = (V_3^1)^T\hat{N}_0^3 = (V_3^2)^T\hat{N}_0^3 = 0. \tag{3.28}$$

In particular the SITC has the group of symmetries of the square generated by the flip ($0 \leftrightarrow 1, 2 \leftrightarrow 3$) and the rotation ($0 \to 1 \to 3 \to 2 \to 0,\ z_1 \leftrightarrow z_2$) (consider the square with vertices $0, 1, 2, 3$ and sides labelled $z_1,\ z_2$ such that both the sums of opposite vertices and the labels of opposite sides are equal; $0, 1, 2, 3$ do not yet respectively correspond, up to some rigid motions, to $(R_0^1, t_0^1)x_0^0, x_0^1, (R_3^0, t_3^0)x_0^2, (R_3^1, t_3^1)x_0^3$, but the symmetries can be seen at the level of (3.28)).

The next result (proved by Bianchi in (122,§ 26)) plays a fundamental rôle in the proof of the BPT:

If $x^0_{z_1},\ x^3_{z_2} \in T_{x_0^1}x_0$ and for a certain choice of x_0^2 such that $x^0_{z_2},\ x^3_{z_1} \in T_{x_0^2}x_0$, then the rulings $x^1_{0u_1}, R_3^0x^2_{0u_2}$ cut the line $[x^0_{z_1}\ x^3_{z_2}]$ with cross ratio $\frac{z_1}{z_2}$.



From (3.15) we get: $\frac{Y(v_1)^T\sqrt{R_{z_1}}X_0}{1-\frac{1}{2}Y'(v_1)^T\sqrt{R_{z_1}}X_0} = u_1 - v_1 = \frac{Y(v_1)^T\sqrt{R_{z_2}}X_3}{1-\frac{1}{2}Y'(v_1)^T\sqrt{R_{z_2}}X_3}$ for QC, $\frac{v_1x_{z_1v}(0,0)^T\hat{N}_0^0 - f^Tx^0_{z_1}}{v_1 - x_{z_1u}(0,0)^T\hat{N}_0^0} = u_1 = \frac{v_1x_{z_2v}(0,0)^T\hat{N}_0^3 - f^Tx^3_{z_2}}{v_1 - x_{z_2u}(0,0)^T\hat{N}_0^3}$ for (I)QWC and other similar equations, obtained by ($0 \leftrightarrow 3, 1 \leftrightarrow 2$). These become:

$$\frac{1}{2}Y(v_1)^T(\sqrt{R_{z_2}}X_3 - \sqrt{R_{z_1}}X_0 - i(\sqrt{R_{z_2}}X_3) \times (\sqrt{R_{z_1}}X_0)) = 0 \text{ for QC},$$
$$v_1^2(x_{z_1v}(0,0)^T\hat{N}_0^0 - x_{z_2v}(0,0)^T\hat{N}_0^3) + v_1(f^TV_0^3 - x_{z_1v}(0,0)^T\hat{N}_0^0x_{z_2u}(0,0)^T\hat{N}_0^3 +$$
$$x_{z_1u}(0,0)^T\hat{N}_0^0x_{z_2v}(0,0)^T\hat{N}_0^3) - x_{z_1u}(0,0)^T\hat{N}_0^0f^Tx^3_{z_2} + x_{z_2u}(0,0)^T\hat{N}_0^3f^Tx^0_{z_1} = 0$$
$$\text{for (I)QWC.} \tag{3.29}$$

When $v_j,\ j = 0, 1, 3$ are fixed $u_0,\ u_3$ vary preserving (3.29), so a homography (invariant under ($0 \leftrightarrow 3, 1 \leftrightarrow 2$)) is established between the rulings $x^0_{z_1u_0}, x^3_{z_2u_3}$. The quadric Q^{301} (Q^{032}) generated by $x^3_{z_2u_3}, x^0_{z_1u_0}, x^1_{0u_1}$ ($x^0_{z_2u_0}, x^3_{z_1u_3}, x^2_{0u_2}$) cuts x_0 along $x^1_{0u_1}$ ($x^2_{0u_2}$) (and possibly other sets we exclude from our discussion). The points on $x^0_{z_1u_0}, x^3_{z_2u_3}$ ($x^0_{z_2u_0}, x^3_{z_1u_3}$) corresponding under the homography lie on the other ruling family of Q^{301} (Q^{032}), so $(R_0^3, t_0^3)Q^{301} = Q^{032}$ (if a line l_1 intersects two non-co-planar lines l_2, l_3, then there is a 2-dimensional family of quadrics having l_2, l_3 in a ruling family and l_1 in the other ruling family; each quadric is uniquely determined by the homography it establishes between l_2, l_3 and it corresponds to the choices of an arbitrary line l_4 at a fixed point on l_1; l_4 will belong to the first ruling family), which is equivalent to the rulings $x^0_{z_1u_0},\ x^3_{z_2u_3}$ cutting the line $[x_0^1\ (R_3^0, t_3^0)x_0^2]$ (again with cross-ratio $\frac{z_1}{z_2}$, because four rulings of a ruling family on a quadric cut any ruling of the other ruling family with same cross ratio). This is equivalent to $R_1^0R_3^0V_2^0 \in T_{x_0^0}x_0,\ R_1^3R_3^0V_2^3 \in T_{x_0^3}x_0$, from which follow the co-cycle relations

$$(R_0^1, t_0^1)(R_2^0, t_2^0) = (R_3^0, t_3^0) = (R_3^1, t_3^1)(R_2^3, t_2^3),\ (R_1^0, t_1^0)(R_3^1, t_3^1) = (R_2^1, t_2^1) = (R_2^0, t_2^0)(R_3^2, t_3^2). \tag{3.30}$$

To prove this let $(R_0, t_0) := (R_1^0, t_1^0)(R_3^0, t_3^0)(R_0^2, t_0^2),\ (R_1, t_1) := (R_0^1, t_0^1)(R_2^1, t_2^1)(R_1^3, t_1^3),\ (R_2, t_2) := (R_3^2, t_3^2)(R_1^2, t_1^2)(R_2^0, t_2^0),\ (R_3, t_3) := (R_2^3, t_2^3)(R_0^3, t_0^3)(R_3^1, t_3^1)$; note: $(R_j, t_j)(x_0^j, x^j_{0u_j}) = (x_0^j, x^j_{0u_j})$, $j = 0, ..., 3$, so R_j are rotations around $x^j_{0u_j},\ j = 0, ..., 3$. But $R_0V_0^2 \in T_{x_0^0}x_0,\ R_3^{-1}V_3^2 \in T_{x_0^3}x_0$; thus $R_0(T_{x_0^0}x_0) = T_{x_0^0}x_0,\ R_3(T_{x_0^3}x_0) = T_{x_0^3}x_0$; since $\det R_0 = \det R_3 = 1$ we get $R_0 = R_3 = I$, so $t_0 = t_3 = 0$. From the first four relations of (3.28) we get $-R_0^1(I - R_0)V_0^2 + R_3^1(I - R_3^{-1})V_3^2 + (I -$

$R_1^{-1})R_0^1V_1^2=0$, so $R_1=I$ and $R_2^{-1}=R_0^2R_1^0R_1R_0^1R_0R_2^0R_3^2R_3R_2^3=I$, so $t_1=t_2=0$. Another way to prove $(R_1,t_1)=(R_2,t_2)=(I_3,0)$ once we know $(R_0,t_0)=(R_3,t_3)=(I_3,0)$ is to apply to the SITC the symmetry $(0\to 1\to 3\to 2\to 0,\ z_1\leftrightarrow z_2)$.

Applying the rigid motions $(R_0^3,t_0^3),\ (R_1^0,t_1^0),\ (R_1^3,t_1^3)$ to the configuration
$((R_0^1,t_0^1)(x_0^0,x_{0u_0}^0),(x_0^1,x_{0u_1}^1),(R_3^0,t_3^0)(x_0^2,x_{0u_2}^2),(R_3^1,t_3^1)(x_0^3,x_{0u_3}^3))$ we get the configurations
$((R_0^2,t_0^2)(x_0^0,x_{0u_0}^0),(R_0^3,t_0^3)(x_0^1,x_{0u_1}^1),(x_0^2,x_{0u_2}^2),(R_3^2,t_3^2)(x_0^3,x_{0u_3}^3))$,
$((x_0^0,x_{0u_0}^0),(R_1^0,t_1^0)(x_0^1,x_{0u_1}^1),(R_2^0,t_2^0)(x_0^2,x_{0u_2}^2),(R_2^1,t_2^1)(x_0^3,x_{0u_3}^3))$,
$((R_1^2,t_1^2)(x_0^0,x_{0u_0}^0),(R_1^3,t_1^3)(x_0^1,x_{0u_1}^1),(R_2^3,t_2^3)(x_0^2,x_{0u_2}^2),(x_0^3,x_{0u_3}^3))$.

These rigid motions realize the symmetries $(0\leftrightarrow 3,1\leftrightarrow 2),(0\leftrightarrow 1,2\leftrightarrow 3),(0\leftrightarrow 2,1\leftrightarrow 3)$; the symmetry $(0\leftrightarrow 3,z_1\leftrightarrow z_2)$ is realized by the identity (at the level of the square it is the flip in the diagonal [1 2]).

Because of (3.16) the discriminant of the quadratic equation (3.29) which defines v_1 is $\mathbf{\Delta}:=(X_0^T\sqrt{R_{z_1}}\sqrt{R_{z_2}}X_3-1)^2-z_1z_2|\hat{N}_0^0|^2|\hat{N}_0^3|^2$ for QC, $:=(x_{z_1u}(0,0)^T\hat{N}_0^0x_{z_2v}(0,0)^T\hat{N}_0^3+x_{z_1v}(0,0)^T\hat{N}_0^0x_{z_2u}(0,0)^T\hat{N}_0^3-f^T(x_{z_1}^0+x_{z_2}^3))^2-z_1z_2|\hat{N}_0^0|^2|\hat{N}_0^3|^2=(Z_0^T\sqrt{R'_{z_1}}\sqrt{R'_{z_2}}I_{1,2}Z_3-(Z_0+Z_3)^T(u(z_1+z_2)f_1+v(z_1+\epsilon z_2)\bar{f}_1+e_3)-(u(z_1)-u(z_2))(v(z_1)-v(z_2))-\frac{|f|^2}{2}(z_1+z_2))^2-z_1z_2|\hat{N}_0^0|^2|\hat{N}_0^3|^2$ for (I)QWC, where $\epsilon:=-1$ for IQWC (3.9) and $:=1$ otherwise. Thus $(0\leftrightarrow 3,1\leftrightarrow 2)\circ\mathbf{\Delta}=\mathbf{\Delta}$ (note also that the first squared terms are obtained by polarization* from (3.14)) and we have:

$$\frac{1}{2}Y'(v_1)^T(\sqrt{R_{z_2}}X_3-\sqrt{R_{z_1}}X_0-i(\sqrt{R_{z_2}}X_3)\times(\sqrt{R_{z_1}}X_0))=\pm\sqrt{\mathbf{\Delta}}=$$
$$\pm\frac{1}{2}Y'(v_2)^T(\sqrt{R_{z_2}}X_0-\sqrt{R_{z_1}}X_3-i(\sqrt{R_{z_2}}X_0)\times(\sqrt{R_{z_1}}X_3))\text{ for QC,}$$
$$2v_1(x_{z_1v}(0,0)^T\hat{N}_0^0-x_{z_2v}(0,0)^T\hat{N}_0^3)+f^TV_0^3-x_{z_1v}(0,0)^T\hat{N}_0^0x_{z_2u}(0,0)^T\hat{N}_0^3+$$
$$x_{z_1u}(0,0)^T\hat{N}_0^0x_{z_2v}(0,0)^T\hat{N}_0^3=\pm\sqrt{\mathbf{\Delta}}=\pm(2v_2(x_{z_1v}(0,0)^T\hat{N}_0^3-x_{z_2v}(0,0)^T\hat{N}_0^0)+f^TV_3^0-$$
$$x_{z_1v}(0,0)^T\hat{N}_0^3x_{z_2u}(0,0)^T\hat{N}_0^0+x_{z_1u}(0,0)^T\hat{N}_0^3x_{z_2v}(0,0)^T\hat{N}_0^0)\text{ for (I)QWC.}\tag{3.31}$$

For QC (3.29) has the symmetry $(u_j\leftrightarrow v_j,\ j=0,1,3)$ (since reflection in the origin preserves the SITC), so the lhs of (3.31) is skew wrt $(u_j\leftrightarrow v_j,\ j=0,1,3)$; for (I)QWC the lhs of (3.31) is skew wrt $u_1\leftrightarrow v_1$ (use $x_{z_ju}(0,0)\leftrightarrow x_{z_jv}(0,0),\ j=1,2$ and (3.17)).

We first choose the $-$ sign in the rhs of (3.31) (this is the choice of x_0^2). Replacing $u_0,\ u_3$ from (3.15), (3.31) becomes a quadratic polynomial of u_1 being identically 0; on account of the homography between the rulings $x_{z_1u_0}^0,\ x_{z_2u_3}^3$ it is enough to give u_1 a particular value (the relation thus found between $v_j,\ j=0,...,3$ will guarantee that the required quadratic polynomial in u_1 is identically 0). If we let $u_1:=\infty$, then we obtain for all quadrics except (3.3) with $a_1\neq a_2$ a homography (as a polynomial it has degree 2) between $v_j,\ j=0,...,3:\ \phi=\phi(z_1,z_2,v_0,v_1,v_2,v_3)=0$ having the required symmetries of the SITC; for (3.3) with $a_1\neq a_2$ we obtain a polynomial of degree 6 : $\varphi=\varphi(z_1,z_2,v_0,v_1,v_2,v_3)=0$ and missing some symmetries. Symmetrizing φ we get a homography (as a polynomial it has degree 4) with all the required symmetries ($\varphi':=\varphi-\varphi\circ(v_0\leftrightarrow v_3,v_1\leftrightarrow v_2)$ is still valid and gains the symmetry $(v_0\leftrightarrow v_3,v_1\leftrightarrow v_2)$; finally $\phi:=\frac{\varphi'-\varphi'\circ(v_0\leftrightarrow v_1,v_2\leftrightarrow v_3)}{v_0v_2-v_1v_3}$ is a homography with all the required symmetries). Now one can easily factor φ as the product of ϕ and a polynomial of degree 2 (linear in v_0,v_1,v_3) by accounting the terms containing $v_0v_1v_2v_3$; the remaining terms also satisfy it.

* This polarization is clear at the analytic level from the ansatz $z_1=z_2$, but it may also have another geometric explanation as in higher dimensions probably another discriminant must also behave well. For this reason the computations of Jacobi's higher dimensional version and its corresponding discretization *ray of light* must be similar: there also the discriminant behaved well because a solution (namely 0) of a quadratic equation was known; as a consequence other solutions of a polynomial of degree $n-1$ with 0 among its roots also behaved well. Thus one can consider the polarization of the next geometric statement: a line tangent to x_0 at x_0^1 cuts a confocal quadric x_z at the known point x_z^0 and thus at another computable point x_z^3; now one *geometrically* polarizes by *wiggling* the points x_0^0,x_0^3 and the z (separate z_1's and z_2's stick to each of x_0^0,x_0^3).

The symmetrization process to find (3.33) for QC (3.3) with $a_1 \neq a_2$ is complicated; once (3.33) is known, the factorization of φ as above is easily achieved. Therefore we propose a change in Bianchi's approach: while he first completes the discussion of the SITC & BPT and then proves the cross-ratio property as a corollary, we insert the cross-ratio property in the discussion of the SITC, thus easily arriving at (3.33)-(3.35); further the above approach by choosing $u_1 := \infty$ (and factorization of φ for QC (3.3) with $a_1 \neq a_2$) turns out to be unnecessary.

Consider the cross ratio property: the ruling $x^1_{0u_1}$ cuts the line $[x^0_{z_1}\ x^3_{z_2}]$ at $x_0(\hat{u}_1, v_1) := x^1_0 + \frac{\hat{u}_1 - u_1}{\hat{u}_1 - v_1}(u_1 - v_1)x^1_{0u_1}$ for QC or $x^1_0 + (\hat{u}_1 - u_1)x^1_{0u_1}$ for (I)QWC; therefore $(x_0(\hat{u}_1, v_1) - x^0_{z_1}) \times (x^0_{z_1} - x^3_{z_2}) = 0$, or multiplying on the left with $(x^0_{z_1})^T$: $x_0(\hat{u}_1, v_1)^T(x^0_{z_1} \times x^3_{z_2}) = 0$. Thus $x_0(\hat{u}_1, v_1) = \frac{(x^1_{0u_1} \times x^1_0) \times (x^0_{z_1} \times x^3_{z_2})}{(x^1_{0u_1})^T(x^0_{z_1} \times x^3_{z_2})}$ and $(x_0(\hat{u}_1, v_1) - x^0_{z_1}) \div (x_0(\hat{u}_1, v_1) - x^3_{z_2}) = \frac{(x^0_{z_1} \times x^3_{z_2})^T(x^1_{0u_1} \times V^0_1)}{(x^0_{z_1} \times x^3_{z_2})^T(x^1_{0u_1} \times V^3_1)} = \frac{(\hat{N}^1_0)^T(x^1_{0u_1} \times V^0_1)}{(\hat{N}^1_0)^T(x^1_{0u_1} \times V^3_1)} =^{(3.22)} \frac{(x^1_{z_1u_1})^T \hat{N}^0_0}{(x^1_{z_2u_1})^T \hat{N}^3_0} (=^{(3.15)} \frac{1 - \frac{1}{2}Y'(v_1)^T\sqrt{R_{z_1}}X_0}{1 - \frac{1}{2}Y'(v_1)^T\sqrt{R_{z_2}}X_3}$ for QC). We thus need:

$$\frac{(1 - \frac{1}{2}Y'(v_1)^T\sqrt{R_{z_1}}X_0)(1 - \frac{1}{2}Y'(v_2)^T\sqrt{R_{z_1}}X_3)}{(1 - \frac{1}{2}Y'(v_1)^T\sqrt{R_{z_2}}X_3)(1 - \frac{1}{2}Y'(v_2)^T\sqrt{R_{z_2}}X_0)} = \frac{z_1}{z_2} \text{ for QC,}$$

$$\frac{(x^1_{z_1u_1})^T\hat{N}^0_0(x^2_{z_1u_2})^T\hat{N}^3_0}{(x^1_{z_2u_1})^T\hat{N}^3_0(x^2_{z_2u_2})^T\hat{N}^0_0} = \frac{z_1}{z_2} \text{ for (I)QWC.} \tag{3.32}$$

We have to prove that the cross-ratio property (3.32) or its counterpart obtained by choosing the other ruling family on x^2_0 is equivalent to the SITC. Let $\varphi_+ := \frac{(x^1_{z_1u_1})^T\hat{N}^0_0(x^2_{z_1u_2})^T\hat{N}^3_0}{(x^1_{z_2u_1})^T\hat{N}^3_0(x^2_{z_2u_2})^T\hat{N}^0_0}$, $\varphi^{\pm}_+ := \frac{(x^1_{z_1u_1})^T\hat{N}^0_0}{(x^1_{z_2u_1})^T\hat{N}^3_0} \pm \frac{(x^1_{z_1v_1})^T\hat{N}^0_0}{(x^1_{z_2v_1})^T\hat{N}^3_0}$, $\varphi_- := \frac{(x^1_{z_1u_1})^T\hat{N}^0_0(x^2_{z_1v_2})^T\hat{N}^3_0}{(x^1_{z_2u_1})^T\hat{N}^3_0(x^2_{z_2v_2})^T\hat{N}^0_0}$, $\varphi^{\pm}_- := (0 \leftrightarrow 3, 1 \leftrightarrow 2) \circ \varphi^{\pm}_+$. By (3.12) we have $\frac{(x^1_{z_1v_1})^T\hat{N}^0_0(x^2_{z_1v_2})^T\hat{N}^3_0}{(x^1_{z_2v_1})^T\hat{N}^3_0(x^2_{z_2v_2})^T\hat{N}^0_0} = \frac{z_1^2}{z_2^2\varphi_+}$, $\frac{(x^1_{z_1v_1})^T\hat{N}^0_0(x^2_{z_1u_2})^T\hat{N}^3_0}{(x^1_{z_2v_1})^T\hat{N}^3_0(x^2_{z_2u_2})^T\hat{N}^0_0} = \frac{z_1^2}{z_2^2\varphi_-}$, so we need to prove $\frac{2z_1}{z_2} = \frac{(\varphi^+_+\varphi^+_-) \pm (\varphi^-_+\varphi^-_-)}{2} (= \varphi_+ + \frac{z_1^2}{z_2^2\varphi_+}$ for the + sign or $\varphi_- + \frac{z_1^2}{z_2^2\varphi_-}$ for the − sign). For QC we have $\frac{-z_2|\hat{N}^3_0|^2\varphi^+_+}{2} = X_0^T\sqrt{R_{z_1}}\sqrt{R_{z_2}}X_3 - 1 = \frac{-z_2|\hat{N}^0_0|^2\varphi^+_-}{2}$, $\frac{z_2|\hat{N}^3_0|^2\varphi^-_+}{2} = -iX_1^T((\sqrt{R_{z_2}}X_3) \times \sqrt{R_{z_1}}X_0) =$lhs of (3.31), $\frac{z_2|\hat{N}^0_0|^2\varphi^-_-}{2} = -iX_2^T((\sqrt{R_{z_2}}X_0) \times \sqrt{R_{z_1}}X_3) =$rhs of (3.31) with the + sign; for (I)QWC and using (3.17) we have $\frac{-z_2|\hat{N}^3_0|^2\varphi^+_+}{2} = x_{z_1u}(0,0)^T\hat{N}^0_0x_{z_2v}(0,0)^T\hat{N}^3_0 + x_{z_1v}(0,0)^T\hat{N}^0_0x_{z_2u}(0,0)^T\hat{N}^3_0 - f^T(x^0_{z_1} + x^3_{z_2}) = \frac{-z_2|\hat{N}^0_0|^2\varphi^+_-}{2}$, $\frac{z_2|\hat{N}^3_0|^2\varphi^-_+}{2} =$lhs of (3.31), $\frac{z_2|\hat{N}^0_0|^2\varphi^-_-}{2} =$rhs of (3.31) with the + sign, so $\varphi^+_+\varphi^+_- = \frac{4\boldsymbol{\Delta}}{z_2^2|\hat{N}^0_0|^2|\hat{N}^3_0|^2} + \frac{4z_1}{z_2}$, $\varphi^-_+\varphi^-_- = \pm\frac{4\boldsymbol{\Delta}}{z_2^2|\hat{N}^0_0|^2|\hat{N}^3_0|^2}$, where $\boldsymbol{\Delta}$ is the discriminant of the quadratic equation (3.29) which defines v_1.

Note that (3.31) can be interpreted as a homography between u_0, u_3 : $C_1u_0u_3 + C_2u_0 + C_3u_3 + C_4 = 0$, $C_j = C_j(z_1, z_2, v_0, v_1, v_2, v_3)$, $j = 1, ..., 4$ and (3.32) as the same homography between u_0, u_3 : $D_1u_0u_3 + D_2u_0 + D_3u_3 + D_4 = 0$, $D_j = D_j(z_1, z_2, v_0, v_1, v_2, v_3)$, $j = 1, ..., 4$; thus C_j, D_j must be proportional: $D_j = \eta C_j$, $j = 1, ..., 4$, $\eta = (1 \leftrightarrow 2, z_1 \leftrightarrow z_2) \circ \eta$; this condition establishes the sought homographies between v_j, $j = 0, ..., 3$. For QC accounting the coefficients of u_1, u_2 from $\frac{D_2}{C_2} = \frac{D_3}{C_3}$ we get $\eta = \frac{v_0v_3z_1f_1^T\sqrt{R_{z_2}}f_1(1 + e_3^T\sqrt{R_{z_2}}e_3) - 2z_2\bar{f}_1^T\sqrt{R_{z_1}}f_1(1 - e_3^T\sqrt{R_{z_1}}e_3)}{v_0v_3f_1^T\sqrt{R_{z_2}}f_1(1 - e_3^T\sqrt{R_{z_1}}e_3) - 2\bar{f}_1^T\sqrt{R_{z_1}}f_1(1 + e_3^T\sqrt{R_{z_2}}e_3)} (= \frac{1 - e_3^T\sqrt{R_{z_1}}e_3}{1 + e_3^T\sqrt{R_{z_2}}e_3}z_2$ for QC (3.4),(3.5)); for all (I)QWC except QWC (3.7) $\eta = \frac{D_1}{C_1} = \frac{z_1f_1^T\sqrt{R'_{z_2}}f_1}{2\bar{f}_1^T\sqrt{R'_{z_1}}f_1}$; for QWC (3.7) the homography between v_j, $j = 0, ..., 3$ is obtained directly from (3.32); taking $u_1 = \infty$ in (3.31) we obtain the same homography times a constant multiple of v_1.

We shall use the notation $f(z_1 \wedge z_2)$ for $f(z_1, z_2)$ if $f(z_2, z_1) = -f(z_1, z_2)$ and $f(z_1 \odot z_2)$ for $f(z_1, z_2)$ if $f(z_2, z_1) = f(z_1, z_2)$.

For QC let $M_2(z) := z^{-1}((\mathrm{tr}\sqrt{R_z})I_3 - \sqrt{R_z})(I_3 + \sqrt{R_z}) = z^{-1}(\mathrm{tr}(\det\sqrt{R_z}(\sqrt{R_z})^{-1} + \sqrt{R_z})I_3 - (\det\sqrt{R_z}(\sqrt{R_z})^{-1} + \sqrt{R_z}))$, $M_3(z_1 \wedge z_2) := ((\mathrm{tr}(\sqrt{R_{z_1}})I_3 - \sqrt{R_{z_1}})(\mathrm{tr}(\sqrt{R_{z_2}})I_3 - \sqrt{R_{z_2}}) - - \mathrm{tr}(\sqrt{R_{z_1}}\sqrt{R_{z_2}})I_3 + \sqrt{R_{z_1}}\sqrt{R_{z_2}})(\sqrt{R_{z_2}} - \sqrt{R_{z_1}})^{-1}$, $m_2(z) := e_3^TM_2(z)e_3$, $m_3(z_1 \wedge z_2) :=$

$e_3^TA^{-1}M_3(z_1\wedge z_2)e_3(=(\frac{1}{m_2(z_1)}-\frac{1}{m_2(z_2)})^{-1}$ if $\mathrm{spec}(A)=\{a_1^{-1}\})$, $a:=\bar{f}_1^TA\bar{f}_1e_3^TAe_3$ for QC (3.3),(3.4), $:=a_1^{-3}$ for QC (3.5), $\delta:=1$ for QC (3.3) and $:=0$ otherwise.

For (I)QWC let $M_3(z_1\wedge z_2):=2((\mathrm{tr}(\sqrt{R'_{z_1}})I_3-\sqrt{R'_{z_1}})(\mathrm{tr}(\sqrt{R'_{z_2}})I_3-\sqrt{R'_{z_2}})-\mathrm{tr}(\sqrt{R'_{z_1}}\sqrt{R'_{z_2}})I_3+\sqrt{R'_{z_1}}\sqrt{R'_{z_2}})(z_1-z_2)^{-1}$, $m_2(z):=2z^{-1}(\mathrm{tr}(\sqrt{R'_z})-1)$, $m_3(z_1\wedge z_2):=e_3^TM_3(z_1\wedge z_2)e_3(=(\frac{1}{m_2(z_1)}-\frac{1}{m_2(z_2)})^{-1}$ for IQWC), $a:=f_1^TA'f_1$, $k:=\frac{-1}{2\sqrt{2}}$ for IQWC (3.8).

The homography between $v_j,\ j=0,...,3$ is:

$$\frac{v_0v_3+v_1v_2}{m_3(z_1\wedge z_2)}+\frac{v_0v_2+v_1v_3}{m_2(z_2)}-\frac{v_0v_1+v_2v_3}{m_2(z_1)}+a\frac{4+\delta v_0v_1v_2v_3}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}=0$$
$$\text{for QC (3.3), (3.4), QWC (3.6), (3.7);} \tag{3.33}$$

$$\frac{v_0v_3+v_1v_2}{m_3(z_1\wedge z_2)}+\frac{v_0v_2+v_1v_3}{m_2(z_2)}-\frac{v_0v_1+v_2v_3}{m_2(z_1)}-2a\frac{v_0+v_1+v_2+v_3}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}+(\frac{1}{m_3(z_1\wedge z_2)^2}+\frac{1}{m_2(z_1)m_2(z_2)})\frac{4a^2}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}=0 \text{ for QC (3.5), IQWC (3.9);} \tag{3.34}$$

$$\frac{v_0v_3+v_1v_2}{m_3(z_1\wedge z_2)}+\frac{v_0v_2+v_1v_3}{m_2(z_2)}-\frac{v_0v_1+v_2v_3}{m_2(z_1)}-4k\frac{v_0+v_1+v_2+v_3+k(z_1+z_2)}{a_1m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}-\frac{k^2z_1z_2}{m_3(z_1\wedge z_2)}=0 \text{ for IQWC (3.8).} \tag{3.35}$$

Note that the above homography for QC can be written as

$$X(v_0,v_3)^T(M_2(z_1)-M_2(z_2))(M_2(z_1)+M_2(z_2))^{-1}X(v_1,v_2)=1.$$

The relations (3.31) and (3.32) can be interpreted as the same homography between v_0,v_3, thus establishing a homography between u_0,v_1,v_2,u_3 analogue to (3.33)-(3.35). For QC we just perform the change $(u_0\leftrightarrow v_0,u_3\leftrightarrow v_3,\sqrt{R_{z_j}}\leftrightarrow-\sqrt{R_{z_j}})$; for QWC (3.6) and IQWC (3.8) the change $(u_0\leftrightarrow v_0,u_3\leftrightarrow v_3,\sqrt{R'_{z_j}}\leftrightarrow r_{e_2}\sqrt{R'_{z_j}})$. For QWC (3.7) we obtain a new homography (m_2^-, m_3 have same definition with $\sqrt{R'_{z_j}}$ replaced by $\sqrt{R'_{z_j}}r_{e_2}$ and it is the reason why m_3 acquires the $-$ sign; $a^-:=\bar{f}_1^TA'\bar{f}_1$):

$$-\frac{2v_1v_2}{m_3(z_1\wedge z_2)}+\frac{u_0v_2+v_1u_3}{m_2^-(z_2)}-\frac{u_0v_1+v_2u_3}{m_2^-(z_1)}-\frac{2a^-}{m_2^-(z_1)m_2^-(z_2)m_3(z_1\wedge z_2)}=0$$
$$\text{for QWC (3.7);} \tag{3.36}$$

for IQWC (3.9) we must perform $(u_j\leftrightarrow v_j,\ j=1,2)$ in (3.31) and consider the other ruling families on $x_0^1,\ x_0^2$ in (3.32); the new (3.31), (3.32) are valid and can be interpreted as the same homography between u_0,u_3, thus establishing the homography (m_2^-, m_3 have same definition with $\sqrt{R'_{z_j}}$ replaced by $r_{e_2}\sqrt{R'_{z_j}}$):

$$-\frac{2u_1u_2}{m_3(z_1\wedge z_2)}+\frac{v_0u_2+u_1v_3}{m_2^-(z_2)}-\frac{v_0u_1+u_2v_3}{m_2^-(z_1)}+2\frac{v_0+v_3}{m_3(z_1\wedge z_2)}+2(\frac{1}{m_2(z_1)}+\frac{1}{m_2(z_2)})\frac{u_1+u_2}{m_3(z_1\wedge z_2)}-\frac{4}{m_3(z_1\wedge z_2)^3}-\frac{8}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}=0 \text{ for IQWC (3.9).} \tag{3.37}$$

Because of the symmetry $(0\leftrightarrow1,2\leftrightarrow3)$ of the SITC, the homography between v_0,u_1,u_2,v_3 is valid with a similar formula: for QC we just perform the change $(u_1\leftrightarrow v_1,u_2\leftrightarrow v_2,\sqrt{R_{z_j}}\leftrightarrow-\sqrt{R_{z_j}})$; for QWC (3.6) and IQWC (3.8) the change $(u_1\leftrightarrow v_1,u_2\leftrightarrow v_2,\sqrt{R'_{z_j}}\leftrightarrow r_{e_2}\sqrt{R'_{z_j}})$; (3.36) is valid for $(u_0,v_1,v_2,u_3)\to(u_1,v_0,v_3,u_2)$ and (3.37) is valid for $(v_0,u_1,u_2,v_3)\to(v_1,u_0,u_3,v_2)$.

Because of the first relation of (3.12), (3.32) remains valid if we consider the other ruling families on $x_0^1,\ x_0^2$ and since $(u_j\leftrightarrow v_j,\ j=0,...,3)\circ(3.31)$, $(u_j\leftrightarrow v_j,\ j=1,2)\circ(3.31)$ remain valid with

the same sign respectively for QC, (I)QWC, the new (3.31), (3.32) can be interpreted as the same homography between $v_0,\ v_3$; this establishes homographies between $u_j,\ j=0,...,3$ analogue to (3.33)-(3.35): for QC, QWC (3.6) and IQWC (3.8) we perform just the change $u_j \leftrightarrow v_j,\ j=0,...,3$ (because of the involutory character of the change in $\sqrt{R_{z_j}}$ ($\sqrt{R'_{z_j}}$) and the change being done twice: once for $(u_0 \leftrightarrow v_0, u_3 \leftrightarrow v_3)$ and once for $(u_1 \leftrightarrow v_1, u_2 \leftrightarrow v_2)$); for the remaining quadrics we get a new homography (m_2, m_3 remain the same, since replacing $\sqrt{R'_{z_j}}$ by $r_{e_2}\sqrt{R'_{z_j}}r_{e_2}$ does not change either; $a^- := -f_1^T A' f_1$ for IQWC (3.9)); for IQWC (3.9) it is obtained directly from the new (3.32):

$$\frac{u_0u_3+u_1u_2}{m_3(z_1\wedge z_2)}+\frac{u_0u_2+u_1u_3}{m_2(z_2)}-\frac{u_0u_1+u_2u_3}{m_2(z_1)}+\frac{4a^-}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)}=0 \tag{3.38}$$
$$\text{for QWC (3.7) \& IQWC (3.9).}$$

For the other choice of the sign in (3.31) perform in (3.31) $u_j \leftrightarrow v_j,\ j=0,2,3$ ($u_2 \leftrightarrow v_2$) respectively for QC ((I)QWC) and consider in (3.32) the other ruling family on x_0^2; again the new (3.31), (3.32) can be interpreted as the same homography between $u_0,\ v_3$; this establishes a homography between v_0, v_1, u_2, u_3 analogue to (3.33)-(3.35): for QC we just perform the change $(u_2 \leftrightarrow v_2, u_3 \leftrightarrow v_3, \sqrt{R_{z_2}} \leftrightarrow -\sqrt{R_{z_2}})$; for QWC (3.6) and IQWC (3.8) the change $(u_2 \leftrightarrow v_2, u_3 \leftrightarrow v_3, \sqrt{R'_{z_2}} \leftrightarrow r_{e_2}\sqrt{R'_{z_2}})$. For the remaining quadrics we obtain a new homography (m_3^- has same definition as m_3 with $\sqrt{R'_{z_2}}$ replaced by $\sqrt{R'_{z_2}}r_{e_2}$ for QWC (3.7) and $r_{e_2}\sqrt{R'_{z_2}}$ for IQWC (3.9), but it loses its skew-symmetry):

$$\frac{v_0u_3+v_1u_2}{m_3^-(z_1\odot z_2)}+\frac{v_0u_2+v_1u_3}{m_2^-(z_2)}-\frac{2v_0v_1}{m_2(z_1)}+\frac{2a^-}{m_2(z_1)m_2^-(z_2)m_3^-(z_1\odot z_2)}=0 \tag{3.39}$$
$$\text{for QWC (3.7);}$$

$$\frac{v_0u_3+v_1u_2}{m_3^-(z_1\odot z_2)}+\frac{v_0u_2+v_1u_3}{m_2^-(z_2)}-\frac{2u_2u_3}{m_2(z_1)}+2\frac{v_0+v_1}{m_2(z_1)}+2(\frac{2}{m_2(z_2)}+\frac{1}{m_2(z_1)})\frac{u_2+u_3}{m_2(z_1)}-$$
$$\frac{2}{m_2(z_1)^3}-(\frac{2}{m_2(z_2)}+\frac{1}{m_2(z_1)})^2\frac{2}{m_2(z_1)}=0 \text{ for IQWC (3.9).} \tag{3.40}$$

Choosing $u_1 := \infty$ in (3.31) with the + sign we get the homography between v_0, v_1, u_2, u_3 for QC, QWC (3.6), IQWC (3.8) (this fact follows from the similar statement for the homography between v_0, v_1, v_2, v_3) and IQWC (3.9); for QWC (3.7) we get the left side of (3.39) times a constant multiple of $v_0(\bar{f}_1^T\sqrt{R'_{z_2}}f_1)^2 - v_1\bar{f}_1^T\sqrt{R'_{z_1}}f_1$.

Further we can consider in this case the corresponding homographies between the other ruling families, but since the homography between v_0, v_1, u_2, u_3 is invariant only under the symmetry $(0 \leftrightarrow 1, 2 \leftrightarrow 3)$ of the SITC, all other symmetries of the SITC generate the sought homographies: $(0 \leftrightarrow 3, 1 \leftrightarrow 2)$, $(0 \leftrightarrow 3, z_1 \leftrightarrow z_2)$, $(1 \leftrightarrow 2, z_1 \leftrightarrow z_2)$ respectively generate the homographies between $u_0, u_1, v_2, v_3;\ u_0, v_1, u_2, v_3;\ v_0, u_1, v_2, u_3$.

Thus the rulings of x_0^0, x_0^3 and x_0^1, x_0^2 involved in a homography are simultaneously same or different; this dichotomy gives the two choices of x_0^2 and can be interpreted as a $\mathbb{Z}_2$ co-cycle: if we assign the signature $\epsilon := 1\ (-1)$ to $v\ (u)$ rulings, then $\epsilon_0\epsilon_1\epsilon_2\epsilon_3 = 1$.

If the homographies (3.33)-(3.35) are written as $\phi = \phi(v_0, (v_1, z_1) \wedge (v_2, z_2), v_3) = 0$, then the algebraic relation:

$$\frac{z_1\mathcal{A}(\phi\phi_{v_2v_3}-\phi_{v_2}\phi_{v_3})}{\Delta^-(z_1,v_0,v_1)}=\frac{1}{m_2(z_1)m_2(z_2)m_3(z_1\wedge z_2)} \tag{3.41}$$

for independent variables $v_j,\ j=0,...,3$ plays an important rôle in the analytic computations of the Möbius configurations and in the proof of the BPT; it remains valid if we perform changes $u_j \leftrightarrow v_j$ for some j's according to the previous recipes (in which case Δ^- may change to Δ^+ or

$\Delta'^{\pm}$; m_2 to m_2^-; m_3 to $-m_3$ or $\pm m_3^-$). Bianchi (122,§25) assumes its use in the proof of the BPT without explicitly stating it.

The homography H of Bianchi II preserves the Ivory affinity, the RMPIA and the (SI)TC; therefore all results of the last three subsections. The preservation of the SITC is suggested by the form of the homographies (3.33)-(3.40) and can be confirmed analytically if one takes into consideration the change of $z_k,\ k=1,2,\ A,\ u_j,\ v_j,\ j=0,...,3$ under H, but an obvious geometric argument can replace these analytic computations.

3.6. **Möbius configurations and an application of the Menelaus theorem.**

A tetrahedron consists of 2^2 points and 2^2 planes, each point (plane) belonging to (containing) $2+1$ planes (points). Möbius considered configurations $\mathcal{M}_3$ of two tetrahedra inscribed one into the other, that is configurations of 2^3 points and 2^3 planes, each point (plane) belonging to (containing) $3+1$ planes (points); two $\mathcal{M}_3$ configurations inscribed one into the other gives raise to a configuration of 2^4 points and 2^4 planes, each point (plane) belonging to (containing) $4+1$ planes (points). Therefore Bianchi ([5],Vol **5**,(117)) calls a configuration of 2^n points and 2^n planes such that any point (plane) belongs to (contains) $n+1$ planes (points) a Möbius configuration $\mathcal{M}_n$. We can consider the TC as $\mathcal{M}_1$, the SITC as $\mathcal{M}_2$ and further iterates as $\mathcal{M}_n$'s. Let $n\geq 2,\ k=1,...,n,\ j,j_1,j_2,...=0,...,2^n-1,\ x_0^0\in x_0,\ z_k\in\mathbb{C},\ x_{z_k}^{2^{k-1}}\in x_{z_k}\cap T_{x_0^0}x_0$; choose rulings $\hat{v}_0(=v_0 \text{ or } u_0)$ at x_0^0 and $\hat{v}_{2^{k-1}}(=u_{2^{k-1}} \text{ or } v_{2^{k-1}})$ at $x_0^{2^{k-1}}$; as before we can order $0<z_1<...<z_n$. We would like to get by repeated application of the SITC well defined points $x_0^j\in x_0$ and rulings $\hat{v}_j$ at those points (and thus RMPIA's $(R_{j_1}^{j_2},t_{j_1}^{j_2}),\ \log_2|j_1-j_2|\in\mathbb{N}$; the Ivory affinities are between $x_0,\ x_{z_{\log_2|j_1-j_2|+1}}$). The rulings are well defined (the signature $\epsilon_j=\epsilon_0^{s-1}\epsilon_{k_1}...\epsilon_{k_s}$ for $0<k_1<...<k_s,\ j=\Sigma_{p=1}^s 2^{k_p-1}$ is independent of the path from x_0^0 to x_0^j). To define x_0^j apply repeatedly the SITC to a path $x_0^{j_0},x_0^{j_1},...x_0^{j_s},\ 0=j_0<j_1<...<j_s=j,\ \log_2(j_p-j_{p-1})\in\mathbb{N},\ p=1,...,s$; this definition must be independent of the chosen path (assume this for the moment); $\mathcal{M}_n$ is formed by facets $(R_{j_1}^{j_0},t_{j_1}^{j_0})\circ(R_{j_2}^{j_1},t_{j_2}^{j_1})\circ...\circ(R_{j_s}^{j_{s-1}},t_{j_s}^{j_{s-1}})(x_0^j,dx_0^j)$ (because of the co-cycle relations (3.30) these are independent of the sequence $j_0,j_1,j_2,...,j_s$ as above). If the rulings are not prescribed, then for $n\geq 2$ we can choose $\hat{v}_0:=v_0$ (since considering other rulings on all $x_0^0,\ x_{z_k}^{2^{k-1}}$ does not change $\mathcal{M}_n$), so we have $\frac{2^n}{2}=2^{n-1}$ choices of $\mathcal{M}_n$ (among the 2^n choices of rulings on $x_{z_k}^{2^{k-1}}$ we have to remove $\frac{1}{2}$ of them, since the dichotomy 'simultaneously same or different' allows to consider the other rulings on all $x_{z_k}^{2^{k-1}}$); for $n=1$ we have two choices of $\mathcal{M}_1$.

We need now prove that the definition of x_0^j does not depend on the path from x_0^0 to x_0^j; it is enough to consider $n=3$, since by induction if all $x_0^j,\ j=0,...,2^{n+1}-2$ are well defined, then we can uniquely define $x_0^{2^{n+1}-1}$ from the quadrilateral $x_0^{2^{n-1}-1},x_0^{2^n-1},x_0^{2^n+2^{n-1}-1},x_0^{2^{n+1}-1}$; if we separately add each $x_0^{2^{n+1}-2^{k-1}-1},\ k=1,...,n-1$ to this quadrilateral and complete it to an $\mathcal{M}_3$, then $x_0^{2^{n+1}-1}$ is well defined for each $\mathcal{M}_3$ obtained, so $x_0^{2^{n+1}-1}$ is well defined for all paths from x_0^0 to $x_0^{2^{n+1}-1}$. For $n=3:\ k=1,2,3,\ z_k\in\mathbb{C},\ x_0^0\in x_0,\ x_{z_k}^{2^{k-1}}\in x_{z_k}\cap T_{x_0^0}x_0$ and consider the rulings $v_0,\ v_{2^{k-1}}$; applying the SITC three times we get well defined facets $(x_0^j,dx_0^j),\ j=0,...,6$. To simplify the notation replace x_0^0 with $0,\ x_{z_1}^1$ with 1, etc. If the rulings v_1,v_4,v_2 respectively cut the lines [3 5], [5 6], [6 3] at the points $1_3^5,\ 4_5^6,\ 2_6^3$, then $((1_3^5-3)\div(1_3^5-5))((4_5^6-5)\div(4_5^6-6))((2_6^3-6)\div(2_6^3-3))=\frac{(x_{z_2u_1}^1)^T\hat{N}_0^3}{(x_{z_3u_1}^1)^T\hat{N}_0^5}\frac{(x_{z_1u_4}^4)^T\hat{N}_0^5}{(x_{z_2u_4}^4)^T\hat{N}_0^6}\frac{(x_{z_3u_2}^2)^T\hat{N}_0^6}{(x_{z_1u_2}^2)^T\hat{N}_0^3}=\frac{(x_{z_2u_1}^1)^T\hat{N}_0^3}{(x_{z_1u_2}^2)^T\hat{N}_0^3}\frac{(x_{z_1u_4}^4)^T\hat{N}_0^5}{(x_{z_3u_1}^1)^T\hat{N}_0^5}\frac{(x_{z_3u_2}^2)^T\hat{N}_0^6}{(x_{z_2u_4}^4)^T\hat{N}_0^6}=^{(3.32)}\frac{z_2}{z_1}\frac{(x_{z_1u_1}^1)^T\hat{N}_0^0}{(x_{z_2u_2}^2)^T\hat{N}_0^0}\times\frac{z_1}{z_3}\frac{(x_{z_3u_4}^4)^T\hat{N}_0^0}{(x_{z_1u_1}^1)^T\hat{N}_0^0}\frac{z_3}{z_2}\frac{(x_{z_2u_2}^2)^T\hat{N}_0^0}{(x_{z_3u_4}^4)^T\hat{N}_0^0}=1$. By the converse of the Menelaus theorem (which is essentially a co-cycle theorem) the points $1_3^5,\ 4_5^6,\ 2_6^3$ in the plane [3 5 6] are co-linear. Consider the points $7_3^5,\ 7_5^6,\ 7_6^3$ respectively on the lines [3 5], [5 6], [6 3] such that they and the previous points cut these lines with the required cross-ratios $\frac{z_j}{z_k}$; again by the converse of the Menelaus theorem $7_3^5,\ 7_5^6,\ 7_6^3$ lie on a line l_7; a similar statement holds for the other rulings u_1,u_2,u_4 and we obtain another line l_7': take $7:=l_7\cap l_7'$. By symmetry we need now only prove that 7 is in the facet of 5, since in this case the

rulings v_5, v_6 will cut the line [4 7] with same cross-ratio as the one the rulings l_7, v_4 cut the line [5 6] (v_4, v_5, v_6, l_7 will be rulings of the same ruling family on a quadric), so 7 completes $4, 5, 6$ to an SITC quadrilateral. We have $\frac{(x^0_{z_1u_0})^T\hat{N}^1_0}{(x^0_{z_2u_0})^T\hat{N}^2_0} - \frac{(x^0_{z_1v_0})^T\hat{N}^1_0}{(x^0_{z_2v_0})^T\hat{N}^2_0} =^{(3.22)} \frac{(V^1_0)^TR^0_2\hat{N}^2_0(\hat{N}^0_0)^T(x^0_{0v_0}\times x^0_{0u_0})}{\mathcal{A}|\hat{N}^0_0|^2(x^0_{z_2u_0})^T\hat{N}^2_0(x^0_{z_2v_0})^T\hat{N}^2_0} =^{(3.12)} -\frac{2(V^1_0)^TR^0_2\hat{N}^2_0}{z_2|\hat{N}^2_0|^2}$; since $7-5$ is a multiple of $\frac{z_3}{z_2}(\frac{(x^1_{z_2u_1})^T\hat{N}^3_0}{(x^1_{z_3u_1})^T\hat{N}^5_0} - \frac{(x^1_{z_2v_1})^T\hat{N}^3_0}{(x^1_{z_3v_1})^T\hat{N}^5_0})(6-4) - \frac{z_1}{z_2}(\frac{(x^4_{z_2u_4})^T\hat{N}^6_0}{(x^4_{z_1u_4})^T\hat{N}^5_0} - \frac{(x^4_{z_2v_4})^T\hat{N}^6_0}{(x^4_{z_1v_4})^T\hat{N}^5_0})(3-1)$, the condition $0=(7-5)^TR^0_1R^1_5\hat{N}^5_0$ becomes $0=(\hat{N}^5_0)^T(R^5_1R^1_0R^0_4V^6_4(R^5_1V^3_1)^T - R^5_1R^1_0R^0_1V^3_1(R^5_4V^6_4)^T)\hat{N}^5_0 =^{(3.30)} (\hat{N}^5_0)^T(R^5_4V^6_4(R^5_1V^3_1)^T - R^5_1V^3_1(R^5_4V^6_4)^T)\hat{N}^5_0$.

By an obvious geometric argument and induction on n, $\mathcal{M}_n$ has the symmetries of an n-dimensional cube, the static n-soliton $\hat{v}_{2^n-1}$ is a homography of $\hat{v}_0$, $\hat{v}_{2^{k-1}}$: $\hat{v}_{2^n-1} = \mathcal{S}_n(\hat{v}_0, (\hat{v}_1, z_1), ..., (\hat{v}_{2^{n-1}}, z_n))$; moreover symmetries of the n-dimensional cube which do not induce symmetries of this homography generate valid homographies between other rulings. Note that $\hat{v}_0$ does not appear in $\hat{v}_{2^n-1}$ for n odd: take for example $n=3$. To fix the ideas consider $\hat{v}_j := v_j$, $j=0,...,7$ satisfying the homographies imposed by the SITC; if we replace v_0, v_3, v_5, v_6 with u_0, u_3, u_5, u_6 we have similar homographies imposed by the SITC. Let now v_0 vary and keep u_0, v_1, v_2, v_4 constant; thus u_3, u_5 are constant; because of the homography between v_1, u_3, u_5, v_7 we conclude that v_7 is constant.

The 'addition' of static solitons now becomes clear: $\hat{v}_{j_1} \oplus_{\hat{v}_0} \hat{v}_{j_2} = \hat{v}_{j_1+j_2}$, where $1 \le j_1, j_2 \le 2^n-1$ have nonzero digits in base 2 on different positions. The SITC defines the law of 'addition' and proves its commutativity on static 1-solitons: $\hat{v}_1 \oplus_{\hat{v}_0} \hat{v}_2 = \hat{v}_2 \oplus_{\hat{v}_0} \hat{v}_1 = \hat{v}_3$; the symmetry of the TC provides an inverse. The existence of $\mathcal{M}_3$ assures the associativity of the 'addition'. Once these are checked on static 1-solitons $\hat{v}_{2^{k-1}}$, they are true for static n-solitons (the existence of $\mathcal{M}_2$, $\mathcal{M}_3$ assures the existence of $\mathcal{M}_n$).

An analytic confirmation of the above properties is more complicated, but all symmetries are revealed in the implicit form (homography) $\phi_n(\hat{v}_0, \bigwedge_{k=1}^n(\hat{v}_{2^{k-1}}, z_k), \hat{v}_{2^n-1}) = 0$ of $\mathcal{S}_n$: $\mathcal{S}_n(\hat{v}_0, \bigodot_{k=1}^n(\hat{v}_{2^{k-1}}, z_k)) = -\frac{\phi_n(\hat{v}_0, \bigwedge_{k=1}^n(\hat{v}_{2^{k-1}}, z_k), 0)}{\partial_v\phi_n(\hat{v}_0, \bigwedge_{k=1}^n(\hat{v}_{2^{k-1}}, z_k), v)}$. We use the notation $f(\bigodot_{k=1}^n x_k)$ for $f(x_1, ..., x_n)$ if $f(x_{\sigma(1)}, ..., x_{\sigma(n)}) = f(x_1, ..., x_n)$ and $f(\bigwedge_{k=1}^n x_k)$ for $f(x_1, ..., x_n)$ if $f(x_{\sigma(1)}, ..., x_{\sigma(n)}) = (-1)^{\epsilon(\sigma)}f(x_1, ..., x_n)$ for σ permutation of $1, ..., n$; although ϕ_n has skew-symmetries (and thus $\mathcal{S}_n$ has symmetries) only when separately permuting u rulings or v rulings, we still maintain for simplicity the $\bigwedge$ ($\bigodot$) notation for all rulings $\hat{v}_{2^{k-1}}$, $k=1, ..., n$. Also ϕ_n (and thus $\mathcal{S}_n$) depends on the signatures ϵ_0, $\epsilon_{2^{k-1}}$, $k=1, ..., n$; only when strictly necessary we shall use the notation ϕ^ε_n, $\varepsilon := 2^{n-1}(1-\epsilon_0) + \sum_{k=1}^n(1-\epsilon_{2^{k-1}})2^{k-2}$ to point out the explicit dependence on rulings. Since for n odd $\hat{v}_{2^n-1}$ does not depend on $\hat{v}_0$ we drop in this case $\hat{v}_0$ from $\phi_{n+1}(...)$.

Since IQWC (3.9) is metrically the most degenerate, (3.38) provides the simplest formulae for ϕ_n; thus it is natural to study it first and conjecture relations valid for all quadrics and rulings. We have $\phi_2(u_0, \bigwedge_{k=1,2}(u_{2^{k-1}}, z_k), u_3) = \begin{vmatrix}(u_1-u_0) & z_1(u_3-u_1)\\ (u_2-u_0) & z_2(u_3-u_2)\end{vmatrix} + \frac{1}{4}\begin{vmatrix}z_1 & z_1^2\\ z_2 & z_2^2\end{vmatrix}$,
$\phi_{2m+1}(u_0, \bigwedge_{k=1}^{2m+1}(u_{2^{k-1}}, z_k), u_{2^{2m+1}-1}) = \sum_{j=1}^{2m+1}(-1)^{j-1}\phi_{2m}(u_0, \bigwedge_{k=1,\ k\neq j}^{2m+1}(u_{2^{k-1}}, z_k), u_{2^{2m+1}-1})$,
$\phi_{2m+2}(u_0, \bigwedge_{k=1}^{2m+2}(u_{2^{k-1}}, z_k), u_{2^{2m+2}-1}) =$
$(\prod_{j=1}^{2m+2} z_j)\sum_{j=1}^{2m+2}\frac{(-1)^{j-1}(u_{2^{j-1}}-u_0)}{z_j}\phi_{2m+1}(u_0, \bigwedge_{k=1,\ k\neq j}^{2m+2}(u_{2^{k-1}}, z_k), u_{2^{2m+2}-1})$, $m \ge 1$. Thus ϕ_n is the linear combination of 2 determinants respectively with coefficients 1, $\frac{1}{4}$; ϕ_{2m+1} is obtained from ϕ_{2m} by replacing $u_{2^{2m}-1}$ with $u_{2^{2m+1}-1}$, then padding its determinants with a first column of 1's and with a last row of the same type as the previous (z_{2m+1}, $u_{2^{2m}}$ replace z_j, $u_{2^{j-1}}$); ϕ_{2m+2} is obtained from ϕ_{2m+1} by multiplying the j^{th} row of each of its determinants with z_j, replacing $u_{2^{2m+1}-1}$ with $u_{2^{2m+2}-1}$, then padding its determinants with a first column of $(u_{2^{j-1}} - u_0)_{j=1,...,2m+1}$ and with a last row of the same type as the previous (z_{2m+2}, $u_{2^{2m+1}}$ replace z_j, $u_{2^{j-1}}$); for example
$\phi_3(u_0, \bigwedge_{k=1}^3(u_{2^{k-1}}, z_k), u_7) = \begin{vmatrix}1 & u_1 & z_1(u_7-u_1)\\ 1 & u_2 & z_2(u_7-u_2)\\ 1 & u_4 & z_3(u_7-u_4)\end{vmatrix} + \frac{1}{4}\begin{vmatrix}1 & z_1 & z_1^2\\ 1 & z_2 & z_2^2\\ 1 & z_3 & z_3^2\end{vmatrix}$,

$$\phi_4(u_0,\bigwedge_{k=1}^4(u_{2^{k-1}},z_k),u_{15}) = \begin{vmatrix}(u_1-u_0) & z_1 & z_1u_1 & z_1^2(u_{15}-u_1)\\ (u_2-u_0) & z_2 & z_2u_2 & z_2^2(u_{15}-u_2)\\ (u_4-u_0) & z_3 & z_3u_4 & z_3^2(u_{15}-u_4)\\ (u_8-u_0) & z_4 & z_4u_8 & z_4^2(u_{15}-u_8)\end{vmatrix} + \frac{1}{4}\begin{vmatrix}(u_1-u_0) & z_1 & z_1^2 & z_1^3\\ (u_2-u_0) & z_2 & z_2^2 & z_2^3\\ (u_4-u_0) & z_3 & z_3^2 & z_3^3\\ (u_8-u_0) & z_4 & z_4^2 & z_4^3\end{vmatrix}.$$

Note that the separate linearity in $u_0,\ u_{2^{k-1}},\ u_{2^n-1}$ behavior mixes with the Vandermonde determinant in z_k behavior, so powers $\lessapprox \frac{n}{2}$ in z_k suffice to guarantee non-singular determinants.

The determinant formulation of ϕ_n is proven by induction after $n\geq 3$: consider the determinants $D_n^j(u_0,\bigwedge_{k=1}^n(u_{2^{k-1}},z_k)),\ j=1,2,3$, where $\phi_n(u_0,\bigwedge_{k=1}^n(u_{2^{k-1}},z_k),u_{2^n-1})=:$
$u_{2^n-1}D_n^1(u_0,\bigwedge_{k=1}^n(u_{2^{k-1}},z_k))+D_n^2(u_0,\bigwedge_{k=1}^n(u_{2^{k-1}},z_k))+\frac{1}{4}D_n^3(u_0,\bigwedge_{k=1}^n(u_{2^{k-1}},z_k))$. We have
$D_{n+1}^1(u_0,\bigwedge_{k=1}^{n+1}(u_{2^{k-1}},z_k))((u_{2^n}-u_0)^2-\frac{1}{4}z_{n+1}^2)^{[\frac{n}{2}]}(\frac{1+(-1)^n}{2}\prod_{k=1}^n z_k+\frac{1-(-1)^n}{2})=$
$D_n^1(u_{2^n},\bigwedge_{k=1}^n(u_{2^n+2^{k-1}},z_k))\prod_{k=1}^n D_2^1(u_0,(u_{2^{k-1}},z_k)\wedge(u_{2^n},z_{n+1}))$, $n\geq 2$ and a similar relation obtained by replacing $D_{n+1}^1(...),\ D_n^1(...)$ with $D_{n+1}^2(...)+\frac{1}{4}D_{n+1}^3(...),\ D_n^2(...)+\frac{1}{4}D_n^3(...)$; since $\phi_n(u_{2^n},\bigwedge_{k=1}^n(u_{2^n+2^{k-1}},z_k),u_{2^{n+1}-1})=0$ this will suffice. Just as the Vandermonde determinant V_{n+1} in $z_k,\ k=1,...,n+1$ can be interpreted as a polynomial of degree n in z_{n+1}, with roots $z_k,\ k=1,...,n$ and V_n as the dominant coefficient, each of the two relations between the functionally independent $u_0,\ u_{2^{k-1}},\ z_k,\ k=1,...,n+1$ can be interpreted as polynomials in z_n respectively of degrees $[\frac{n+1}{2}],\ [\frac{n+1}{2}]+1$, with roots among $z_k,\ k=1,...,n-1,n+1$ and whose dominant coefficients are 0. The condition that the dominant coefficients of the polynomials in z_n are 0 can be interpreted in turn as two polynomials in z_{n-1} of degrees $[\frac{n+1}{2}]-1,\ [\frac{n+1}{2}]$, with roots among $z_k,\ k=1,...,n-2,n+1$ and whose dominant coefficients are 0 (the dominant coefficients are linear in $u_{2^{n-2}}$; the coefficients of $u_{2^{n-2}}$ being 0 are the relations that must be initially proved, but with $n-2$ replacing n; the other coefficients being 0 are also valid relations proven by backward induction with step two). To prove the fact that roots of the considered polynomials are among $z_k,\ k=1,...,n-1,n+1$ or $k=1,...,n-2,n+1$ we need $z_j=z_k\Rightarrow u_{2^{j-1}}=u_{2^{k-1}}$; take for example $z_1=z_2$; then either $u_1=u_2$ and $u_0,\ u_3$ are functionally independent, or $u_0=u_3$ and $u_1,\ u_2$ are functionally independent; in the second case the n^{th} soliton u_{2^n-1} degenerates to the $(n-2)^{\text{th}}$ soliton u_{2^n-4} and its formula is thus valid; optionally one can apply a symmetry of the n dimensional cube and exchange the functionally independent $u_1,\ u_2$ with $u_0,\ u_3$ and thus still get $u_1=u_2$.

For the other quadrics we cannot expect Vandermonde determinants since z_k do not appear polynomially, but some of the behavior of the previous determinants remains; in particular since $\frac{\Delta'^+(z_{n+1},u_0,u_{2^n})}{4\mathcal{A}}=(u_{2^n}-u_0)^2-\frac{1}{4}z_{n+1}^2$ the use of (3.41) and backward induction by step two naturally appears.

The algebraic relation

$$\phi_n(\hat{v}_{2^n},\bigwedge_{k=1}^n(-\frac{\phi_2(\hat{v}_0,\bigwedge_{j=k,n+1}(\hat{v}_{2^{j-1}},z_j),0)}{\partial_v\phi_2(\hat{v}_0,\bigwedge_{j=k,n+1}(\hat{v}_{2^{j-1}},z_j),v)},z_k),\hat{v}_{2^{n+1}-1})\prod_{k=1}^n\partial_v\phi_2(\hat{v}_0,\bigwedge_{j=k,n+1}(\hat{v}_{2^{j-1}},z_j),v)=$$

$$t_{n+1}(\frac{\hat{\Delta}(z_{n+1},\hat{v}_0,\hat{v}_{2^n})}{z_{n+1}\mathcal{A}})^{[\frac{n}{2}]}\phi_{n+1}(\hat{v}_0,\bigwedge_{k=1}^{n+1}(\hat{v}_{2^{k-1}},z_k),\hat{v}_{2^{n+1}-1}),\ n\geq 2 \tag{3.42}$$

is valid for independent variables $\hat{v}_0,\ \{\hat{v}_{2^{k-1}}\}_{k=1,...,n+1},\ \hat{v}_{2^{n+1}-1}$ and certain constants
$t_n=t_n(z_1,...,z_n)$ ($\hat{\Delta}$ is one of $\Delta^{\pm},\ \Delta'^{\pm}$ according to the choice of the rulings $\hat{v}_0,\ \hat{v}_{2^n}$ and thus the use of (3.41) becomes clear). Since $\hat{v}_{2^n+2^{k-1}}=-\frac{\phi_2(\hat{v}_0,\bigwedge_{j=k,n+1}(\hat{v}_{2^{j-1}},z_j),0)}{\partial_v\phi_2(\hat{v}_0,\bigwedge_{j=k,n+1}(\hat{v}_{2^{j-1}},z_j),v)}$ (more generally for $j:=\Sigma_{s=1}^p 2^{k_s-1},\ 0<k_1<...<k_p\leq n$ we have $\phi_p(\hat{v}_0,\bigwedge_{s=1}^p(\hat{v}_{2^{k_s-1}},z_{k_s}),\hat{v}_j)=0$), (3.42) gives a procedure of generating ϕ_n by induction after n.

From a practical point of view (generate ϕ_n) we don't need to completely prove (3.42): note that the lhs of (3.42) and $\phi_{n+1}(...)$ must be simultaneously annihilated; thus (3.42) must be true with the coefficient of $\phi_{n+1}(...)$ being an a-priori arbitrary polynomial in $\hat{v}_0,\ \hat{v}_{2^n}$ (the remaining variables $\{\hat{v}_{2^{k-1}}\}_{k=1,...,n+1},\ \hat{v}_{2^{n+1}-1}$ appear linearly, since we know that ϕ_{n+1} must be a homography). Using few terms of the highest degree of this polynomial (with the simplification $\hat{v}_0=0$ for

n even, since $\hat{v}_{2^{n+1}-1}$ does not depend on $\hat{v}_0$) and the required skew-symmetryies of $\phi_{n+1}(...)$ wrt $\{(\hat{v}_{2^{k-1}}, z_k)\}_{k=1,...,n+1}$, one can find $\phi_{n+1}(...)$, t_{n+1} and the considered terms (which turn out to be as they should).

By this practical procedure we first find ϕ_n for $n = 3, 4$ and assume it to be found for $n \geq 5$; then (3.42) is proven by backward induction after n with step 2 (the validity of (3.42) for n is reduced to the validity of (3.42) for $n-2$).

For ϕ_3 again there is a formulation for all QC:

$$X(\hat{v}_1, \hat{v}_7)^T (M_3(z_1 \wedge z_2) - M_3(z_1 \wedge z_3))(M_3(z_1 \wedge z_2) + M_3(z_1 \wedge z_3))^{-1} X(\hat{v}_2, \hat{v}_4) = 1;$$

one replaces $\sqrt{R_{z_k}}$ with $-\sqrt{R_{z_k}}$ in M_3 if $\hat{v}_0,\ \hat{v}_{2^{k-1}}$ have different signatures.

We have:

$$\phi_3(\bigwedge_{k=1}^{3} (\hat{v}_{2^{k-1}}, z_k), \hat{v}_7) = \sum_{k=1}^{3} (-1)^{k-1} \phi_2(\hat{v}_7, \bigwedge_{j=1,\ j\neq k}^{3} (\hat{v}_{2^{j-1}}, z_j), \hat{v}_{2^{k-1}}). \tag{3.43}$$

Note that ϕ_3 for rulings of same signature has the most symmetries (the symmetries of a regular tetrahedron), since all rulings are on equal footing (for example for ϕ_2 the relationship between v_0, v_1 is different from that between v_0, v_3).

To prove (3.42) for $n = 2,\ t_3 = -\prod_{k=1}^{3} \frac{1}{m_2(z_k)},\ \hat{v}_0 = 0$ we need

$$I(\bigwedge_{k=1}^{3} z_k) :\ \sum_{k=1}^{3} (-1)^{k-1} \frac{m_2(z_k)}{m_3(\bigwedge_{j=1, j\neq k}^{3} z_j)} = \prod_{k=1}^{3} \frac{m_2(z_k)}{m_3(\bigwedge_{j=1, j\neq k}^{3} z_j)},$$

$$II(\bigwedge_{k=1}^{3} z_k) :\ \sum_{k=1}^{3} (-1)^{k-1} \frac{\prod_{j=1, j\neq k}^{3} m_2(z_j)}{m_3(\bigwedge_{j=1, j\neq k}^{3} z_j)} = 4a^2\delta \prod_{k=1}^{3} \frac{1}{m_3(\bigwedge_{j=1, j\neq k}^{3} z_j)};$$

again we maintain the $\wedge$ notation for m_3, although sometimes m_3 loses the skew-symmetry and we skip the $-$ when it appears in $m_2^-,\ m_3^-$. For all (I)Q(W)C except the general QC (3.3) (A^{-1} having distinct eigenvalues) and all rulings one can provide a justification of (3.43) without using (3.42) (and thus I& II are consequences and do not need proof): from (3.33)-(3.40),(3.43) we have $\sum_{k=1}^{3} (-1)^{k-1} (\phi_2(\hat{v}_0, \bigwedge_{j=1,\ j\neq k}^{3} (\hat{v}_{2^{j-1}}, z_j), \hat{v}_{7-2^{k-1}}) + \phi_2(\hat{v}_7, \bigwedge_{j=1,\ j\neq k}^{3} (\hat{v}_{7-2^{j-1}}, z_j), \hat{v}_{2^{k-1}})) = \phi_3(\bigwedge_{k=1}^{3} (\hat{v}_{2^{k-1}}, z_k), \hat{v}_7) + \phi_3(\bigwedge_{k=1}^{3} (\hat{v}_{7-2^{k-1}}, z_k), \hat{v}_0)$. By the same argument as when was proven that $\hat{v}_7$ does not depend on $\hat{v}_0$, one can keep $\hat{v}_1, \hat{v}_2, \hat{v}_4, \hat{v}_7$ constant and vary $\hat{v}_0, \hat{v}_3, \hat{v}_5, \hat{v}_6$ and conversely; thus $\phi_3(\bigwedge_{k=1}^{3} (\hat{v}_{2^{k-1}}, z_k), \hat{v}_7) = c(z_1, z_2, z_3)$; by symmetry $\phi_3(\bigwedge_{k=1}^{3} (\hat{v}_{7-2^{k-1}}, z_k), \hat{v}_0) = c(z_1, z_2, z_3)$, so $c(z_1, z_2, z_3) = 0$.

The reader familiar with the proof of the BPT for the B transformation of the sine-Gordon equation (see Bianchi ([5],Vol **5**,(46))) may recall similar cancellations and rightfully so: we shall see later that these algebraic computations encode all necessary information for the proof of the BPT.

Because ϕ_2 for QWC (3.7), IQWC (3.9) are missing some quadratic terms for mixed rulings, (3.43) and the above argument have to be slightly modified. For example:

$$\begin{aligned}
&\phi_3^4((v_1, z_1) \wedge (v_2, z_2), (u_4, z_3), u_7) = \phi_2^2(u_7, (v_2, z_2), (u_4, z_3), v_1) - \phi_2^2(u_7, (v_1, z_1), (u_4, z_3), v_2) + \\
&\qquad 2\phi_2^0(u_7, (v_1, z_1) \wedge (v_2, z_2), 0) \text{ for QWC (3.7)}, \\
&\phi_3^4((v_1, z_1) \wedge (v_2, z_2), (u_4, z_3), u_7) = \phi_2^2(v_1, (v_2, z_2), (u_4, z_3), u_7) - \phi_2^2(v_2, (v_1, z_1), (u_4, z_3), u_7) + \\
&\qquad \phi_2^7(0, (0, z_1) \wedge (0, z_2), 0) - \phi_2^7(u_7, (v_1, z_3) \wedge (v_2, z_3), u_4) \text{ for IQWC (3.9)};
\end{aligned} \tag{3.44}$$

further $\phi_2^2(v_0, (v_2, z_2), (u_4, z_3), u_6) - \phi_2^2(v_0, (v_1, z_1), (u_4, z_3), u_5) + \phi_2^2(v_3, (v_1, z_2), (u_7, z_3), u_5) - \phi_2^2(v_3, (v_2, z_1), (u_7, z_3), u_6) + 2\phi_2^0(v_0, (v_1, z_1) \wedge (v_2, z_2), v_3) = \phi_3^4((v_1, z_1) \wedge (v_2, z_2), (u_4, z_3), u_7) + \phi_3^4((v_0, z_1) \wedge (v_3, z_2), (u_5, z_3), u_6)$ for QWC (3.7); for IQWC (3.9) replace $\phi_2^0(v_0, (v_1, z_1) \wedge (v_2, z_2), v_3)$ with $\phi_2^7(u_7, (u_6, z_1) \wedge (u_5, z_2), u_4)$. We don't need (3.44) to find ϕ_3^4: because of the symmetries of the 3-dimensional cube we have $\phi_3^4((v_1, z_1) \wedge (v_2, z_2), (u_4, z_3), u_7) = \phi_3^6((v_1, z_3), (u_7, z_2) \wedge (u_4, z_1), v_2)$; for ϕ_3^6 (3.43) and the above argument remain valid without modification. Note also

$\sum_{k=1}^{3}(-1)^{k-1}(\phi_2^3(v_0,\bigwedge_{j=1,\ j\neq k}^{3}(u_{2^{j-1}},z_j),v_{7-2^{k-1}})+\phi_2^4(u_7,\bigwedge_{j=1,\ j\neq k}^{3}(v_{7-2^{j-1}},z_j),u_{2^{k-1}}))=$
$-2\phi_3^0(\bigwedge_{k=1}^{3}(v_{7-2^{k-1}},z_k),v_0)$ for QWC (3.7), $=-2\phi_3^{15}(\bigwedge_{k=1}^{3}(u_{2^{k-1}},z_k),u_7)$ for IQWC (3.9). The $-$ sign appears in the rhs (for all quadrics) because $m_3(z_{k_1}\wedge z_{k_2})$ acquires a $-$ sign for $\phi_2^3,\ \phi_2^4$.

We have $t_n=-\frac{1}{m_2(z_n)^{n-2}}\prod_{k=1}^{n-1}\frac{1}{m_2(z_k)}$; to get ϕ_4 from (3.42) for $n=3$ as previously described we need the consequences rhs $\mathrm{II}(\bigwedge_{k=1}^{3}z_k)\prod_{k=1}^{3}\frac{1}{m_3(\bigwedge_{j=1,j\neq k}^{3}z_j)}=\sum_{k=1}^{3}\frac{(-1)^{k-1}\mathrm{lhs\ I}(\bigwedge_{j=1,j\neq k}^{4}z_j)}{m_2(z_4)m_3(\bigwedge_{j=k,4}z_j)}(=\sum_{k=1}^{3}\frac{(-1)^{k-1}}{m_3(\bigwedge_{j=1,\ j\neq k}^{3}z_j)m_3(\bigwedge_{j=k,4}z_j)})$, $\sum_{k=1}^{4}\frac{(-1)^{k-1}\mathrm{rhs\ I}(\bigwedge_{j=1,j\neq k}^{4}z_j)}{\prod_{j=1,\ j\neq k}^{4}m_2(z_j)}=\sum_{k=1}^{4}\frac{(-1)^{k-1}\mathrm{lhs\ I}(\bigwedge_{j=1,j\neq k}^{4}z_j)}{\prod_{j=1,\ j\neq k}^{4}m_2(z_j)}$ $(=0)$ of I,II; in fact one can get a whole hierarchy of consequences of I,II for $n\geq 5$.

3.7. **Doubly ruled Möbius configurations.**

We consider *doubly ruled Möbius configurations* (DRMC) as a natural generalization of Möbius configurations, apparent at the level of analytic computations. Let $k=1,2,3$; consider rulings $\hat{v}_0(=u_0$ or $v_0),\hat{v}_{2^{k-1}}(=u_{2^{k-1}}$ or $v_{2^{k-1}})\in\mathbb{C}$, signatures $\epsilon_0(=1$ if $\hat{v}_0=v_0$ and -1 otherwise), $\epsilon_{2^{k-1}}(=1$ if $\hat{v}_{2^{k-1}}=v_{2^{k-1}}$ and -1 otherwise) associated to these rulings and $z_k\in\mathbb{C}$. The homographies below depend a-priori in an arbitrary fashion on z_k.

Assume:

I The 1st *DRMC* $\mathcal{M}_1$ *is a homography* $\phi_1((u_0,v_0)\odot(u_1,v_1),z_1)=0$.

II Given $\phi_1((u_0,v_0)\odot(u_1,v_1),z_1)=0,\ \phi_1((u_0,v_0)\odot(u_2,v_2),z_2)=0$, *there are two choices of* (u_3,v_3) *(according to* $\epsilon_1\epsilon_2=\pm1$*) with* $\phi_1((u_3,v_3)\odot(u_1,v_1),z_2)=0,\ \phi_1((u_3,v_3)\odot(u_2,v_2),z_1)=0$.

The 2nd DRMC $\mathcal{M}_2$ is a homography $\phi_2(\hat{v}_0,(\hat{v}_1,z_1),(\hat{v}_2,z_2),\hat{v}_3)=0,\ \epsilon_3=\epsilon_0\epsilon_1\epsilon_2$. Moreover symmetries of the square 0123 below which do not induce symmetries of ϕ_2 generate valid versions of ϕ_2 (for simplicity we drop in the notation ϕ_2 the dependence on $\epsilon_0,\epsilon_1,\epsilon_2$).

III Given $\phi_2(\hat{v}_0,(\hat{v}_2,z_2),(\hat{v}_4,z_3),\hat{v}_6)=0,\ \phi_2(\hat{v}_0,(\hat{v}_1,z_1),(\hat{v}_4,z_3),\hat{v}_5)=0,$
$\phi_2(\hat{v}_0,(\hat{v}_1,z_1),(\hat{v}_2,z_2),\hat{v}_3)=0,\ \epsilon_{2^{k_1-1}+2^{k_2-1}}=\epsilon_0\epsilon_{2^{k_1-1}}\epsilon_{2^{k_2-1}}$, *there is a unique* $\hat{v}_7,\ \epsilon_7=\epsilon_1\epsilon_2\epsilon_4$ *with* $\phi_2(\hat{v}_1,(\hat{v}_3,z_2),(\hat{v}_5,z_3),\hat{v}_7)=0,\ \phi_2(\hat{v}_2,(\hat{v}_3,z_1),(\hat{v}_6,z_3),\hat{v}_7)=0,\ \phi_2(\hat{v}_4,(\hat{v}_5,z_1),(\hat{v}_6,z_2),\hat{v}_7)=0$.

The 3rd DRMC $\mathcal{M}_3$ is a homography $\phi_3((\hat{v}_1,z_1),(\hat{v}_2,z_2),(\hat{v}_4,z_3),\hat{v}_7)=0,\ \epsilon_7=\epsilon_1\epsilon_2\epsilon_4$. Moreover symmetries of the cube below which do not induce symmetries of ϕ_3 generate valid versions of ϕ_3.



As in the previous subsection these three assumptions allows us to generate the n^{th} DRMC $\mathcal{M}_n$ as a homography $\phi_n(\hat{v}_0,(\hat{v}_1,z_1),...,(\hat{v}_{2^n-1},z_n),\hat{v}_{2^n-1})=0$. Moreover symmetries of the n-dimensional cube which do not induce symmetries of ϕ_n generate valid versions of ϕ_n.

Thus the natural question arises if all DRMC can be reduced, via Möbius transformations $u\mapsto\frac{au+b}{cu+d}$ (applied to all rulings), to one of the cases of the previous subsection.

In particular, is the relation $\phi_3((\hat{v}_1,z_1),(\hat{v}_2,z_2),(\hat{v}_4,z_3),\hat{v}_7)=\phi_2(\hat{v}_7,(\hat{v}_2,z_2),(\hat{v}_4,z_3),\hat{v}_1)-\phi_2(\hat{v}_7,(\hat{v}_1,z_1),(\hat{v}_4,z_3),\hat{v}_2)+\phi_2(\hat{v}_7,(\hat{v}_1,z_1),(\hat{v}_2,z_2),\hat{v}_4)$ still valid (after proper normalization of ϕ_2 and with some modifications in exceptional cases)?

4. The rolling problem

4.1. **Submanifolds in $\mathbb{C}^n$.**

We shall recall without many details the concept of sub-manifolds of $\mathbb{C}^n$ (called henceforth sub-manifolds). In the early development of the geometry of sub-manifolds in scalar product spaces geometers made liberal use of imaginary numbers, until this geometry has been divided into Lorentz and Riemannian.

We would like to have the GW and the GCMR equations for a sub-manifold x in $\mathbb{C}^n$ just as the GW and GCMR equations for a real sub-manifold in $\mathbb{R}^n$. The important thing in creating these formulae is the existence of the normal bundle; thus the natural definition for k-dimensional sub-manifolds is that they have k-dimensional complex tangent spaces (called henceforth tangent spaces), $1 \leq k \leq n-1$ and thus $(n-k)$-dimensional normal spaces. We exclude the case of isotropic tangent spaces (spaces having tangency with the isotropic cone), when the linear element degenerates and one obtains among others sub-Riemannian manifolds (the linear element is real and positive semi-definite). The real dimension of the real tangent spaces is a-priori any number between k and $2k$ (we exclude the case when the dimension of the real tangent spaces varies while the dimension of the complex tangent spaces remains constant). The distribution $T \cap iT$ formed by intersecting real tangent spaces T with iT is Frobenius integrable (by its own definition); on its leaves the Nijenhuis condition holds, the complex structure being induced by the surrounding space; from Nirenberg-Newlander the leaves are holomorphic sub-manifolds. The Frobenius theorem gives a parametrization holomorphic on these leaves; in this parametrization it appears clearly that the manifold is essentially k-dimensional, although apparently initially could have had real dimension between k and $2k$. If it so happens that after possibly applying a change of variables the parametrization in some of the remaining real parameters reveals itself to be real analytic, it can be extended by analyticity to a k-dimensional sub-manifold with more complex parameters (and less real).

Take for example the IQWC (3.9) $x_0 = uvf_1 + u\bar{f}_1 + ve_3$ and let $x_0^0 \subseteq x_0$ be a surface of real dimension 3 obtained by restricting (u, v) to the domain $(u, v) = (x + iy, s + ixy),\ x, y, s \in \mathbb{R}$; this is chosen so that $K(x_0^0) = \frac{1}{(u^2-2v)^2} = \frac{1}{(x^2-y^2-2s)^2} > 0$. We know that there is a parametrization $(z, t),\ z \in \mathbb{C},\ t \in \mathbb{R}$ of x_0^0; since in this case $K(x_0^0)$ is holomorphic in z while real valued, we get $K(x_0^0) = f(t)$; we can choose $t := u^2 - 2v = x^2 - y^2 - 2s$ and $x_0^0 = u\frac{u^2-t}{2}f_1 + u\bar{f}_1 + \frac{u^2-t}{2}e_3$, so we can choose $z := u$. In this case $T \cap iT = \mathbb{C}x_{0z}^0 = \mathbb{C}(x_{0u}^0 + ux_{0v}^0)$ is real 2-dimensional.

We need the normal bundle (and the fundamental forms of the sub-manifold) to be holomorphic in the complex parameters. This is done as follows: along the holomorphic leaves the tangent spaces to the sub-manifold are spanned by the derivatives of the position vector in the complex parameters and in the real parameters, but because along such leaves the real parameters are set to constants, the vectors spanning the tangent spaces are holomorphic in the complex parameters, so we can find in the normal spaces orthonormal frames which vary holomorphically in the complex parameters. Thus the linear element and the second fundamental form are holomorphic in the complex parameters and the sub-manifold is recovered (the Gauß-Bonnet theorem) modulo rigid motions of $\mathbb{C}^n$ from the knowledge of its linear element and the second fundamental form through the integration of a Ricatti equation (the GW equations) and a quadrature.

For two applicable sub-manifolds of $\mathbb{C}^n$ one can choose orthonormal frames for both normal bundles and define the rotation of the rolling as the family of orthogonal transformations which takes the bases of the tangent spaces formed by parametric vectors and the orthonormal frames for the normal bundles at corresponding points one onto the other. The rotation can be arbitrarily chosen so as to take any orthonormal frame of the first normal bundle into any other of the second normal bundle, so we take advantage and define it to take a holomorphic (in the complex parameters) orthonormal frame into a holomorphic one. In this way the rotation is holomorphic in the complex parameters (and unique modulo right transformations through sections of $\mathbf{O}_{n-k}(\mathbb{C})$ bundles holomorphic in the complex parameters) and so will turn out to be the translation.

For example if $x(u, v)$ is a holomorphic surface and we impose a nonanalytic condition $f(u, \bar{u}) = 0$ or explicitly $u(t) = g(t),\ t \in \mathbb{R}$, then the resulting surface $y(t, v) := x(u(t), v)$ is nonanalytic in t, but can be extended to the holomorphic surface $x(u, v)$.

Ruled surfaces with real ruling or real surfaces of revolution extend by analyticity to ruled surfaces or surfaces of revolution.

However if we take for example a holomorphic ruled surface $x(u, v) := a(u)v + b(u)$, non-analytically restrict the complex variable u to a curve $u(t)$ and then one deforms the resulting

$y(t,v) := x(u(t),v)$ (and analytically inconsistent with $u(t)$'s non-analyticity) to a ruled surface $z(t,v)$ with the same ruling, then $z(t,v)$ cannot be extended to a holomorphic ruled surface applicable to the initial one.

Intersecting such a k-dimensional sub-manifold of $\mathbb{C}^n$ with a totally real subspace (here we use the definition of a totally real space as being obtained after a rigid motion of $\mathbb{C}^n$ from an n-dimensional real subspace of $\mathbb{C}^n$ having real or purely imaginary entries in the standard basis; all definitions of totally real subspaces coincide) we get a real sub-manifold of a Lorentz space (the imaginary part of the scalar product in $\mathbb{C}^n$ disappears), possibly analytic in some parameters.

A real analytic k-dimensional sub-manifold of $\mathbb{R}^n$ extends by analyticity to a holomorphic k-dimensional sub-manifold of $\mathbb{C}^n$. One can iterate this procedure: a holomorphic k-dimensional sub-manifold in $\mathbb{C}^n$ can be considered as a real analytic $2k$-dimensional sub-manifold of $\mathbb{R}^{2n}$ which can be complexified to a holomorphic $2k$-dimensional sub-manifold in $\mathbb{C}^{2n}$, but the Euclidean scalar product on $\mathbb{C}^n$ becomes a pair of indefinite scalar products on $\mathbb{C}^{2n}$, so it is not interesting from a metric point of view.

CMC(minimal) surfaces have analytic data in conformal parametrization, so they extend to holomorphic CMC(minimal) surfaces. The Lawson or Bianchi-Bryant-Weierstrass isometric correspondence between minimal surfaces and CMC surfaces in space forms always has as domain a space form of smaller curvature; for the domain to be a space form of higher curvature we need time-like mean curvature vector, so we need deformations in Lorentz spaces.

According to Bianchi ([5],Vol **4**,(202)), Lie noticed that the formula developed by Monge for minimal surfaces can be stated as: all minimal surfaces are surfaces of translation with isotropic generating curves, that is $x = f(z) + \bar{f}(\bar{z})$, $f_z^T f_z = 0$ and this is one step from the Weierstrass formula, only that at that time Cauchy had not yet rigorously developed the theory of holomorphic functions.

Bianchi found in ([5],Vol **4**,(108)) a conformal correspondence between CMC surfaces in $\mathbb{R}^3$ and CMC surfaces in $\mathbb{S}^3$, ($\mathbb{H}^3$): if one replaces the Euclidean metric on a ball containing two parallel CMC surfaces (the existence of this configuration was proven by Bonnet) with the metric induced by considering the boundary of the given ball as the Cayley's absolute, we obtain two parallel surfaces in $\mathbb{S}^3$, ($\mathbb{H}^3$) (that is they are normal to a normal congruence of geodesics) in conformal correspondence (the conformal correspondence from the Euclidean space is preserved), so they must have CMC.

While this conformal correspondence is not of applicability, it is good enough to study the topology of these surfaces; for example CMC tori (Wente, Pinkall- Sterling [25], Bobenko's CMC tori in terms of θ-functions) or CMC compact surfaces of higher genus (Kapouleas), including the case when these surfaces are tangent to the boundary of the ball in discussion or when they have singularities (for example cusps or degenerate linear element) at this boundary.

4.2. **Rolling surfaces.**

The study of the rolling problem was initiated by Ribacour and has been extensively pursued in Bianchi [4],([5],Vol **7**) and Darboux [13].

Let $(u^1,u^2) \ni D$ with D a domain in $\mathbb{R}^2$, $\mathbb{C}\times\mathbb{R}$ or $\mathbb{C}^2$ and $x : D \longmapsto \mathbb{C}^3$ be a surface. Then $|dx|^2 = dx^T dx = g_{jk}du^j \otimes du^k$ is the first fundamental form. Since $dx^T \wedge dx = 0$, $|dx|^2$ is a symmetric $(0,2)$ tensor. We have $d^2x^T dx = \Gamma_{jk,l}du^j \otimes du^k \otimes du^l$, where $\Gamma_{jk,l}$ are the Christoffel symbols of first type. Since d^2x is symmetric we have $\Gamma_{jk,l} = \Gamma_{kj,l}$. Denote $\partial x^2 := g^{jk}\partial_{u^j} \otimes \partial_{u^k}$, so $\partial x^2 \lrcorner |dx|^2 = |dx|^2 \llcorner \partial x^2 = \partial_{u^j} \otimes du^j$ is the identity element of both $\lrcorner$, $\llcorner$ (with a,b tensors by $a \lrcorner b = b \llcorner a$ we contract the first contravariant entry of a with the first covariant entry of b; one can stack the contractions in wait for the contravariant or covariant entries without problems when working with symmetric tensors).

For a function $f : D \longmapsto \mathbb{C}$ we have $\nabla f := df \llcorner \partial x^2 = f_{u^j} g^{jk}\partial_{u^k}$ the tensor dual to df, so we have $\Gamma^l_{jk}\partial_{u^l} \otimes du^j \otimes du^k = \nabla x^T d^2x$. Denote $\nabla^2 f := d^2 f - df \llcorner \nabla x^T d^2x$ the second covariant derivative of f. We have $0 = d(|dx|^2 \llcorner \partial x^2)$ so $d(\partial x^2) = -\partial x^2 \lrcorner d(|dx|^2) \llcorner \partial x^2 = -\partial x^2 \lrcorner d^2x^T \nabla x - \nabla x^T d^2x \llcorner \partial x^2$.

If N is a unit normal frame of the surface, then the second fundamental form is $-dx^T dN = d^2x^T N$ and the GW equations are: $d^2x = (d^2x)^\top + (d^2x)^\perp = dx\llcorner\nabla x^T d^2x + NN^T d^2x$, or equivalently: $\nabla^2 x = NN^T d^2x$, $dN = (dN)^\top = dx\llcorner\nabla x^T dN = -\nabla x\lrcorner d^2x^T N$.

For $\omega_1,\ \omega_2$ vector 1 forms in $\mathbb{C}^3$ and $a, b \in \mathbb{C}^3$, we have $a^T\omega_1\wedge b^T\omega_2 = ((a\times b)\times\omega_1 + b^T\omega_1 a)^T\wedge\omega_2 = (a\times b)^T\omega_1\times\wedge\omega_2 + b^T\omega_1\wedge a^T\omega_2$. In particular $a^T\omega\wedge b^T\omega = \frac{1}{2}(a\times b)^T\omega\times\wedge\omega$. Since both $\times$ and $\wedge$ are skew-symmetric, we have $\omega_1\times\wedge\omega_2 = \omega_1\times\omega_2 + \omega_2\times\omega_1 = \omega_2\times\wedge\omega_1$.

Consider the scalar product $<,>$ on $\mathbf{M}_3(\mathbb{C})$: $<X, Y> := \frac{1}{2}\mathrm{tr}(X^T Y)$. We have the isometry

$$\alpha : \mathbb{C}^3 \to \mathbf{o}_3(\mathbb{C}),\ \alpha(\begin{bmatrix} x^1 \\ x^2 \\ x^3 \end{bmatrix}) = \begin{bmatrix} 0 & -x^3 & x^2 \\ x^3 & 0 & -x^1 \\ -x^2 & x^1 & 0 \end{bmatrix},\ x^T y =< \alpha(x), \alpha(y) >= \tfrac{1}{2}\mathrm{tr}(\alpha(x)^T\alpha(y)),$$

$\alpha(x\times y) = [\alpha(x), \alpha(y)] = \alpha(\alpha(x)y)$.

Note $< \alpha(x)^2, \alpha(y) >= \frac{1}{2}\mathrm{tr}((\alpha(x)^2)^T\alpha(y))) = -\frac{1}{2}\mathrm{tr}(\alpha(y)^T\alpha(x)^2) = 0$.

Let $x \subset \mathbb{C}^3$ be a surface non-rigidly applicable to a surface $x_0 \subset \mathbb{C}^3$:

$$(x, dx) = (R, t)(x_0, dx_0) := (Rx_0 + t, Rdx_0), \tag{4.1}$$

where (R, t) is sub-manifold in $\mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ (in general surface, but it is a curve if $x_0,\ x$ are ruled and the rulings correspond under the applicability). The sub-manifold R gives the rolling of x_0 on x, that is if we rigidly roll x_0 on x such that points corresponding under the applicability will have the same differentials, R will dictate the rotation of x_0; the translation t will satisfy $dt = -dRx_0$.

Therefore $\mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ acts on 2-dimensional integrable distributions of facets (x_0, dx_0) in $T^*\mathbb{C}^3$ as: $(R, t)(x_0, dx_0) = (Rx_0 + t, Rdx_0)$ a rolling is a sub-manifold $(R, t) \in \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$ such that $(R, t)(x_0, dx_0)$ is still integrable.

Consider first the case $[R^{-1}dR, \wedge R^{-1}dR] \neq 0$ (R is a surface).

Applying the compatibility condition $d\wedge$ to (4.1) we get:

$$R^{-1}dR \wedge dx_0 = 0 = dRR^{-1} \wedge dx. \tag{4.2}$$

Let b be an right infinitesimal deformation of R:

$R_\epsilon := R\exp(\epsilon b)$, $\frac{d}{d\epsilon}|_{\epsilon=0}(R_\epsilon^{-1}dR_\epsilon) = db - [b, R^{-1}dR]$, $0 = \frac{d}{d\epsilon}|_{\epsilon=0} < R_\epsilon^{-1}dR_\epsilon, R_\epsilon^{-1}dR_\epsilon >=< R^{-1}dR, db > + < db, R^{-1}dR >$, so $db = [R^{-1}dR, \alpha(y)]$ for some surface y in $\mathbb{C}^3$, or $d\alpha^{-1}(b) = R^{-1}dRy$. Imposing the compatibility condition $d\wedge$ we get $R^{-1}dR\wedge dx_0 = 0$ for $x_0 := y + \alpha^{-1}(b)$ and this is just (4.2). Thus any infinitesimal deformation of a surface R is put into correspondence with pairs (x_0, x) of applicable surfaces in the Euclidean space $\mathbb{C}^3$ (this is essentially due to Bianchi; see ([4],§ 494*)). Note that $t := -R\alpha^{-1}(b)$ will be the translation of the rolling of x_0 on $x := Ry$, $dx = Rdx_0$. Also $R^{-1}N_R := \frac{[R^{-1}dR, R^{-1}dR]}{|[R^{-1}dR, R^{-1}dR]|}$ (the left Gauß map of R) is parallel to $\alpha(N_0)$, $N_0 := \frac{dx_0\times\wedge dx_0}{|dx_0\times\wedge dx_0|}$ and $N_R R^{-1}$ to $\alpha(N := RN_0)$, so any statement about $R,\ x_0$ becomes a statement about $R^{-1},\ x$. This dichotomy into left and right geometries is due to Clifford; he got interesting geometric results (Clifford left and right translations) of what is basically currently know as $\mathbf{o}_4(\mathbb{C}) = \mathbf{o}_3(\mathbb{C}) \oplus \mathbf{o}_3(\mathbb{C})$; this can be lifted for all practical purposes (locally) to the Lie groups and thus provided a tool to attack the geometry of 3-dimensional space forms. Sym's formula from [30] basically uses a combination of both right and left differentiation: $d(R^{-1}\partial_z R) = R^{-1}\partial_z(dRR^{-1})R$ for $R = R(u, v, z)$, $d\cdot := (\partial_u\cdot)du + (\partial_v\cdot)dv$ when dRR^{-1} is linear in z (or more generally rational in z); most surfaces of interest to the classical geometers and arising as integrable systems (CGC, CMC) come in spectral families (Lie's spectral family for CGC -1 and Bonnet's family for CMC); for such families dRR^{-1} is linear in z and the family of surfaces is $R^{-1}\partial_z R \subset \mathbf{o}_3(\mathbb{R}) \simeq \mathbb{R}^3$.

Note that any two infinitesimal isometries of R will give pairs of applicable surfaces such that the corresponding normals are parallel.

Bianchi calls two surfaces $x,\ y \subset \mathbb{C}^3$ associate if $dx\times\wedge dy = 0$ (this has to do, as expected, with infinitesimal deformations of both x and y- see § 4.5).

Applying $R^{-1}d$ to (4.1) we get

$$R^{-1}d^2x = R^{-1}dRdx_0 + d^2x_0. \tag{4.3}$$

Also since $N = RN_0$ we have:

$$R^{-1}dRN_0 = R^{-1}dN - dN_0. \tag{4.4}$$

Thus for a parametrization (u, v) on $(x_0,\ x)$ with the same off-diagonal second fundamental forms (equivalent to $R_v x_u = 0$; for example the common conjugate system on $(x_0,\ x)$ or u (v) asymptotic parameter on x_0 (x)) the axes of rotation of the rotation R along a parametric line are the tangents to x_0 along the other parametric line. Bianchi calls such coordinates kinematically conjugate. The rotation of the rolling with the other face of x_0 (or on the other face of x) is $R' := R(I - 2N_0N_0^T) = (I - 2NN^T)R$.

Since $R^{-1}dR$ is skew-symmetric and using (4.2) we have

$$dx_0^T R^{-1}dRdx_0 = 0. \tag{4.5}$$

For $a \in \mathbb{C}^3$ we have $R^{-1}dRa = R^{-1}dR(a^\perp + a^\top) = a^T N_0 R^{-1}dRN_0 - a^T R^{-1}dRN_0N_0 = \omega \times a,\ \omega := N_0 \times R^{-1}dRN_0 = (\det R)R^{-1}(N \times dN) - N_0 \times dN_0$. Thus $R^{-1}dR = \alpha(\omega)$ and

$$d \wedge \omega + \frac{1}{2}\omega \times \wedge\omega = 0,\ \omega \times \wedge dx_0 = 0. \tag{4.6}$$

Therefore any such flat connection form ω in the tangent bundle of x_0 is of the form $\omega = R^{-1}dR,\ R^{-1}dR \wedge dx_0 = 0$ and we get all surfaces x applicable to x_0 by integrating the Ricatti equation $R^{-1}dR = \omega$ and a quadrature for $dx = Rdx_0$. To show that $R^{-1}dR = \omega$ is a Ricatti equation consider $X(a, b) := R^{-1}c$, where c is a constant unit vector, so $0 = R^{-1}dc = \omega \times X + dX$. Multiplying on the left with $Y(a)^T,\ Y(b)^T$ we get $Y(a)^T\omega = 2ida,\ Y(b)^T\omega = 2idb$; from any 2 solutions a, b of the Ricatti equation $Y(a)^T\omega = 2ida$ we can find $X(a, b)$, then R is uniquely determined by the constraint $d(RX) = 0$ and the initial condition (since c depends only on two constants, one adds another constant for the rotations which preserve c). Note that the integrability condition is satisfied: $2id \wedge da = Y(a)^T d \wedge \omega + \frac{i}{2}Y'(a)^T\omega \wedge Y(a)^T\omega = Y(a)^T(d \wedge \omega + \frac{1}{2}\omega \times \wedge\omega) = 0$. This Ricatti equation will also characterize the B_∞ transformation of QC.

In this vein Darboux showed in ([13],§ 725) that the problem of finding all surfaces applicable to a given one is equivalent to finding all virtual asymptotic coordinates. If we infinitesimally know a surface (that is the first and second fundamental forms), then we need to integrate a Ricatti equation and a quadrature to find the surface (for example from the GW equations we know $F^{-1}dF$, where $F := [N\ \frac{x_u}{|x_u|}\ N \times \frac{x_u}{|x_u|}]$).

If $x = x(u^1, u^2)$ is a given surface and $(\tilde{u}^1, \tilde{u}^2)$ are virtual asymptotic coordinates, then keeping account of the change of Christoffel symbols $\frac{\partial u^l}{\partial \tilde{u}^c}\tilde{\Gamma}^c_{ab} = (\frac{\partial^2 u^l}{\partial \tilde{u}^a \partial \tilde{u}^b} + \frac{\partial u^j}{\partial \tilde{u}^a}\frac{\partial u^k}{\partial \tilde{u}^b}\Gamma^l_{jk})$ and of the Codazzi-Mainardi equations $d\log(K) = 4(\tilde{\Gamma}^2_{12}d\tilde{u}^1 + \tilde{\Gamma}^1_{12}d\tilde{u}^2)$, we get the equations of virtual asymptotic coordinates:

$$\frac{\partial^2 u^l}{\partial \tilde{u}^1 \partial \tilde{u}^2} + \frac{\partial u^j}{\partial \tilde{u}^1}\frac{\partial u^k}{\partial \tilde{u}^2}\Gamma^l_{jk} + \frac{1}{4}(\frac{\partial u^j}{\partial \tilde{u}^1}\frac{\partial u^l}{\partial \tilde{u}^2} + \frac{\partial u^l}{\partial \tilde{u}^1}\frac{\partial u^j}{\partial \tilde{u}^2})\frac{\partial \log(K)}{\partial u^j} = 0,\ l = 1, 2. \tag{4.7}$$

The virtual asymptotic coordinates may not admit level curves; for example real surfaces of positive Gauß curvature have imaginary asymptotic directions and thus do not admit integral curves if the surface is not analytic, but one can use in this case *isothermal-conjugate systems* (conformal parametrization of the second fundamental form); their level curves are integral curves of the real and imaginary parts of the asymptotic directions.

Let $|dx_0|^2 =: g_{jk}du^j \otimes du^k,\ g := \frac{\det(|dx_0|^2)}{(du^1 \wedge du^2)^2}$ and $s := s_{jk}du^j \otimes du^k := (x_{jk} - x_{0jk})du^j \otimes du^k := N_0^T(R^{-1}d^2x - d^2x_0)$ be the difference of the second fundamental forms of $x,\ x_0$. Note:

$$\sqrt{g}\omega = (s_{12}x_{0u^1} - s_{11}x_{0u^2})du^1 + (s_{22}x_{0u^1} - s_{21}x_{0u^2})du^2; \tag{4.8}$$

($s_{12} = s_{21}$ is equivalent here to $\omega \times \wedge dx_0 = 0$; $d \wedge \omega + \frac{1}{2}\omega \times \wedge\omega = 0$ encodes the difference of the GCM equations of x_0 and x).

From (4.4) we get the first and left second fundamental forms of R : $|R^{-1}dR|^2 = |\omega|^2 = |R^{-1}dRN_0|^2 = s \llcorner \partial x_0^2 \lrcorner s = s_{jk}g^{kl}s_{lm}du^j \otimes du^m,\ < R^{-1}N_R, d(R^{-1}dR) >= N_0^T d\omega = N_0^T(dN_0 \times$

$R^{-1}dRN_0) = (N_0 \times dN_0)^T(R^{-1}dN - dN_0) = \frac{1}{\sqrt{g}}(x_{01j}s_{2k} - x_{02j}s_{1k})du^j \otimes du^k$. Note that the left second fundamental form of R is not symmetric; its symmetric part is the second fundamental form of R and its antisymmetric part accounts for the curvature $\frac{1}{4}$ of $\mathbf{O}_3(\mathbb{C})$.

Thus the common conjugate system on (x_0, x) corresponds to asymptotic coordinates on R (Bianchi) and asymptotic coordinates on x_0 (x) give a conjugate system on R. This is true again for associated surfaces; in fact this property together with parallel normals is Bianchi's definition of associated surfaces.

Gauß theorem states that $dN \times \wedge dN = 2K(x)Nda$ (da is the area form). Because $R^{-1}N_R$ and N_0 are parallel, we thus get $2K(x_0)\sqrt{\det(|dx_0|^2)} = \pm 2K(R)\sqrt{\det(|R^{-1}dR|^2)}$, so rotations of rollings of flat surfaces in $\mathbb{C}^3$ are flat surfaces in $\mathbf{O}_3(\mathbb{C})$ (also surfaces associated to flat surfaces are flat).

We have the mean curvature $H(R) = -\frac{\sqrt{g}}{\det s}g^{jk}(x_{01j}s_{2k} - x_{02j}s_{1k})$ and $db = \alpha(\omega \times R^{-1}x)$, $\frac{d}{d\epsilon}|_{\epsilon=0}(R_\epsilon^{-1}dR_\epsilon) = \alpha(\omega \times x_0)$, $\frac{d}{d\epsilon}|_{\epsilon=0}(R_\epsilon^{-1}N_{R_\epsilon}) = \frac{[db,R^{-1}dR]+[R^{-1}dR,db]}{|[R^{-1}dR,R^{-1}dR]|} - [b, R^{-1}N_R] = \alpha(N_0 \times R^{-1}x - (x_0 - R^{-1}x) \times N_0) = \alpha(N_0 \times x_0)$, $\frac{d}{d\epsilon}|_{\epsilon=0} < R_\epsilon^{-1}N_{R_\epsilon}, d(R_\epsilon^{-1}dR_\epsilon) >= N_0^T(\omega \times dx_0) = s$.

Therefore $H(R_\epsilon) = H(R) + \epsilon\frac{g}{\det s}(H(x) - H(x_0)) + O(\epsilon^2)$, so infinitesimal deformations of an isothermic R preserving its mean curvature are put in correspondence with Bonnet pairs in $\mathbb{C}^3$ (Bianchi ([4],§ 494*)).

We are interested in the mean and Gauß curvatures $H(t)$, $K(t)$ of the surface t described by a point (which can be considered the origin) rigidly attached to x_0 in this rolling. We thus have $t = x - Rx_0$ and $R^{-1}dt = -R^{-1}dRx_0 = -x_0^T dx_0 \llcorner R^{-1}dR\nabla x_0 - x_0^T N_0 R^{-1}dRN_0 = -x_0^T N_0 R^{-1}dRN_0 + x_0^T dx_0 \llcorner \nabla x_0^T R^{-1}dRN_0N_0$, so $|dt|^2 = (s\llcorner\partial x_0^2 \lrcorner x_0^T dx_0)^2 + (x_0^T N_0)^2|s\llcorner\partial x_0^2 \lrcorner dx_0|^2 = s\llcorner B \lrcorner s$,
$B := (\nabla\frac{|x_0|^2}{2})^2 + (x_0^T N_0)^2\partial x_0^2 = B^{jk}\partial_{u^j} \otimes \partial_{u^k} = (x_0^T g^{jl}\partial_l x_0 x_0^T g^{km}\partial_m x_0 + (x_0^T N_0)^2 g^{jk})\partial_{u^j} \otimes \partial_{u^k}$. $N_t = \frac{1}{|x_0|}Rx_0$ is the normal to t and the second fundamental form of t is: $N_t^T d^2 t = \frac{1}{|x_0|}x_0^T(R^{-1}d^2Rx_0 - R^{-1}dRdx_0) = \frac{1}{|x_0|}(x_0^T(R^{-1}d^2R - d(R^{-1}dR))x_0 - x_0^T R^{-1}dRdx_0) = \frac{1}{|x_0|}(-|dt|^2 + x_0^T N_0 s)$.

Using a kinematically conjugate system on (x_0, x) we have: $2H(t) = \frac{-1}{|x_0|}(2 - \frac{x_0^T N_0 s_{jj} B^{jj}}{s_{11}s_{22}(B^{11}B^{22} - (B^{12})^2)})$, $K(t) = \frac{1}{|x_0|^2}(1 + \frac{x_0^T N_0(x_0^T N_0 - s_{jj}B^{jj})}{s_{11}s_{22}(B^{11}B^{22} - (B^{12})^2)})$. But $B^{11}B^{22} - (B^{12})^2 = g^{-1}(x_0^T N_0)^2|x_0|^2$ and using the Gauß equation for both x_0, x we get in general coordinates, with
$n := 2(H(t)|x_0| + 1)|x_0|^2 x_0^T N_0$, $q := (K(t)|x_0|^2 - 1)|x_0|^2 x_0^T N_0$:

$$(2K(x)n + x_{0jk}B^{jk}) - x_{jk}(n(-1)^{j+k}x_{0(j+1)(k+1)}g^{-1} + B^{jk}) = 0,$$
$$(2K(x)q - x_0^T N_0 - x_{0jk}B^{jk}) - x_{jk}(q(-1)^{j+k}x_{0(j+1)(k+1)}g^{-1} - B^{jk}) = 0. \tag{4.9}$$

We have $|d\frac{x_0}{|x_0|}|^2 = |dN_t|^2 + \frac{1}{|x_0|^2}|dt|^2 - \frac{2}{|x_0|}N_t^T d^2 t =: d\sigma^2$.

Therefore a surface t appears as the translation of a rolling if $d\sigma^2$ has curvature 1. This constitutes a second order PDE (linear in the second order terms) for $\frac{1}{|x_0|}$; once a solution is known, x_0 can be recovered by the integration of a Ricatti equation. Any surface t appears as the translation of a rolling in a fashion depending on two arbitrary functions of one variable and to the lines of curvature parametrization of t corresponds on x_0 coordinates appearing orthogonal from the origin (Bianchi ([4],§ 323)).

We are interested in totally real deformations x of x_0 : $\bar{x} = \epsilon x$, $\epsilon := \mathrm{diag}[\pm 1\ \pm 1\ \pm 1]$. A-priori x_0 does not have to be totally real; since $|dx|^2$ is real, so it must be $|dx_0|^2$. Consider $x_0 \subset \mathbb{C}^3$ surface with real linear element of signature $\epsilon' := \mathrm{diag}[\epsilon_1\ \epsilon_2]$; by Sylvester's it must have real dimension 2 choosing $v_j := \sqrt{\epsilon_j}e_j$, $j = 1, 2, 3$, $V := [v_1\ v_2\ v_3]$, (ϵ_3 is undetermined at this point), we have $dx_0 = RV\psi$, $R \subset \mathbf{O}_3(\mathbb{C})$ where ψ is $\mathbb{R}^3$ valued 1− form, $e_3^T\psi = 0$. If $dx_0 = R'V\psi'$ is a similar decomposition, then $R'^T R \subset \mathbf{O}_3(\mathbb{C})$, $R'^T Re_3 = \pm e_3$, $\overline{R'^T R} = \epsilon R'^T R\epsilon$, $\psi' = r\psi$, $r := \epsilon V R_1^T RV$ is a curve of real orthogonal transformations of $\mathbb{R}^3$ preserving the signature ϵ, $re_3 = \pm e_3$, so we can consider $r \subset \mathbf{O}_2(\mathbb{R})$ if $\det\epsilon' = 1$ and $r \subset \mathbf{O}_{1,1}(\mathbb{R})$ if $\det\epsilon' = -1$. We have $d\bar{x}_0 = \bar{R}\epsilon V\psi = \bar{R}\epsilon R^T dx_0$, so $\bar{R}\epsilon R^T$ (which does not change if R changes as above) is the rotation of the rolling of x_0 on $\bar{x}_0$. Note

that if $d\bar{x}_0 = Rdx_0,\ R \subset \mathbf{O}_3(\mathbb{C})$, then $\bar{R}^T = R$, so such surfaces x_0 are put in correspondence with special right infinitesimal deformations of surfaces $R \subset \mathbf{O}_3(\mathbb{C}),\ \bar{R}^T = R$: with $y := R^{-1}\bar{x}_0$ we have $d(R^{-1}\bar{y}) - R^{-1}d(Ry) = 0$; equivalently the special right infinitesimal deformation $b := \alpha(Ry - \bar{y})$ of R satisfies its requirement $db = [R^{-1}dR, \alpha(y)]$. If $d\bar{x}_0 = Rdx_0,\ R \in \mathbf{O}_3(\mathbb{C})$, then $\bar{x}_0 = (R,t)x_0$, (R,t) rigid motion with $\bar{R}^T = R,\ \bar{t} + \bar{R}t = 0$. Should x_0 be totally real, then there would be signature ϵ and rigid motion (R_0^T, t_0) such that $\overline{(R_0^T, t_0)x_0} = \epsilon((R_0^T, t_0)x_0)$; equivalently $R = \bar{R}_0\epsilon R_0^T,\ \bar{t}_0 - \epsilon t_0 + \bar{R}_0^T t = 0$; once we found R_0 with $R = \bar{R}_0\epsilon R_0^T,\ \bar{R}_0^T t$ will satisfy the compatibility condition $\overline{\bar{R}_0^T t} = -\epsilon\bar{R}_0^T t$ needed in order to prescribe t_0. Note that $R_0,\ t_0$ can be prescribed modulo rigid motions of the corresponding totally real space, which satisfy $\bar{R}_0 = \epsilon R_0\epsilon,\ \bar{t}_0 = \epsilon t_0$. Now the signature ϵ' can be found from the linear element of x_0, the last component ϵ_3 of the signature ϵ (so far undetermined) can be found from $\det R = \det\epsilon$. The usual argument with the dimension of kernel and codomain and connectedness of the codomain of the map $\mathbf{O}_3(\mathbb{C}) \ni R_0 \mapsto \bar{R}_0\epsilon R_0^T$ applies to prove the existence of R_0.

Thus all totally real surfaces x_0 are recognized from $d\bar{x}_0 = Rdx_0,\ R \in \mathbf{O}_3(\mathbb{C}),\ \bar{R}^T = R$.

Note that there are non-totally real surfaces with real linear element. To see this note that there are curves in arc-length parametrization (that is $c = c(s),\ s \in \mathbb{R},\ |dc|^2 = \pm ds^2$) which are not totally real. All real and real analytic 2-dimensional linear elements admit local isometric embedding in totally real spaces and depending on two functions of a variable as long as the signature of the ambient space allows the signature of the linear element. If one takes a real analytic curve in arc-length parametrization on the surface and one deforms it to a general position (non-totally real), then by Cauchy-Kovalewskaia a small portion of the surface around the curve can be deformed to the new general position: thus is non-totally real. In general Cauchy-Kovalewskaia fails for characteristic data either in non-existence or non-uniqueness of solutions, but in our case it fails in the good way: if the general position of the deformed curve is asymptote on the new surface, it is asymptote on an 1-dimensional family of deformations of the considered surface (Darboux proved that the characteristics of the deformation problem are the asymptotes; thus it becomes clear that Bianchi's notion of W congruence is the correct tool in studying the deformation problem since although it does not give in general the applicability correspondence, it gives the correspondence of characteristics). Here Eisenhart in ([19],§ 139) puts the condition that for real surfaces and deformations the curvature of the deformed curve should be greater than or equal to the geodesic curvature of the curve on the surface (equality is obtained when the deformed curve becomes asymptote on the deformed surface). For the same problem (real surface and deformation of a curve on the surface) Darboux does not put this condition; he just mentions instead that imaginary deformations are obtained if this condition is not satisfied. If one takes in Darboux's imaginary deformation case the deformed curve in discussion to be non-planar (one can change the ambient totally space along planes), then the imaginary surface thus obtained is non-totally real and has positive definite linear element.

The unit normal field N_0 of a totally real surface $x_0 \subset V$, where $\bar{v} = \epsilon v,\ \forall v \in V$, satisfies $N_0 \subset V$ if $\epsilon_3 = 1$ (in which case the second fundamental form is real) and $N_0 \subset iV$ otherwise (in which case the second fundamental form is purely imaginary). Thus x_0 has asymptotic lines iff the sign of the Gauß curvature K_0 is $-\epsilon_1\epsilon_2\epsilon_3$; if $K_0 \equiv 0$, then the asymptotic lines coincide (the rulings of the developable x_0).

Take for example surfaces $x_0 \subset \mathbb{R}^2 \times i\mathbb{R},\ x \subset \mathbb{R}^3,\ (x, dx) = (R,t)(x_0, dx_0),\ (R,t) \subset \mathbf{O}_3(\mathbb{C}) \ltimes \mathbb{C}^3$. Then with $\epsilon := \mathrm{diag}[1\ 1\ -1]$ we have $iN_0 \subset \mathbb{R}^2 \times i\mathbb{R},\ \epsilon(I_3 - 2N_0N_0^T)\bar{R}^T R = I_3$. If $K_0 > 0$, then x_0 has asymptotic lines while x does not; if $K_0 < 0$, then x has asymptotic lines while x_0 does not.

We shall now prove that the only surfaces in unit complex (pseudo-)spheres having real linear element are the totally real (pseudo-)spheres. Assume $x \subset \mathbb{C}^3,\ |x|^2 = \epsilon_1,\ \epsilon_1 := \pm 1,\ d\bar{x} = Rdx,\ R \subset \mathbf{O}_3(\mathbb{C}),\ \bar{R}^T = R$. Thus $\bar{N} = \epsilon_2 RN$, where $\epsilon_2 := \pm 1,\ |N|^2 = 1,\ N^T dx = 0$; since $N = \sqrt{\epsilon_1}x$, we get $\bar{x} = \epsilon_1\epsilon_2 Rx$. If $\epsilon_1\epsilon_2 = -1$, then one can replace R with $R(I_3 - 2NN^T) = R(I_3 - 2\epsilon_1 xx^T)$ to get $\epsilon_1\epsilon_2 = 1$; if $\epsilon_1\epsilon_2 = 1$, then $dRx = 0$ so $0 = R^{-1}dRx = \omega \times x$, or $\omega \times N = 0$; since $\omega = \omega^\top$, we get $\omega = 0$ and R is constant.

4.3. **Curves corresponding under the applicability.**
Consider $c_0,\ c$ curves on $x_0,\ x$ corresponding under the applicability. We would like to see how the curvatures and torsions transform under the applicability. If $F_c := [T_c := \frac{dc}{|dc|}\quad N_c := \frac{(dc\times d^2c)\times dc}{|(dc\times d^2c)\times dc|}\quad B_c := \frac{dc\times d^2c}{|dc\times d^2c|}]$, $\tilde F_c := [T_c\quad N\quad T_c\times N]$, $k_c := \frac{|dc\times d^2c|}{|dc|^3}$, $\tau_c := \frac{(dc\times d^2c)^T d^3c}{|dc\times d^2c|^2}$, $k_{nc} := \frac{N^T d^2c}{|dc|^2}$, $k_{gc} := -\frac{(dc\times d^2c)^T N}{|dc|^3}$, $\tau_{gc} := \frac{(N\times dN)^T dc}{|dc|^2}$ are the usual frames along c, curvatures and torsions of c (and similar quantities for c_0), then we have $F_c^{-1}dF_c = |dc|\alpha([\tau_c\ 0\ k_c]^T)$, $\tilde F_c^{-1}d\tilde F_c = |dc|\alpha([\tau_{gc}\ -k_{gc}\ k_{nc}]^T)$, $|dc|\tilde F_c^{-1}N_c = \frac{1}{k_c}\tilde F_c^{-1}dT_c = |dc|[0\ \frac{k_{nc}}{k_c}\ \frac{k_{gc}}{k_c}]^T$, $|dc|\tilde F_c^{-1}B_c = \frac{1}{\tau_c}(d(\tilde F_c^{-1}N_c) + \tilde F_c^{-1}d\tilde F_c\tilde F_c^{-1}N_c + |dc|k_c\tilde F_c^{-1}T_c) = \frac{1}{\tau_c}[0\quad d(\frac{k_{nc}}{k_c}) - |dc|\frac{\tau_{gc}k_{gc}}{k_c}\quad d(\frac{k_{gc}}{k_c}) + |dc|\frac{\tau_{gc}k_{nc}}{k_c}]^T$. Therefore $|dc|\tilde F_c^{-1}F_c = \begin{bmatrix}|dc| & 0 & 0\\ 0 & |dc|\frac{k_{nc}}{k_c} & \frac{1}{\tau_c}(d(\frac{k_{nc}}{k_c}) - |dc|\frac{\tau_{gc}k_{gc}}{k_c})\\ 0 & |dc|\frac{k_{gc}}{k_c} & \frac{1}{\tau_c}(d(\frac{k_{gc}}{k_c}) + |dc|\frac{\tau_{gc}k_{nc}}{k_c})\end{bmatrix} = |dc|\begin{bmatrix}1 & 0 & 0\\ 0 & \frac{k_{nc}}{k_c} & -\frac{k_{gc}}{k_c}\\ 0 & \frac{k_{gc}}{k_c} & \frac{k_{nc}}{k_c}\end{bmatrix}$ and we have $|dc|\tau_c = |dc|\tau_{gc} + \frac{k_{nc}dk_{gc} - k_{gc}dk_{nc}}{k_c^2}$ (Bonnet).

We have $\tilde F_c = R|_{c_0}\tilde F_{c_0}$, so $\tilde F_c^{-1}d\tilde F_c = \tilde F_{c_0}^{-1}d\tilde F_{c_0} + \tilde F_{c_0}^{-1}R^{-1}dR|_{c_0}\tilde F_{c_0}$. Therefore

$$k_{nc} = k_{nc_0} + (\frac{dc_0}{|dc_0|}\times N_0)^T\frac{\omega|_{c_0}}{|dc_0|},$$
$$k_{gc} = k_{gc_0},\ \tau_{gc} = \tau_{gc_0} - \frac{dc_0^T}{|dc_0|}\frac{\omega|_{c_0}}{|dc_0|}. \tag{4.10}$$

Note that if we choose (u,v) coordinates on x such that $c(u) = x(u,0)$ and $v =$const are parallel, $u =$const are geodesics, then $\tau_{gc} = \frac{N^T x_{uv}}{|x_u||x_v|}$, so $\tau_{gc} = \tau_c = \sqrt{-K}$ for c asymptote on x (Enneper). In particular the asymptotes of CGC surfaces are curves of constant torsion.

Conversely, we can compute the curvatures and torsions of the corresponding curve c_R on R in function of the curvatures and torsions of $c_0,\ c$:
$|dc_R| = |c_R^{-1}dc_R| = |\omega| = \sqrt{(\tau_{gc}-\tau_{gc_0})^2 + (k_{nc}-k_{nc_0})^2}|dc_0|$, $k_{nc_R} = \frac{k_{nc_0}\tau_{gc} - \tau_{gc_0}k_{nc}}{(\tau_{gc}-\tau_{gc_0})^2 + (k_{nc}-k_{nc_0})^2}$,
$k_{gc_R}|dc_0| = \frac{(k_{nc}-k_{nc_0})d(\tau_{gc}-\tau_{gc_0}) - (\tau_{gc}-\tau_{gc_0})d(k_{nc}-k_{nc_0})}{((\tau_{gc}-\tau_{gc_0})^2 + (k_{nc}-k_{nc_0})^2)^{\frac{3}{2}}}$, $\tau_{gc_R} = \frac{\tau_{gc_0}(\tau_{gc}-\tau_{gc_0}) + k_{nc_0}(k_{nc}-k_{nc_0})}{(\tau_{gc}-\tau_{gc_0})^2 + (k_{nc}-k_{nc_0})^2}$.

4.4. **Ruled surfaces.**
If $[R^{-1}dR, \wedge R^{-1}dR] = 0$, then by a change of coordinates we can suppose $R_u = 0$, so $x_{0u} = \phi_u a(v)$, $R^{-1}dRa = 0$, $|a|^2 = 1$ or $a = Y(v)$, $\phi_u \neq 0$ and $x_0(u,v) = \phi(u,v)a(v) + b(v)$, $b(v)$ arbitrary such that $a\times(\phi a_v + b_v) \neq 0$. With the change of coordinates $(u',v') = (\phi(u,v),v)$, we get the formula for ruled surfaces: $x_0(u,v) = ua(v) + b(v)$ (for convenience after all these changes of coordinates we still keep the notation (u,v) for our parametrization).

In this case $x = uRa + c$, where $c(v) = \int Rdb = Rb - \int\varphi R(a\times b)|db|$ and R is a solution of the ODE $R^{-1}dR = \varphi|db|\alpha(a)$. The line of striction b of x_0 is defined as $da^Tdb = 0$ (the curve along which the rulings are closest) can be realized after a change of the u parameter. In particular if x_0 is developable, then its line of striction coincides with the line of regression. Because $d(Ra)^TRdb = db^T(da + R^{-1}dRa) = 0$, the line of striction is invariant under deformations preserving the rulings.

We can prescribe various conditions on c by choosing suitable φ's (see Eisenhart ([19],§ 144)).

In our case $N_0 = \frac{a\times db}{|a\times db|}$, so we have $k_{nc} = \frac{d^2b^T(a\times db)}{|a\times db||db|^2} + \varphi\frac{|a\times db|}{|db|}$, $\tau_{gc} = (1 - \frac{a^Tdb}{|db|})\frac{(a\times db)^T}{|db|}(\frac{a^Tdbd^2b}{|db|^3} - (1 + \frac{a^Tdb}{|db|})\frac{da}{|db|}) - \varphi\frac{a^Tdb}{|db|}$.

Therefore we can prescribe all quantities except the torsion by solving an ODE; for the torsion we need to solve two ODE's.

According to Bonnet, if two surfaces $x_0,\ x$ are applicable with corresponding asymptotes of parameter u, then $R_ux_{0u} = R_ux_{0v} = 0$, so $R_u = 0$, so $x_0,\ x$ are ruled with the rulings corresponding under the applicability.

Also (see Eisenhart ([19],§ 144)) if $x_0,\ x$ are ruled and applicable with rulings not corresponding under the applicability, then choosing parametrization (u,v) such that the rulings are the curves $u=$const, $v=$const, (u,v) will be virtual asymptotes which are geodesics. Thus there is a surface on which (u,v) are asymptotes and geodesics, so $x_0,\ x$ are applicable (with correspondence of rulings) to a quadric.

According to a theorem of Chieffi, if we have a deformation of a ruled surface, then the ruled surface formed by the tangents (to the geodesics corresponding to rulings) along an asymptotic line is applicable to the initial ruled surface.

If $x_0(u_0,v_0)=u_0a(v_0)+b(v_0),\ (x,dx)=(R,t)(x_0,dx_0)$ and (u,v) is the parametrization of x with v asymptotic parameter, then setting $u:=$const the ruled surface $\tilde{x}(w,v):=wx_{u_0}+x=R(wa+x_0)+t$ is applicable to $\tilde{x}_0(w,v):=wa+x_0=(w+u_0)a+b=x_0(w+u_0,v_0)$, because $d\tilde{x}=Rd\tilde{x}_0+wdRx_{0u_0}=Rd\tilde{x}_0$. Here we used the fact that $dRx_{0u_0}=R_vx_{0u_0}dv=0$, which follows from (4.3) (because $u_0=$const, $v=$const are asymptotes on x_0 respectively x, the Gauß curvature information is recorded only in the off-diagonal terms of the second fundamental forms of $x_0,\ x$, so $0=N_0^T(R^{-1}x_{u_0v}-x_{0u_0v})=R^{-1}R_vx_{0u_0}$).

Because $u=$const is an asymptote on x, its osculating spaces coincide with the tangent spaces of x along $u=$const; therefore $u=$const is an asymptote also on $\tilde{x}$.

For example, because the catenoid is applicable to the minimal helicoid, it is applicable to the ruled surface formed by the tangents (to the geodesics corresponding to the rulings of the minimal helicoid) along one of its asymptotes. This example (for surfaces applicable to the catenoid) is due to Bianchi and was the inspiration of Chieffi's result.

Note that we have the more general result, due to Bianchi: if $x,\ x_0$ are applicable surfaces with rolling $(x,dx)=(R,t)(x_0,dx_0)$ and we have $(u_0,v_0),\ (u,v)$ parameterizations on $x_0,\ x$ with (u_0,v) kinematically conjugate, then setting $u:=$const the ruled surface $\tilde{x}(w,v):=wx_{u_0}+x$ is applicable to the ruled surface $\tilde{x}_0(w,v):=wx_{0u_0}+x_0$ with the same rolling as that of $(x_0,\ x)$ and correspondence of rulings. We have $\tilde{x}=R\tilde{x}_0+t,\ d\tilde{x}=Rd\tilde{x}_0+wdRx_{0u_0}=Rd\tilde{x}_0$, because $dRx_{0u_0}=R_vx_{0u_0}dv=0$. Darboux in ([13],§ 932) has a result in the same vein: if $(c_0,\ c)$ are curves of $x_0,\ x$ corresponding under the applicability, then since $0=R^{-1}dR|_{c_0}\frac{\omega|_{c_0}}{|dc_0|}$, the ruled surfaces $c_0+w\frac{\omega|_{c_0}}{|dc_0|},\ c+wR\frac{\omega|_{c_0}}{|dc_0|}$ are applicable with rolling $(R,t)|_{c_0}$.

Conversely, if $(u_0,v_0),\ (u,v)$ are a-priori arbitrary parameterizations on $x_0,\ x$ and a point-wise correspondence is established between $x_0,\ x$ such that for any $u=$const the ruled surface $\tilde{x}(w,v):=wx_{u_0}+x$ is applicable to the ruled surface $\tilde{x}_0(w,v):=wx_{0u_0}+x_0$, $(\tilde{x},d\tilde{x})=(R,t)(\tilde{x}_0,d\tilde{x}_0)$ with correspondence of rulings, then $x_{u_0}dw+wdx_{u_0}+dx=d\tilde{x}=Rd\tilde{x}_0=Rx_{0u_0}dw+wRdx_{0u_0}+Rdx_0$, so $x_{u_0}=Rx_{0u_0},\ wx_{u_0v}+x_v=wRx_{0u_0v}+Rx_{0v}$. Setting $w=0$ we get $x_v=Rx_{0v}$, so x is applicable to x_0. Moreover we get $x_{u_0v}=Rx_{0u_0v}$, so $R_vx_{0u_0}=0$ and thus (u_0,v) are kinematically conjugate.

In this vein according to Bianchi (122) Lie has reduced the study of B transformations of the (pseudo-)sphere to the study of B transformations of their asymptotes, curves of constant torsion. Also, according to Bianchi (122,§ 67), Serret has integrated the equation of ruled CGC surfaces: all are obtained by choosing the ruling as one of the isotropic directions in the osculating plane of a curve of constant torsion. This can be easily justified with Chieffi's and Enneper's results; such surfaces x are of the form $x(u,v)=\frac{1}{u-v}R(v)Y(v)-\int R(v)f_1dv,\ R\subset\mathbf{O}_3(\mathbb{C}),\ R'(v)Y(v)=0$ for the CGC 1 case and $ix,\ x$ as above for the CGC -1 case. In particular if we let $u\bar{v}=-2\ (=2)$ for the CGC 1 (-1) case, then the linear element of $x\ (ix)$ is real positive definite; such surfaces $x\ (ix)$ are real analytic. Further for the CGC 1 (-1) case we have $d\bar{x}=\overline{R(v)}R(v)^Tdx\quad(d\overline{(ix)}=\overline{R(v)}(I_3-2e_3e_3^T)R(v)^Td(ix))$; should $\overline{R(v)}R(v)^T\quad(\overline{R(v)}(I_3-2e_3e_3^T)R(v)^T)$ be constant, $\overline{R(v)}$ would be holomorphic in v and in $\bar{v}$, so it would be a constant. Thus $x\ (ix)$ is non-totally real except when $x\ (ix)$ is up to rigid motions the standard totally real unit (pseudo-)sphere.

4.5. **Infinitesimal deformations, Weingarten congruences, Darboux's 12 surfaces and transformations of surfaces.**

(see also Sabitov [27])

Let s_1 be an infinitesimal deformation of the surface s_0 : $0 = \frac{d}{d\epsilon}|_{\epsilon=0}|d(s_0+\epsilon s_1)|^2 = ds_0^T ds_1 + ds_1^T ds_0$. Therefore there is s_2 (the infinitesimal rotation of the infinitesimal deformation) such that $ds_0 = s_2 \times ds_1$. With $s_3 := s_0 - s_2 \times s_1$ we have $ds_3 = s_1 \times ds_2$ and $ds_2^T ds_3 + ds_3^T ds_2 = 0$, so there is s_4 such that $ds_2 = s_4 \times ds_3$ and we can iterate this construction (replace $s_k,\ k = 0, 1$ with $s_{2j+k}, j = 1, 2, ...$). Darboux proved in ([13],§ 883-§ 924) that this process ends after 12 steps ($s_j = s_{j+12}$) and found interesting relationships between these 12 surfaces.

We have $ds_{2j+3} = s_{2j+1} \times ds_{2j+2} = s_{2j+1} \times (s_{2j+4} \times ds_{2j+3}) = -s_{2j+1}^T s_{2j+4} ds_{2j+3}$, so $s_{2j+1}^T s_{2j+4} + 1 = 0$. Also $ds_{2j+2} = s_{2j+4} \times ds_{2j+3} = s_{2j+4} \times (s_{2j+1} \times ds_{2j+2}) = -s_{2j+1}^T s_{2j+4} ds_{2j+2} + ds_{2j+2}^T s_{2j+4} s_{2j+1}$, so $ds_{2j+2}^T s_{2j+4} = 0$.

But $ds_{2j+2} \times \wedge ds_{2j+1} = d \wedge ds_{2j} = 0$, so $s_{2j+1},\ s_{2j+2}$ are associate; in particular they have parallel tangent planes, so $ds_{2j+1}^T s_{2j+4} = 0$, so $s_{2j+1},\ s_{2j+4}$ are *polar reciprocal with respect to the pseudo-sphere* (prwrtps).

From $s_{2j+4}^T ds_{2j+1} = s_{2j-1}^T ds_{2j+1} = 0$ we get $s_{2j+4} = \varphi s_{2j-1}$ for some function φ. Multiplying this last relation on the left with $s_{2j+1}^T,\ s_{2j+2}^T$ we get $s_{2j+4} = -\frac{s_{2j-1}}{s_{2j+1}^T s_{2j-1}},\ s_{2j-1} = -\frac{s_{2j+4}}{s_{2j+2}^T s_{2j+4}}$. Therefore $s_{2j+3} = s_{2j} + \frac{s_{2j-3} \times s_{2j+1}}{s_{2j-3}^T s_{2j-1}}$ and $s_{2j-3}^T s_{2j+3} + 1 = 0$. Then $s_{2j+7} = s_{2j+4} + \frac{s_{2j+1} \times s_{2j+5}}{s_{2j+1}^T s_{2j+3}} = -\frac{s_{2j-1}}{s_{2j+1}^T s_{2j-1}} + \frac{s_{2j+1} \times (s_{2j-1}^T s_{2j+1} s_{2j+2} + s_{2j-1} \times s_{2j+3})}{s_{2j+1}^T s_{2j+3} s_{2j-1}^T s_{2j+1}} = -\frac{s_{2j-1}}{s_{2j+1}^T s_{2j-1}} + \frac{s_{2j+1} \times s_{2j+2}}{s_{2j}^T s_{2j+1}} - \frac{s_{2j+3}}{s_{2j+1}^T s_{2j+3}} + \frac{s_{2j-1}}{s_{2j-1}^T s_{2j+1}} = -\frac{s_{2j}}{s_{2j}^T s_{2j+1}} = -\frac{s_{2j}}{s_{2j}^T s_{2j-2}} = s_{2j-5}$ and $s_{2j+6} = -\frac{s_{2j+1}}{s_{2j+3}^T s_{2j+1}} = -\frac{s_{2j-11}}{s_{2j-9}^T s_{2j-11}} = s_{2j-6}$.

The prwrtps is uniquely determined: $s_{2j+1} = \frac{-N_{2j+4}}{s_{2j+4}^T N_{2j+4}},\ s_{2j+4} = \frac{-N_{2j+1}}{s_{2j+1}^T N_{2j+1}}$; s_{2j+1} and s_{2j+2} are associate (parallel tangent planes and to asymptotes on one surface corresponds on the other a conjugate system). Conversely from two associate surfaces $s_1,\ s_2$ ($ds_1 \times \wedge ds_2 = 0$) we can recover s_0 by a quadrature from $ds_0 = s_2 \times ds_1$.

We have $s_{2j+3} - s_{2j} = s_{2j+1} \times s_{2j+2}$; since $s_{2j+2}^T ds_{2j} = s_{2j+1}^T ds_{2j+3} = 0$ we see that $s_{2j+3} - s_{2j}$ is tangent to both $s_{2j},\ s_{2j+3}$. With $\phi := s_{2j+2}^T s_{2j+4}$ we have $0 = ds_{2j+3}^T ds_{2j+1} (\phi s_{2j+1} + s_{2j+2})^T s_{2j+4} = (s_{2j+4} \times ds_{2j+3})^T ((\phi s_{2j+1} + s_{2j+2}) \times ds_{2j+1}) = ds_{2j+2}^T (\phi s_{2j+1} \times ds_{2j+1} + ds_{2j}) = \phi d^2 s_{2j+3}^T s_{2j+1} - d^2 s_{2j}^T s_{2j+2}$; therefore the second fundamental forms of $s_{2j},\ s_{2j+3}$ are proportional and $s_{2j},\ s_{2j+3}$ are the focal surfaces of a W congruence.

Conversely Guichard proved that if $s_0,\ s_3$ are the focal surfaces of a W congruence, then we can recover infinitesimal deformations $s_1, (s_2)$ of $s_0, (s_3)$ such that $s_1^T ds_3 = s_2^T ds_0 = 0,\ s_3 - s_0 = s_1 \times s_2$. These $s_j,\ j = 0, 3$ will satisfy $ds_0 - s_2 \times ds_1 = ds_3 - s_1 \times ds_2,\ 0 = s_2^T (ds_0 - s_2 \times ds_1) = s_1^T (ds_3 - s_1 \times ds_2) = ds_1^T (ds_0 - s_2 \times ds_1 + ds_1^T (ds_0 - s_2 \times ds_1) = ds_2^T (ds_3 - s_1 \times ds_2) + ds_2^T (ds_3 - s_1 \times ds_2)$, so $0 = ds_0 - s_2 \times ds_1 = ds_3 - s_1 \times ds_2$.

To prove Guichard's result we refer $s_0,\ s_3$ to their asymptotic coordinates $u,\ v$: $\theta_j := |\theta_j| N_j,\ |\theta_j|^2 := \frac{|s_{ju}|}{|N_{ju}|} = \frac{|s_{jv}|}{|N_{jv}|}$, so $ds_j = \theta_j \times (\theta_{ju} du - \theta_{jv} dv),\ N_j = \frac{\theta_j}{|\theta_j|},\ \theta_j \times \theta_{juv} = 0,\ j = 0$ or 3 (Lelieuvre formulae). All infinitesimal deformations s_1 of s_0 are given by: $ds_1 = \theta_0 (\tilde{\theta}_{0u} du - \tilde{\theta}_{0v} dv) - \tilde{\theta}_0 (\theta_{0u} du - \theta_{0v} dv)$, where $\tilde{\theta}_0,\ (\theta_0)$ are scalar (vector) solutions of the Laplace equation $\theta_{uv} = M_0(u, v)\theta$; then $s_2 := \frac{\theta_0}{\theta_0}$ satisfies $ds_0 - s_2 \times ds_1 = 0$. We have $s_3 - s_0 = \rho \theta_0 \times \theta_3$ for some function ρ, so $\rho d\theta_0^T (\theta_0 \times \theta_3) = -\theta_0^T d(s_3 - s_0) = -(\theta_0 \times \theta_3)^T (\theta_{3u} du - \theta_{3v} dv)$ and $\rho d\theta_3^T (\theta_0 \times \theta_3) = -\theta_3^T d(s_3 - s_0) = -(\theta_0 \times \theta_3)^T (\theta_{0u} du - \theta_{0v} dv)$. Therefore $(\rho^2 - 1) d\theta_0^T (\theta_0 \times \theta_3) = (\rho^2 - 1) d\theta_3^T (\theta_0 \times \theta_3) = 0$ and $\rho = 1$ (we can absorb the sign in θ_0; also $\theta_0 \times \theta_3 = 0$ is impossible). Thus $0 = ds_3 - ds_0 - d\theta_0 \times \theta_3 - \theta_0 \times d\theta_3 = (\theta_3 - \theta_0) \times (\theta_3 + \theta_0)_u du - (\theta_3 + \theta_0) \times (\theta_3 - \theta_0)_v dv$ and $(\theta_3 + \theta_0)_u = \log(\tilde{\theta})_u (\theta_3 - \theta_0),\ (\theta_3 - \theta_0)_v = \log(\theta)_v (\theta_3 + \theta_0)$ for some functions $\tilde{\theta},\ \theta$. Differentiating these two last relations wrt $v,\ u$ and using the relations themselves we get $0 = \theta_3 (M_3 - \log(\tilde{\theta})_{uv} - \log(\tilde{\theta})_u \log(\theta)_v) + \theta_0 (M_0 + \log(\tilde{\theta})_{uv} - \log(\tilde{\theta})_u \log(\theta)_v) = \theta_3 (M_3 - \log(\theta)_{uv} - \log(\tilde{\theta})_u \log(\theta)_v) - \theta_0 (M_0 + \log(\theta)_{uv} - \log(\tilde{\theta})_u \log(\theta)_v)$, or $(\log(\theta) - \log(\tilde{\theta}))_{uv} = 0,\ M_0 = -\log(\tilde{\theta})_{uv} + \log(\tilde{\theta})_u \log(\theta)_v,\ M_3 = \log(\tilde{\theta})_{uv} + \log(\tilde{\theta})_u \log(\theta)_v$. Therefore $\tilde{\theta}_3 := f(v)\tilde{\theta} = g(u)\theta =: \frac{1}{\theta_0}$ for some functions $f,\ g,\ \tilde{\theta}_{0uv} = M_0 \tilde{\theta}_0,\ \tilde{\theta}_{3uv} = M_3 \tilde{\theta}_3$ and

$s_1 := \frac{\theta_3}{\theta_3}$, $s_2 := \frac{\theta_0}{\theta_0}$ are the surfaces we are looking for. To the geometric W congruence corresponds analytically the Moutard transformation of solutions of the Laplace equations with functions M_0, M_3. Note that $K(s_0) = -\frac{1}{|\theta_0|^4}$, so a congruence is W iff its focal surfaces s_0, s_3 satisfy $K(s_0)K(s_3) = \frac{|N_0 \times N_3|^4}{|s_0 - s_3|^4}$ (Ribacour).

Thus the infinitesimal deformation problem becomes equivalent to W congruences. These in turn become *Ribacour sphere congruences* (2-dimensional families of spheres upon whose envelopes the lines of curvature correspond) via Lie's contact transformation. The envelopes of the Ribacour sphere congruences correspond to the focal surfaces of the W congruence. Any transformation preserving W congruences (homographies, correlations) preserves the problem of infinitesimal deformation.

If we take any two surfaces of the 12 and which are not prwrtps one of the other, then by taking their prwrtps we get four surfaces among which there will be 2 who are in the relation: one infinitesimal deformation of the other, associate or focal surfaces of a W congruence, so we can recover the 12 surfaces.

Since a linear isomorphism of a plane preserves the cross-ratio of four lines through the origin, a correspondence between two surfaces preserves the cross-ratio of four vector fields. Thus if asymptotic directions correspond under the correspondence (which is the case for the focal surfaces of a W congruence), then all conjugate systems correspond (because conjugate directions and asymptotic directions have cross-ratio 1).

On the 12 surfaces we have three systems of coordinates of interest; each will give on each surface asymptotic or conjugate systems with equal (tangential) invariants (that is $\Gamma^2_{12v} = \Gamma^1_{12u}$, where the Christoffel symbols are taken wrt the linear element of the surface (Gauß map)). On two surfaces one infinitesimal deformation of the other the asymptotic coordinates on one correspond to a conjugate system with equal invariants on the other and the common conjugate system has equal tangential invariants on both. On two associate surfaces the asymptotic coordinates on one correspond to a conjugate system with equal tangential invariants on the other and the common conjugate system has equal invariants on both surfaces. On the focal surfaces of a W congruence (or on a surface and its prwrtps) the asymptotic coordinates correspond and the conjugate system with equal invariants on one corresponds to a conjugate system with equal tangential invariants on the other. Knowledge of a conjugate system with equal invariants gives by a quadrature the associate surface (unique modulo translations) upon which the given conjugate system is still conjugate with equal invariants, so knowledge of such systems is equivalent to the problem of infinitesimal deformation.

Consider the congruence $s_1 + t(s_2 - s_1)$; it cuts its focal surfaces and s_1, s_2 with cross-ratio 1. The developables of the congruence (inducing conjugate systems with equal invariants on the focal surfaces) correspond to the common conjugate system on s_1, s_2; therefore Koenigs considered such surfaces s_1, s_2 as transforms one of the other (a particular case is the D transformation of isothermic surfaces). Eisenhart in [20] and Jonas have generalized this transformation to the *fundamental* (F) transformation (they do not require equal invariants). According to Eisenhart [20] this transformation F together with Bianchi's generalized *Bäcklund* (B) transformation (the focal surfaces of a W congruence are considered as transforms one of the other) account for most if not all of transformations of surfaces for which an interesting theory has been developed (including for example variants of the BPT).

4.6. **Rolling congruences.**
All results of this subsection in the particular case when the curve c is a line appear in Bianchi ([5], Vol**7**,(157),(166)).

Let $c := c(w)$ be a curve in $\mathbb{C}^3$. As x_0 rolls on x and rigidly moving c the curve c will describe a congruence of congruent curves $(R(u,v), t(u,v))c(w)$. We are interested in the focal surfaces (envelopes) of such a congruence, that is prescribing $w(u,v)$ such that the surface $(R(u,v), t(u,v))c(w(u,v))$ meets tangentially the curve $w \to (R(u,v), t(u,v))c(w)$. We thus need

$0 = (Rdc)^T(d((R,t)c) \times \wedge d((R,t)c)) = dc^T((dc + \omega \times (c - x_0)) \times \wedge(dc + \omega \times (c - x_0))) = dc^T((\omega \times (c - x_0)) \times \wedge(\omega \times (c - x_0))) = N_0^T(\omega \times \wedge\omega)N_0^T(c - x_0)dc^T(c - x_0)$. Therefore the focal surfaces are given by $N_0^T(c - x_0) = 0,\ dc^T(c - x_0) = 0$ and are thus independent of the rolling (we exclude the case $N_0^T(\omega \times \wedge\omega) = 0$ of $x_0,\ x$ ruled with correspondence of rulings).

Thus the focal surfaces are given by the intersection(s) of the tangent plane of x_0 with c in the first case and the perpendicular(s) from x_0 to c in the second case, since the normals to the focal surfaces are multiples of $R(N_0 \times dc)$ in the first case and of $R(c - x_0)$ in the second case, the focal planes are the tangent plane to c normal to the tangent plane of x_0 in the first case and normal to $c - x_0$ in the second case.

Conversely, if a congruence $C(u, v, w) := (R(u, v), t(u, v))c(w)$ of congruent curves has the property that on the normals of a focal surface $F_2 := C(u, v, w_2(u, v))$ one can choose a surface x such that the tangent space of the surface x is normal to the tangent space of another focal surface $F_1 := C(u, v, w_1(u, v))$, then this congruence is in general a congruence of rolling. If c does not admit a continuous group of symmetries (that is c does not have constant curvature and torsion; equivalently c is not a line, circle or helix; note however that a line does not have curvature and torsion defined geometrically; equivalently from a geometric point of view a line can have arbitrarily prescribed curvature and torsion), then we can recover (R, t) (up to finitely many choices, as the curve c may have a finite group of symmetries) from the knowledge of the congruence C. With $x_0 := R^{-1}(x - t)$ and because $F_1,\ F_2$ are focal surfaces we get $0 = d(c(w_2))^T(c(w_2) - x_0) = (R^{-1}dF_2)^T(c(w_2) - x_0) = (R^{-1}N)^T(c(w_1) - x_0) = (R^{-1}dF_1)^T((R^{-1}N) \times d(c(w_1)))$. Since $R^{-1}dF_j = R^{-1}dRc(w_j) + d(c(w_j)) + R^{-1}dt,\ j = 1, 2$ we get $0 = (c(w_2) - x_0)^T(R^{-1}dRx_0 + R^{-1}dt)$...

4.7. **Rolling distributions.**
Ribaucour proved that if we take a tangential congruence of curves (each curve in a tangent space of x_0), then the integrability condition of the distribution of facets formed by the normal planes of the curves depends only on the linear element of x_0, so if the distribution in the initial position was integrable and we roll x_0 on an applicable surface x, then the rolled distribution remains integrable. The statement includes the degenerate case of a congruence of curves in the tangent planes of a curve x_0, in which case the integrability depends on the arc-length and curvature of x_0, so if we deform the tangent surface of x_0, then the distribution remains integrable if it was integrable in the initial position (see Bianchi ([4],§ 359)).

In the particular case when the curves are circles and the distribution is integrable such a congruence of circles is called *cyclic system*. In particular cyclic systems with all circles of the same radius 1 exist when x_0 has CGC -1 and the centers of circles are the corresponding points of x_0; in this case the leaves of the distribution are Bianchi's complementary transforms (also CGC -1 surfaces).

Darboux proved that all cyclic systems appear when we intersect the tangent spaces of x_0 with a particular isotropic cone thus highlighting a circle in each tangent space. The leaves of the distribution in this case are the rulings of the isotropic cone and in general will be the focal surfaces of the rolling congruences of rulings (Darboux ([13],§ 936). When x_0 is a quadric and the isotropic cones have umbilics as vertices among the leaves of the rolled distribution are special isothermic surfaces (which envelope the congruence of rolled isotropic rulings).

Bianchi was able to prove in ([5],Vol **4**,(108)) the Permutability Theorem for the D transformation of isothermic surfaces, a thing considered miraculous at the time since the Ricatti character of the Darboux system was not evident. The Ricatti character was recently discovered and involves Clifford multiplication (see Burstall [9]). Bianchi proved in ([5],Vol **4**,(114)) that a D transformation for special isothermic surfaces is a composition of two B transformations of quadrics.

Consider an integrable 3-dimensional distribution of facets $(p, P) = (p, P)(u, v, w)$ in $\mathbb{C}^3$ along the surface $x = x(u, v)$ (thus distributed as a 1-dimensional distribution of facets along each point of x). It is thus natural to inquire when this distribution remains integrable if we roll it as x rolls on any applicable surface x_0 (see Bianchi ([5],Vol **4**,(173))). Let $m = m(u, v, w)\perp P(u, v, w)$ be a

normal field along the distribution. We shall use the notation: $\tilde{df} = f_u du + f_v dv + f_w dw = df + f_w dw$ for $f = f(u, v, w)$. If the distribution is integrable with leaves $p(u, v, w(u, v))$, then $0 = m^T \tilde{dp} = m^T p_w dw + m^T dp$ holds. In particular if $m^T N = N^T(p - c(w)x) = 0$, then the integrability condition depends only on the linear element of x. Imposing the integrability condition $m^T p_w \tilde{d}\wedge$ (suppose $m^T p_w \neq 0$) and using the equation itself we get: $(m_w^T dp - dm^T p_w) \wedge m^T dp + m^T p_w dm^T \wedge dp = 0$, or:

$$(dp \times p_w)^T \wedge (m \times dm) + \frac{1}{2}(m_w \times m)^T (dp \times \wedge dp) = 0. \tag{4.11}$$

If $(x_0, dx_0) = (R, t)(x, dx)$ is the rolling of x on x_0, then the rolled distribution will be $(R, t)(V + x, P) = (RV + x_0, RP)$, $V := p - x$, so (4.11) must be satisfied with RV, Rm replacing V, m: $((d(V+x)+\omega\times V)\times V_w)^T \wedge (m\times(dm+\omega\times m)) + \frac{1}{2}(m_w \times m)^T((d(V+x)+\omega\times V)\times\wedge(d(V+x)+\omega\times V)) = = 0$, or, using (4.11): $((m\times((d(V+x)\times V_w)\times m) - V\times(V_w\times(m\times dm) + d(V+x)\times(m_w\times m)))^T \wedge \omega + (V^T(m_w\times m)V + V_w\times((m\times V)\times m))^T \frac{\omega\times\wedge\omega}{2} = 0$. Using the Gauß theorem ($dN \times \wedge dN = KN\sqrt{g}du \wedge dv$) for both x_0, x we have $\frac{\omega\times\wedge\omega}{2} = \frac{1}{2}N^T((R^{-1}dN_0 - dN) \times \wedge(R^{-1}dN_0 - dN))N = -N^T(dN \times \wedge(R^{-1}dN_0 - dN))N = dN^T \wedge \omega N$ and we thus need: $(m\times((d(V+x)\times V_w)\times m) - V\times(V_w\times(m\times dm) + d(V+x)\times(m_w\times m)) + (V^T(m_w\times m)V + V_w\times((m\times V)\times m))^T N dN)^T \wedge \omega = 0$ for any flat connection form ω. This is equivalent to $(m\times((d(V+x)\times V_w)\times m) - V\times(V_w\times(m\times dm) + d(V+x)\times(m_w\times m)) + (V^T(m_w\times m)V + V_w\times((m\times V)\times m))^T N dN)^T dx = 0$, or using $dx \times dx = Nda = N\sqrt{g}du \wedge dv$:

$$\begin{aligned}(m^T d(V+x)(m\times V)_w - ((m^T d(V+x))_w - d(m^T V_w))m\times V - m^T V_w d(m\times\ V))^T dx - \\ m^T V_w m^T N da + ((m\times V)\times(m\times V)_w - m\times V m^T V_w)^T N N^T d^2x = 0.\end{aligned} \tag{4.12}$$

But (4.12) is satisfied if RV, Rm replace V, m, so it depends only on the linear element of x, as expected. Therefore the distributions with the required properties must satisfy (4.11) and (4.12) in which we ignore the terms involving linearly the second fundamental form of x (these equation can also be obtained from (4.11) (which becomes an affine equation in the second fundamental form of x by an application of the Gauß theorem) by setting all the coefficients of the second fundamental form of x to 0). We have $dV = V^I + V^{II}$, $V^I := d(V^T N)N + d(V^T \nabla x \lrcorner dx^T)\nabla x \lrcorner dx$, $V^{II} := N^T d^2 x_\llcorner((\nabla x \times N)\times V)$, $N \times dx = -\nabla x \lrcorner da$, $m \times V = (m^T \nabla x V^T \nabla x N + \begin{vmatrix} m^T N & m^T \nabla x \\ V^T N & V^T \nabla x \end{vmatrix} \nabla x)\lrcorner\lrcorner da$. Therefore we need:

$$\begin{aligned}((V^I + dx)\times V_w)^T \wedge (m\times m^I) + \frac{1}{2}(m_w\times m)^T((V^I + dx)\times\wedge(V^I + dx)) - \\ KN^T((m\times V)\times(m\times V)_w - m\times V m^T V_w)da = 0, \\ (m^T(V^I + dx)(m\times V)_w - ((m^T(V^I + dx))_w - d(m^T V_w))m\times V - \\ m^T V_w(m\times V)^I)^T dx - m^T V_w m^T N da = 0.\end{aligned} \tag{4.13}$$

In particular if the distribution is integrable for a particular deformation of x (which can be taken as the definition of the distribution), then we need not consider the first equation. If we take the intersections of the tangent planes of x with an isotropic developable (rulings are isotropic vectors, or tangent to $C(\infty)$), then these curves are like in Ribacour's result and the leaves of the distribution in this initial position are the rulings (Darboux ([13],§ 762)).

Suppose that the leaves of the initial distribution (p, P) are curves: $0 = \tilde{d}(V + x) \times \wedge \tilde{d}(V + x) = V_w^T(d(V + x) \times \wedge d(V + x))m^T V_w m$. Since $m^T V_w \neq 0$ this is equivalent to $0 = V_w^T(KN^T V V da + (V^I + dx) \times \wedge(V^I + 2V^{II} + dx))$...

5. Deformations in $\mathbb{C}^3$ of quadrics in $\mathbb{C}^3$

5.1. **The Ricatti equation and the applicability correspondence provided by the Ivory affinity.**

Let $x^0 \subset \mathbb{C}^3$ be non-rigidly applicable to the surface x_0^0 in the quadric x_0: $(x^0, dx^0) = (R_0, t_0)(x_0^0, dx_0^0)$. If the facets of the distribution $\mathcal{D}$ become tangent spaces to surfaces (leaves) $x^1 := (R_0, x^0)V_0^1$ when rigidly moved with x_0^0 when it rolls on x^0, then $0 = (R_0 m_0^1)^T dx^1$ holds.

This will constitute a completely integrable Ricatti equation on the parameter of the conics $T_{x_0^0}x_0 \cap x_z$ (chosen to be v_1), so the arbitrariness of the transformation B_z taking x^0 into x^1 will be given by the choice of a point on an initial conic (for simplicity we ignore this 1-dimensionality in the notation B_z); 4 transforms $B_z(x^0)$ will cut these conics with constant cross-ratio. Therefore to find all transforms $B_z(x^0)$ with fixed z we need only one solution of this Ricatti equation, two of a linear differential equation and further algebraic computations.

We have: $R_0^{-1}dx^1 = d(x_0^0 + V_0^1) + R_0^{-1}dR_0V_0^1 = dx_z^1 + \omega_0 \times V_0^1$, $\omega_0 := N_0^0 \times R_0^{-1}dR_0N_0^0$. But $(\omega_0)^\perp = 0$ and $dx_z^1 = x_{zv_1}^1 dv_1 + x_{zu_1}^1 du_1$, so $0 = (R_0 m_0^1)^T dx^1$ becomes:

$$-(V_0^1)^T(\omega_0 \times N_0^0)(m_0^1)^T N_0^0 + (m_0^1)^T x_{zv_1}^1 dv_1 = 0. \tag{5.1}$$

If similarly to Bianchi's original approach (122,§ 32) we postpone the proof of the complete integrability of (5.1) until later and first try to prove the ACPIA: $|dx^1|^2 = |dx_0^1|^2$, then the necessary algebraic conditions (3.12) appear.

Using the second relation of (3.12), (5.1) becomes:

$$-(V_0^1)^T(\omega_0 \times N_0^0) + 2(x_{zv_1}^1)^T N_0^0 dv_1 = 0. \tag{5.2}$$

Multiplying this by $\mathcal{B}_1(x_{zu_1}^1)^T N_0^0$ and using $-\mathcal{B}_1(x_{zu_1}^1)^T N_0^0 V_0^1 = \mathcal{B}_1(V_0^1 \times x_{zu_1}^1) \times N_0^0 = -m_0^1 \times N_0^0$ we finally get:

$$(m_0^1)^T\omega_0 + 2z dv_1 = 0. \tag{5.3}$$

Thus the Ricatti character of the equation appears clearly. We have $dm_0^1 = m_{0v_1}^1 dv_1 + \mathcal{B}_1 dx_0^0 \times x_{zu_1}^1$, so $d(m_0^1)^T \wedge \omega_0 = dv_1 \wedge (m_{0v_1}^1)^T\omega_0$. Imposing the condition $d\wedge$ on (5.3) and using the equation itself we need: $-(m_0^1)^T\omega_0 \wedge (m_{0v_1}^1)^T\omega_0 + 2z(m_0^1)^T d\wedge\omega_0 = 0$, or using (4.6): $(N_0^0)^T(2zm_0^1 + m_0^1 \times m_{0v_1}^1)(N_0^0)^T(\omega_0 \times \wedge\omega_0) = 0$. Therefore the complete integrability is equivalent to (3.18). We get the transformation B_z' for the other ruling family by replacing (m_0^1, v_1) with $(m_0'^1, u_1)$. Since the distributions $\mathcal{D}$, $\mathcal{D}'$ reflect in Tx_0^0, the rolled distributions $(R_0, t_0)\mathcal{D}$, $(R_0, t_0)\mathcal{D}'$ reflect in Tx^0, so $B_z'(x^0)$ is just $B_z(x^0)$ when x_0^0 rolls on the other face of x^0. This geometric observation can be confirmed analytically if one keeps account of the change of ω_0 to $\omega_0' := -\omega_0 - 2N_0^0 \times dN_0^0$ when we roll x_0^0 on the other face of x^0 and of (3.26). Thus

$$\begin{aligned} &(m_0^1)^T\omega_0 + 2z dv_1 = 0 \;\Leftrightarrow\; (m_0'^1)^T\omega_0' + 2z du_1 = 0 \;\Leftrightarrow\; B_z \text{ transformation,} \\ &(m_0^1)^T\omega_0' + 2z dv_1 = 0 \;\Leftrightarrow\; (m_0'^1)^T\omega_0 + 2z du_1 = 0 \;\Leftrightarrow\; B_z' \text{ transformation.} \end{aligned} \tag{5.4}$$

Note that to obtain (3.18) as the complete integrability of (5.3) we just used $m_0^1 = f(z, v_1) + g(z, v_1) \times x_0^0$. The condition that the vectors f, g satisfy (3.18) imposes certain restrictions on the geometry of x_0^0 (for example if f, g are rational functions in v_1, then we have algebraic restriction on the geometry of x_0^0, which may be over-determined and prevent its existence); also equation (5.3) loses its geometric meaning.

Using $R_0^{-1}dx^1 = dx_z^1 + \omega_0 \times V_0^1$ and (5.2) we have $R_0^{-1}dx^1 = dx_z^1 - 2(x_{zv_1}^1)^T N_0^0 N_0^0 dv_1$ and the ACPIA becomes: $|dx_z^1 - 2(x_{zv_1}^1)^T N_0^0 N_0^0 dv_1|^2 = |dx_0^1|^2$, which boils down the first relation of (3.12). Note that if we conversely impose the ACPIA, then we get (5.3) (or its correspondent for the other ruling family).

We can try to prescribe the function $v_1(u_0, v_0)$ and solve (5.3) for flat connection forms ω_0. If $\frac{\omega_0}{\mathcal{B}_0} =: (\omega_{0u_0}^{u_0} x_{0u_0}^0 - \omega_{0u_0}^{v_0} x_{0v_0}^0)du_0 + (\omega_{0v_0}^{u_0} x_{0u_0}^0 - \omega_{0v_0}^{v_0} x_{0v_0}^0)dv_0$, then $0 = \omega_0 \times \wedge dx_0^0$ is equivalent to $\omega_{0u_0}^{u_0} = \omega_{0v_0}^{v_0}$ and the first line of (5.4) becomes:

$$\begin{bmatrix} v_{1u_0} \\ v_{1v_0} \end{bmatrix} = \begin{bmatrix} \omega_{0u_0}^{v_0} & \omega_{0u_0}^{u_0} \\ \omega_{0u_0}^{u_0} & \omega_{0v_0}^{u_0} \end{bmatrix} \begin{bmatrix} \frac{\Delta^+}{2z} \\ \frac{\Delta^-}{2z} \end{bmatrix}; \begin{bmatrix} u_{1u_0} \\ u_{1v_0} \end{bmatrix} = \begin{bmatrix} -\omega_{0u_0}^{v_0} & -(\omega_{0u_0}^{u_0} + 2\mathcal{N}_0) \\ -(\omega_{0u_0}^{u_0} + 2\mathcal{N}_0) & -\omega_{0v_0}^{u_0} \end{bmatrix} \begin{bmatrix} \frac{\Delta'^+}{2z} \\ \frac{\Delta'^-}{2z} \end{bmatrix}. \tag{5.5}$$

With $\rho_0^2 := \frac{-1}{K(x_0^0)} = \mathcal{A}^2|\hat{N}_0^0|^4,\ g_0^0 := \frac{\det(|dx_0^0|^2)}{(du_0\wedge dv_0)^2} = \frac{4\mathcal{A}^2|\hat{N}_0^0|^2}{\mathcal{B}_0^2}$ we have $\mathcal{N}_0 = \frac{1}{\mathcal{B}_0\rho_0}$ and the Christoffel symbols of x_0^0 are $\Gamma_{22}^1 = \Gamma_{11}^2 = 0,\ -\Gamma_{11}^1 = \Gamma_{22}^2 = \partial_{u_0}\log\mathcal{B}_0,\ \Gamma_{12}^2 = \partial_{u_0}\log\sqrt{g_0^0} - \Gamma_{11}^1 = \partial_{u_0}\log\sqrt{\rho_0},\ \Gamma_{12}^1 = \partial_{v_0}\log\sqrt{g_0^0} - \Gamma_{22}^2 = \partial_{v_0}\log\sqrt{\rho_0}$. Keeping account of (4.8), the condition $d\wedge\omega_0 + \frac{1}{2}\omega_0\times\wedge\omega_0 = 0$ becomes:

$$(\omega_{0u_0}^{u_0})^2 + 2\mathcal{N}_0\omega_{0u_0}^{u_0} = \omega_{0v_0}^{u_0}\omega_{0u_0}^{v_0},$$
$$\partial_{v_0}\omega_{0u_0}^{v_0} = \omega_{0u_0}^{u_0}\partial_{u_0}\log\frac{\omega_{0u_0}^{u_0}}{\mathcal{N}_0},\ \partial_{u_0}\omega_{0v_0}^{u_0} = \omega_{0u_0}^{u_0}\partial_{v_0}\log\frac{\omega_{0u_0}^{u_0}}{\mathcal{N}_0} \tag{5.6}$$

(the difference of the GCM equations for $x^0,\ x_0^0$). In particular if $v_{1u_0} = 0$, then $\omega_{0u_0}^{u_0} = \omega_{0u_0}^{v_0} = R_{0u_0} = t_{0u_0} = 0$ and $x^0 = (R_0(v_0), t_0(v_0))x_0^0$ is a ruled surface, in which case $\omega_{0v_0}^{u_0} = \varphi_0(v_0)$, $\varphi_0(v_0) := \frac{2zv_{1v_0}}{\Delta^-}$. The nonzero diagonal term of the second fundamental form of x^0 is $\mathcal{B}_0\sqrt{g_0^0}\varphi_0(v_0)$. Using (5.9) we see that x^1 is also ruled with rulings corresponding to $x_{0u_1}^1$. Therefore as we roll a quadric on a ruled applicable surface, the ruling families of a confocal quadric will generate congruences decomposable in ruled surfaces applicable to the initial quadric. Note that since Δ^+ is linear in v_1 for IQWC (3.9), if x^0 is ruled with rulings corresponding to $x_{0v_0}^0$, then its B_z transforms can be found by quadratures only (therefore its B_z' transforms can also be found by quadratures only).

Using first equations of (5.5) and (5.6) we get:

$$\omega_{0u_0}^{u_0} = \frac{2z^2v_{1u_0}v_{1v_0}}{\mathcal{N}_0\Delta^+\Delta^- + zv_{1u_0}\Delta^+ + zv_{1v_0}\Delta^-},\ \omega_{0u_0}^{v_0} = \frac{2zv_{1u_0}(\mathcal{N}_0\Delta^- + zv_{1u_0})}{\mathcal{N}_0\Delta^+\Delta^- + zv_{1u_0}\Delta^+ + zv_{1v_0}\Delta^-},$$
$$\omega_{0v_0}^{u_0} = \frac{2zv_{1v_0}(\mathcal{N}_0\Delta^+ + zv_{1v_0})}{\mathcal{N}_0\Delta^+\Delta^- + zv_{1u_0}\Delta^+ + zv_{1v_0}\Delta^-}. \tag{5.7}$$

From the compatibility condition $(v_{1u_0})_{v_0} = (v_{1v_0})_{u_0}$ applied to the first equation of (5.5), using $((u_1, \Delta'^-, \Delta'^+) \leftrightarrow (v_1, \Delta^-, \Delta^+))\circ$ the first relation of (3.27) and the first relation of (5.6) (which ω_0 as defined by (5.7) automatically satisfies) we get the fact that the last two relations of (5.6) are linearly dependent with coefficients $-\Delta^+,\ \Delta^-$; thus either of them must be satisfied. The second equation of (5.6) becomes: $(\mathcal{N}_0\Delta^- + zv_{1u_0})(\log\frac{v_{1u_0}(\mathcal{N}_0\Delta^- + zv_{1u_0})}{\mathcal{N}_0\Delta^+\Delta^- + zv_{1u_0}\Delta^+ + zv_{1v_0}\Delta^-})_{v_0} = zv_{1v_0}(\log\frac{v_{1u_0}v_{1v_0}}{\mathcal{N}_0(\mathcal{N}_0\Delta^+\Delta^- + zv_{1u_0}\Delta^+ + zv_{1v_0}\Delta^-)})_{u_0}$, or using again $((u_1, \Delta'^-, \Delta'^+) \leftrightarrow (v_1, \Delta^-, \Delta^+))\circ$ the first relation of (3.27):

$$z(\mathcal{N}_0\Delta^+ + zv_{1v_0})(\log\frac{v_{1u_0}}{v_{1v_0}})_{u_0} + z(\mathcal{N}_0\Delta^- + zv_{1u_0})(\log\frac{v_{1v_0}}{v_{1u_0}})_{v_0} - \mathcal{N}_0^2\Delta^+\Delta^-\frac{v_{1u_0}v_0}{v_{1u_0}v_{1v_0}} -$$
$$z^2(v_{1v_0}(\log\mathcal{N}_0)_{u_0} + v_{1u_0}(\log\mathcal{N}_0)_{v_0}) - \mathcal{N}_0\Delta^+\Delta^-(\frac{z}{\Delta^-}\partial_{u_0}\log(\mathcal{N}_0\Delta^+) -$$
$$\mathcal{N}_0\partial_{v_1}\log\Delta^+ + \frac{z}{\Delta^-}(\log\mathcal{N}_0)_{u_0} + \frac{z}{\Delta^+}(\log\mathcal{N}_0)_{v_0}) = 0. \tag{5.8}$$

Therefore v_1 must satisfy a second order PDE linear in the second order terms and having the symmetry $(u_0, \Delta^+) \leftrightarrow (v_0, \Delta^-)$ (because of this the third equation of (5.6) and (5.8) are equivalent). Since (5.6) is valid with $(\omega_{0u_0}^{v_0}, \omega_{0u_0}^{u_0}, \omega_{0v_0}^{u_0}) \leftrightarrow (-\omega_{0u_0}^{v_0}, -(\omega_{0u_0}^{u_0} + 2M_0), -\omega_{0v_0}^{u_0})$, (5.8) is valid for $(u_1, \Delta'^-, \Delta'^+) \leftrightarrow (v_1, \Delta^-, \Delta^+)$.

5.2. **Inversion of the Bäcklund transformation and the Weingarten congruence.**
Let $x_0^0,\ x_0^1$ be in the TC $(V_0^1)^T N_0^0 = 0$ and v_1 be given by (5.3), where $(x^0, dx^0) = (R_0, t_0)(x_0^0, dx_0^0)$.

If we roll x^0 on x_0^0, then the tangent space of $x^1 := (R_0, x^0)V_0^1$ will be applied to the space at x_z^1 generated by $x_{zu_1}^1,\ V_0^1$. Further applying (R_0^1, t_0^1), this space will be applied to $T_{x_0^1}x_0$ with $(R_0^1, t_0^1)(x_z^1, x_{zu_1}^1, V_0^1) = (x_0^1, x_{0u_1}^1, -V_1^0)$. If $(x^1, dx^1) = (R_1, t_1)(x_0^1, dx_0^1)$ is the rolling of x_0^1 on x^1 with $\det(R_1^{-1}R_0) = 1$, then we have just proved $(R_0^1, t_0^1)(R_0, t_0)^{-1}(x^1, dx^1) = (R_1, t_1)^{-1}(x^1, dx^1)$, so

$$(R_0^1, t_0^1) = (R_1, t_1)^{-1}(R_0, t_0). \tag{5.9}$$

If we make the ansatz $x^0 = x_0^0$, so $(R_0, t_0) = (I_3, 0)$, $\omega_0^0 = 0$, $v_1 = \text{const}$ or $(R_0, t_0) = (I_3 - 2N_0^0(N_0^0)^T, 2N_0^0(N_0^0)^T x_0^0)$, $\omega_0^0 = -2N_0^0 \times dN_0^0$, $v_1 = \text{const}$, then we get in a natural geometric way all interesting algebraic identities of the static part (including the explanation for the necessity of the existence of the RMPIA).

Moreover, $x^0 = (R_0, x^1)(-V_0^1) = (R_0, x^1)(R_0^1)^{-1}V_1^0 = (R_1, x^1)V_1^0$ and thus x^0 reveals itself as a B_z transform of x^1. Therefore we can find all the B transforms $B_z(x^1)$ for fixed z only by two quadratures and further algebraic computations. We shall see later that we don't need the quadratures once we know the B transforms $B_z(x^0)$ for all z as one can take a derivative in the spectral parameter z in a L'Hospital situation of the BPT (although Sym's formula was not known to the classical geometers, tricks similar to it were). Also equations (5.5)-(5.8) admit the symmetry $(u_0, v_0, \Delta^+, \Delta'^-, \omega_0, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \Delta'^-, \Delta^+, \omega_1, \mathcal{N}_1)$. We shall now provide an analytic confirmation of the inversion of the B transformation, equivalently of the fact that $((u_0, v_0, \Delta^+, \Delta'^-, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \Delta'^-, \Delta^+, \mathcal{N}_1))\circ(5.8)$ is still valid. This analytic confirmation will play an important rôle in the discussion of the B_∞ transformation, since the easy geometric justification will not be available.

We have $\begin{bmatrix} u_{0u_1} & u_{0v_1} \\ v_{0u_1} & v_{0v_1} \end{bmatrix} = \frac{1}{J}\begin{bmatrix} v_{1v_0} & -u_{1v_0} \\ -v_{1u_0} & u_{1u_0} \end{bmatrix} = \frac{1}{J}\begin{bmatrix} v_{1v_0} & \frac{\Delta'^-}{\Delta^-}(\frac{\mathcal{N}_0\Delta^+}{z} + v_{1v_0}) \\ -v_{1u_0} & -\frac{\Delta'^-}{\Delta^-}(\frac{\mathcal{N}_0\Delta^-}{z} + v_{1u_0}) \end{bmatrix}$, where $J := \frac{du_1 \wedge dv_1}{du_0 \wedge dv_0} = \frac{\mathcal{N}_0\Delta'^+}{z}v_{1u_0} - \frac{\mathcal{N}_0\Delta'^-}{z}v_{1v_0}$ (assume $J \neq 0$).

Now $((u_0, v_0, \Delta^+, \Delta'^-, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \Delta'^-, \Delta^+, \mathcal{N}_1))\circ(5.8)$ becomes

$z(\mathcal{N}_1\Delta'^- + zv_{0v_1})((\log\frac{v_{0u_1}}{v_{0v_1}})_{u_0}u_{0u_1} + (\log\frac{v_{0u_1}}{v_{0v_1}})_{v_0}v_{0u_1}) + z(\mathcal{N}_1\Delta^- + zv_{0u_1})((\log\frac{v_{0v_1}}{v_{0u_1}})_{u_0}u_{0v_1} + (\log\frac{v_{0v_1}}{v_{0u_1}})_{v_0}v_{0v_1}) - \mathcal{N}_1^2\Delta'^-\Delta^-((\log v_{0u_1})_{v_0} + \frac{u_{0v_1}}{v_{0v_1}}(\log v_{0u_1})_{u_0}) - z^2(v_{0v_1}(\log\mathcal{N}_1)_{u_1} + v_{0u_1}(\log\mathcal{N}_1)_{v_1}) - \mathcal{N}_1\Delta'^-\Delta^-(\frac{z}{\Delta^-}\partial_{u_1}\log(\mathcal{N}_1\Delta'^-) - \mathcal{N}_1\partial_{v_0}\log\Delta'^- + \frac{z}{\Delta^-}(\log\mathcal{N}_1)_{u_1} + \frac{z}{\Delta'^-}\log(\mathcal{N}_1)_{v_1}) = 0$. Using (3.24), the coefficient of $(\log\frac{v_{0u_1}}{v_{0v_1}})_{u_0}$ becomes 0; the remaining part becomes

$0 = -\frac{zv_{1u_0}\mathcal{N}_0\mathcal{N}_1(\Delta^-)^2}{J(\mathcal{N}_0\Delta^- + zv_{1u_0})}(\frac{\mathcal{N}_0\Delta'^-}{zv_{1u_0}}(\partial_{v_0}\log(\mathcal{N}_0\Delta'^-) + u_{1v_0}\partial_{u_1}\log\Delta'^- - (\log v_{1u_0})_{v_0}) + \frac{\Delta'^-}{\Delta^-}(\partial_{v_0}\log\frac{\Delta'^-}{\Delta^-} + u_{1v_0}\partial_{u_1}\log\Delta'^- - v_{1v_0}\partial_{v_1}\log\Delta^-)) + \frac{\mathcal{N}_1^2\Delta'^-\Delta^-v_{1u_0}}{J}(\frac{\mathcal{N}_0\Delta'^+}{z}((\log\mathcal{N}_0)_{v_0} + u_{1v_0}\partial_{u_1}\log\Delta'^+) - \frac{\mathcal{N}_0\Delta'^-v_{1v_0}}{zv_{1u_0}}(\partial_{v_0}\log(\mathcal{N}_0\Delta'^-) + u_{1v_0}\partial_{u_1}\log\Delta'^- + (\log\frac{v_{1v_0}}{v_{1u_0}})_{v_0}) - \frac{\mathcal{N}_0\Delta^+ + zv_{1v_0}}{\mathcal{N}_0\Delta^- + zv_{1u_0}}(\frac{\mathcal{N}_0\Delta'^+}{z}(\partial_{u_0}\log(\mathcal{N}_0\Delta'^+) + u_{1u_0}\partial_{u_1}\log\Delta'^+) - \frac{\mathcal{N}_0\Delta'^-v_{1v_0}}{zv_{1u_0}}((\log\mathcal{N}_0)_{u_0} + u_{1u_0}\partial_{u_1}\log\Delta'^- - (\log\frac{v_{1u_0}}{v_{1v_0}})_{u_0}))) - \frac{z^2}{J}(u_{1u_0}(\log\mathcal{N}_1)_{u_1} - v_{1u_0}(\log\mathcal{N}_1)_{v_1}) - \mathcal{N}_1\Delta'^-\Delta^-(\frac{z}{\Delta^-}\partial_{u_1}\log(\mathcal{N}_1\Delta'^-) - \mathcal{N}_1\partial_{v_0}\log\Delta'^- + \frac{z}{\Delta^-}(\log\mathcal{N}_1)_{u_1} + \frac{z}{\Delta'^-}\log(\mathcal{N}_1)_{v_1})$

Comparing the coefficients of $(\log\frac{v_{1u_0}}{v_{1v_0}})_{u_0}$, $(\log\frac{v_{1v_0}}{v_{1u_0}})_{v_0}$, $\frac{v_{1u_0}v_0}{v_{1u_0}v_{1u_0}}$ we see that so far this is just (5.8) multiplied by $-\frac{\mathcal{N}_0\mathcal{N}_1^2\Delta^-(\Delta'^-)^2v_{1v_0}}{z^2J(\mathcal{N}_0\Delta^- + zv_{1u_0})}$. Replacing u_{1u_0}, u_{1v_0}, J with their values (note that the terms in the second parenthesis containing u_{1u_0}, u_{1v_0} cancel), the remaining part is the same multiple of (5.8) (this boils down to a quadratic polynomial in v_{1u_0}, v_{1v_0} being identically 0; no further expansion of derivatives is needed; one uses instead the algebraic identities (3.24), (3.27) and its versions under symmetries).

Another way to prove the inversion of the B transformation is to show that ω_1 as defined by $((u_0, v_0, \Delta^+, \Delta'^-, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \Delta'^-, \Delta^+, \mathcal{N}_1))\circ(5.7)$ satisfies the last two equations of $((u_0, v_0, \omega_0, \mathcal{N}_0) \leftrightarrow (u_1, v_1, \omega_1, \mathcal{N}_1))\circ(5.6)$ (the first equation is automatically satisfied); by symmetry it is enough to prove only the second one. The differential of x^1 depends on the shape (but not its derivatives) of x^0; a-priori the shape of x^1 depends also on the first derivatives of the shape of x^0, but it actually depends only on the shape of x^0. Using the GCM equations and the first derivatives of the Gauß equation of x^0, the required equation is algebraic in $u_0, v_0, u_1, v_1, \omega_{0u_0}^{v_0}, \omega_{0u_0}^{u_0}, \partial_{u_0}\omega_{0u_0}^{u_0}, \partial_{v_0}\omega_{0u_0}^{u_0}$; accounting the coefficients of the four terms involving the shape of x^0 and its derivatives they will be identically 0.

Note that we have provided another proof for the ACPIA by effectively finding the rolling. This proof is the one susceptible for generalization and the essence of the theory of deformations of quadrics revealed itself: take a surface $x_0 = x_0(u, v)$ in $\mathbb{C}^3$ and a 3-dimensional integrable distribution of facets $(p, P) = (p, P)(u, v, w)$, whose integrability condition depends only on the linear

element of x_0. If $x^0 \subset \mathbb{C}^3$ is applicable to $x_0^0 := x_0(u_0, v_0)$ with rolling $(x^0, dx^0) = (R_0, t_0)(x_0^0, dx_0^0)$, then the rolled distribution will still be integrable with leaves

$x^1 := (R_0(u_0, v_0), x^0(u_0, v_0))(p(u_0, v_0, w(u_0, v_0)) - x^0(u_0, v_0))$. If for each triple $f := (u, v, w)$ we have a triple of functions $\tilde{f} := (\tilde{u}, \tilde{v}, \tilde{w})$ of (u, v, w), a rigid motion $(R_{f\tilde{f}}, t_{f\tilde{f}})$ taking $(p, P)(u, v, w)$ to $(x_0, dx_0)(\tilde{u}, \tilde{v})$ and $(R_{f\tilde{f}} R_0^{-1} dx_1)(u_1, v_1) = dx_0(u_1, v_1)$, where

$(u_1, v_1) := (\tilde{u}(u_0, v_0, w(u_0, v_0)),\ \tilde{v}(u_0, v_0, w(u_0, v_0)))$, so by a change of coordinates (when (u_1, v_1) are functionally independent) we can consider the (u_1, v_1) coordinates on x^1, $x_0^1 := x_0(u_1, v_1)$, then x^1 is applicable to x_0^1 with explicit rolling. If further $(R_{f\tilde{f}}, t_{f\tilde{f}})$ takes $(x_0, dx_0)(u, v)$ to $(p, P)(\tilde{u}, \tilde{v}, \tilde{w})$, then the relationship between x^0, x^1 is symmetric. To find such $(R_{f\tilde{f}}, t_{f\tilde{f}})$ we need a correspondence between the given distribution of facets (p, P) and the distribution of facets (x_0, dx_0); therefore we need a submersion between the two distributions. In the case of quadrics this submersion factors through the leaves (a ruling family on the confocal quadric) and becomes the Ivory affinity. It is thus natural for this submersion to factor through the leaves of the initial distribution (when these are curves) to a diffeomorphism between the surface described by these curves and x_0. In this case $(R_{f\tilde{f}}, t_{f\tilde{f}})$ are not uniquely determined and it is natural that the extra condition should require that it takes the tangent vector to the curve (leaf) to a certain vector of the tangent space of x_0 at the point corresponding under the diffeomorphism.

However, it may be the case that only the B transformation for quadrics satisfies such requirements. Bianchi (for example in ([5], Vol **4**,(173)) tried generalizations (natural from a geometric point of view) of various aspects of the B transformation for quadrics, but most of such configurations arise only for quadrics.

According to R. Calapso in § 5.17 of the introduction to Bianchi ([5], Vol **4**, part 1), it may be an interesting problem to find the surfaces for which a theory of deformations similar to that for quadrics can be developed.

We are interested in the case when x^1 degenerates to a curve, that is $0 = dx^1 \times \wedge dx^1$, or using (5.9): $du_1 \wedge dv_1 = 0$, or replacing du_1 from (3.26): $v_{1u_0}\Delta^+ - v_{1v_0}\Delta^- = 0$, which must be adjoined to (5.8). In general $v_{1u_0} v_{1v_0} = 0$ is not possible, so we need $v_{1u_0}, v_{1v_0} \neq 0$. Using (5.8) we thus need:

$$\mathcal{N}_0^2 (\frac{\Delta^-}{v_{1u_0}})^2 v_{1u_0v_0} + (\mathcal{N}_0\Delta^+\Delta^- + z v_{1u_0}\Delta^+)(\frac{z}{\Delta^-}\partial_{u_0}\log(\mathcal{N}_0\Delta^+) + \frac{z}{\Delta^+}\partial_{v_0}\log(\mathcal{N}_0\Delta^-)) +$$

$$\mathcal{N}_0\Delta^+\Delta^-(\frac{z}{\Delta^-}\partial_{u_0}\log(\mathcal{N}_0\Delta^+) - \mathcal{N}_0\partial_{v_1}\log\Delta^+) = 0,\ v_{1u_0}\Delta^+ - v_{1v_0}\Delta^- = 0. \tag{5.10}$$

Differentiating the second relation of (5.10) wrt u_0 and v_0 and using the first one we get $v_{1u_0u_0}$, $v_{1v_0v_0}$, $v_{1u_0v_0}$ in terms of v_1, v_{1u_0} and so on; therefore all derivatives of v_1 can be written in function of v_1, v_{1u_0}, so in general (formally) such a v_1 depends on two constants besides the parameter z. An application of Cauchy-Kovalewskaia assures in general the existence of such solutions. Because of the first relation of (3.27) all derivatives of v_1 admit the symmetry $(u_0, \Delta^+) \leftrightarrow (v_0, \Delta^-)$. Once the solution v_1 is found, (5.7) gives the shape of the 1-solitons. Note that the family of 1-solitons may depend up to rigid motions on less than three constants; for example Dini's CGC -1 helicoids depend up to rigid motions only on a constant (instead of three; in fact the axis of the tractrix as a limiting case of a Dini helicoid is a good picture of it as a CGC -1 surface). Note however that Serret's ruled surfaces generated by taking isotropic directions in tangent planes along asymptotes of Dini's helicoids still provide another 1-dimensional family of ruled imaginary 1-solitons (again they cannot be seen at the level of the sine-Gordon equation); this must be also the case for general quadrics (by Chieffi's), but at that level one cannot see the analytic behavior. If we take according to Rogers-Schieff ([26],(1.85)) the Dini helicoids $x = x(u, v) := \cosh(\frac{u - v\cos(\zeta)}{\sin(\zeta)})^{-1}\sin(\zeta)(\cos(v)e_1 + \sin(v)e_2) + (u - \sin(\zeta)\tanh(\frac{u - v\cos(\zeta)}{\sin(\zeta)}))e_3$, then it becomes clear that for $\zeta = 0$ we obtain the e_3-axis; for $u = \frac{c+s}{2}$, $v = \epsilon_1\frac{c-s}{2}$, $\epsilon_1 := \pm 1$, c constant we have ruled CGC -1 surfaces $S(s, t) := x + t(\partial_u x + i\epsilon_2\sinh(\frac{u - v\cos(\zeta)}{\sin(\zeta)})\partial_v x)$, $\epsilon_2 := \pm 1$; on Rogers-Schieff ([26],

fig 1.5 pag 36) asymptotes are obtained by drawing diagonals of the rectangles obtained by lines of curvature u, v.

According to a theorem of Ribacour $x^0,\ x^1$ are the focal surfaces of a W congruence iff $K(x^0)K(x^1) = \frac{\sin^4(\beta)}{d^4}$, where β (respectively d) is the angle (the distance) between the tangent planes at the corresponding points (the corresponding points). This is basically a statement about tangent curves (an asymptote on x^0 (x^1) lies in the osculating bundle of its corresponding asymptote on x^1 (x^0) and the relation between their torsions becomes via Enneper's formula a relation between the curvatures of x^0 and x^1; see Bianchi ([5],Vol **5**,(117))). In our case that becomes: $K(x^0)K(x^1) = \frac{|m_0^1 \times N_0^0|^4}{|m_0^1|^4|V_0^1|^4} = \frac{((N_0^0)^T x^1_{z u_1})^4}{|x^1_{z u_1} \times V_0^1|^4} =^{(3.22)} \frac{1}{\mathcal{A}^4 |\hat{N}_0^0|^4 |\hat{N}_0^1|^4}$, which is true since $K(x_0^j) = \frac{-1}{\mathcal{A}^2 |\hat{N}_0^j|^4},\ j = 0, 1.$

Bianchi (122) has a beautiful geometric argument: using Cheffi's result it is enough to prove the W congruence property only for ruled surfaces applicable to quadrics, in which case the rulings correspond under the B transformation (use (5.9)). Thus asymptotes of one system (rulings) correspond, so Ribacour's theorem applies and therefore the asymptotes of the other system correspond. Note that although the rulings are plane curves, they do not have zero torsion because the Frenet trihedral is not defined; therefore the torsion is undefined, or equivalently arbitrarily defined so as to suit our purposes.

Again the geometric picture gives the correct answer: the curvature of a space curve measures its bending in the normal direction (the normal is naturally defined from a geometric point of view in the osculating plane of the curve) and the torsion measures its twisting (tendency to leave its osculating plane). Since a line does not admit a well defined osculating plane, the curvature and torsion of a line cannot be defined from a geometric point of view, or equivalently they can be arbitrarily defined (usually as a limiting process) so as to suit our purposes. For example the family of helices $x = x(s) = \lambda \cos(\frac{s}{\sqrt{\lambda a}})e_1 + \lambda \sin(\frac{s}{\sqrt{\lambda a}})e_2 + \frac{\sqrt{\lambda(a-\lambda)}}{\sqrt{\lambda a}} s e_3$ has constant curvature $\frac{1}{a}$, so we get the curvature of the line se_3 to be $\frac{1}{a}$ for $\lambda \to 0$ and the tangent surfaces of the family (all deformations one of the other) further contains as a deformation the tangent bundle of the line se_3. One can perform an experiment for the real plane region between concentric circles of radii $a,\ a + \epsilon$ with a cut: this can be tightly coiled around itself; as $\epsilon \to 0$ we get the required line*. The same experiment can be done with any one-sided tubular neighborhood of a plane curve and thus a line can have any curvature. Of course such a statement is ridiculous, but statements and structures whose validity depend only on the linear element of any of such tangent surfaces may remain valid and interesting in the limit.

Because of the correspondence of asymptotes on $x^0,\ x^1$ we can reduce (similarly to Lie's approach for CGC surfaces) the B transformation of deformations of quadrics to the B transformation of their asymptotes (if a curve corresponds with preserved arc-length with a curve on a quadric and is an asymptote on a deformation of that quadric, then its torsion is determined by Enneper's formula) and conversely we can extend B transformations of such curves to B transformations of deformations of quadrics. Similarly we can extend B transformations (of deformations of quadrics) to B transformations of triply conjugate families of surfaces (and containing a family of deformations of quadrics). In fact according to Bianchi ([5],Vol **4**,(143)) the family $x^1(c) := x^1(u_1(u_0, v_0, c), v_1(u_0, v_0, c)) = B_{a_1}(x^0)$ of singular B transforms of x^0 gives a family of surfaces (all applicable to that same quadric) which can be completed to a triply conjugate system

* Again this discussion took place in Calculus III; since students did not a-priori see the fact that the projection of helices on the xy plane become circles with smaller radius I saw myself put into the position of further deforming the piece of paper so it becomes clear that the projection becomes a point in the limit (I was not prepared to attack the point picture, but my students forced me to); thus they spelled out the truth that the circles become smaller and in the limit we obtain a point. I then asked *the natural question* about the curvature of the line without being prepared with the correct answer; when a student spelled out the truth that the line has curvature $\frac{1}{a}$ I knew that he was right (Bianchi proved him right) but that the discussion went longer than I was prepared and I had to clarify it on the next meeting by trimming the piece of paper to a smaller one-sided tubular neighborhood of the given circle, because the piece of paper just with the disk removed and a cut did not leave me coil it tightly and just pulling it apart (as one would do with a coiled wire) would break it.

of surfaces. All such deformations of the considered quadric have as the considered conjugate system the permanent conjugate system, so for CGC surfaces we obtain triply orthogonal family of surfaces containing a family of CGC surfaces.

Since $s_0 := x^0, s_3 := x^1$ are the focal surfaces of a W congruence, there exists $s_1 := r_1 R_0 m_0^1$ infinitesimal deformation of x^0 : $0 = 2zr_1^{-1}(ds_1^T dx^0 + (dx^0)^T ds_1) = 2z(d(\log r_1)(m_0^1)^T dx_0^0 + (dx_0^0)^T m_0^1 d(\log r_1) + (dm_0^1)^T dx_0^0 + (dx_0^0)^T dm_0^1 - 2(m_0^1)^T(\omega_0 \times dx_0^0)) = (2zd(\log r_1) - (m_{0v_1}^1)^T\omega_0)(m_0^1)^T dx_0^0 + (dx_0^0)^T m_0^1(2zd(\log r_1) - (m_{0v_1}^1)^T\omega_0)$ (use $(dm_0^1)^T dx_0^0 + (dx_0^0)^T dm_0^1 = dv_1(m_{0v_1}^1)^T dx_0^0 + (dx_0^0)^T m_{0v_1}^1 dv_1$, (3.18) and $0 = (dx_0^0)^T(\omega_0 \times dx_0^0)$). Therefore r_1 should be given by $2zd(\log r_1) = (m_{0v_1}^1)^T\omega_0$, which must be integrable. But differentiating (3.18) wrt v_1 we get $(N_0^0)^T(2czm_{0v_1}^1 + m_0^1 \times m_{0v_1 v_1}^1) = 0$, which ensures $d\wedge(m_{0v_1}^1)^T\omega_0 = 0$. Thus we can say that r_1^{-1} is an infinitesimal deformation of v_1 : $r_1^{-1} = \delta v_1$, or $r_1^{-1} = \partial_c v_1$ if $v_1 = v_1(z, u_0, v_0, c)$. Therefore s_1 and Darboux's 12 surfaces can be generated by a quadrature (or no quadrature if we know the general solution of (5.3) for fixed z). We have $r_1^{-1}R_0^{-1}ds_1 = dx_0^0 \times (\mathcal{B}_1 x_{zu_1}^1 - \omega_0 \lrcorner \frac{(\nabla x_0^0)^T(2zm_0^1 + m_0^1 \times m_{0v_1}^1)}{2z})$, so $s_{-1} := r_1 R_0(-\mathcal{B}_1 x_{zu_1}^1 + \omega_0 \lrcorner \frac{(\nabla x_0^0)^T(2zm_0^1 + m_0^1 \times m_{0v_1}^1)}{2z})$ is the prwrtps of $s_2 := r_2 R_1 m_1^0$ and $0 = 1 + s_{-1}^T s_2 = 1 + r_1 r_2 \Delta^-$ (use (3.22) and the first equation of (3.23) for $R_1 m_1^0$). Thus

$$1 + r_1 r_2 \Delta^- = 0. \tag{5.11}$$

5.3. **Bianchi Permutability Theorem and moving Möbius configurations.**
Let $B_{z_1}(x^0) = x^1 = B_{z_2}(x^3)$. The BPT states that there is $B_{z_2}(x^0) = x^2 = B_{z_1}(x^3)$; moreover $x^3 = (B_{z_1} \circ B_{z_2})(x^0) = (B_{z_2} \circ B_{z_1})(x^0)$ is obtained from (x^0, x^1, x^2) only by algebraic computations.



Let $(R_j, t_j)(x_0^j, dx_0^j) = (x^j, dx^j),\ j = 0, 1, 3$ be the rollings known so far. One can find x_0^2 according to the algebraic computations of the SITC; the sought surface $x^2 := (R_0, x^0)V_0^2 = (R_3, x^3)V_3^2$ is recovered using only algebraic computations (we need to take derivatives just to find R_0). From the cross ratio property we get $x^3 = x^0$ for $z_1 = z_2$, so the Permutability Theorem does not a-priori get rid of the quadratures for the iteration of the transformations B_z with the same z. However an application of L'Hospital will suffice once we know all $B_z(x^0)$ (see Bianchi (122)).

We have $x^2 = (R_1, t_1)(R_3^0, t_3^0)x_0^2 = (R_2, t_2)x_0^2,\ R_2 := R_1 R_3^0,\ t_2 := (R_1, t_1)t_3^0$ and we need to prove $dx^2 = R_2 dx_0^2$, that is $R_1^{-1}dR_1 R_0^1 V_1^2 + d(R_3^0, t_3^0)x_0^2 = 0$. Using (5.3) this becomes an algebraic relation between $x_0^j,\ j = 0, ..., 3$ and linear in the second fundamental form of x^1. Using (5.5) for $\frac{\omega_1}{\mathcal{B}_1} =: (\omega_{1u_1}^{u_1} x_{0u_1}^1 - \omega_{1u_1}^{v_1} x_{0v_1}^1)du_1 + (\omega_{1v_2 1}^{u_1} x_{0u_1}^1 - \omega_{1v_1}^{v_1} x_{0v_1}^1)dv_1$ we thus need the algebraic relations:

$$\begin{aligned}
\mathcal{B}_1 x_{0u_1}^1 \times R_0^1 V_1^2 + (\frac{\Delta^-(z_1, v_0, v_1)}{2z_1}(R_3^0, t_3^0)_{v_0} + \frac{\Delta^-(z_2, v_1, v_3)}{2z_2}(R_3^0, t_3^0)_{v_3})x_0^2 = 0,\\
-\mathcal{B}_1 x_{0v_1}^1 \times R_0^1 V_1^2 + (\frac{\Delta^+(z_1, v_0, u_1)}{2z_1}(R_3^0, t_3^0)_{v_0} + \frac{\Delta^+(z_2, u_1, v_3)}{2z_2}(R_3^0, t_3^0)_{v_3})x_0^2 = 0.
\end{aligned} \tag{5.12}$$

If $x^j,\ j = 0, ..., 3$ are ruled with rulings corresponding to $x_{0u_j}^j,\ j = 0, ..., 3$ only the first relation of (5.12) appears when the rulings correspond to $x_{0u_0}^0,\ x_{0v_1}^1,\ x_{0v_2}^2,\ x_{0u_3}^3$ only the second one appears.

Therefore we have reduced the proof of the BPT to the case of ruled surfaces (Bianchi in (122) applies the inverse of Chieffi's result for deformations of quadrics). Suppose $x^j,\ j = 0, ..., 3$ are ruled (with rulings corresponding to $x_{0u_j}^j,\ j = 0, ..., 3$) satisfying the BPT. Using (5.5) we have $2z_1 dv_0 = \Delta^-(z_1, v_0, v_1)\varphi_1(v_1)dv_1,\ 2z_2 dv_3 = \Delta^-(z_2, v_1, v_3)\varphi_1(v_1)dv_1,\ 2z_2 dv_0 = \Delta^-(z_2, v_0, v_2)\varphi_2(v_2)dv_2,$

$2z_1dv_3=\Delta^-(z_1,v_2,v_3)\varphi_2(v_2)dv_2$, so $\frac{z_2\Delta^-(z_1,v_0,v_1)}{z_1\Delta^-(z_2,v_1,v_3)}=\frac{dv_0}{dv_3}=\frac{z_1\Delta^-(z_2,v_0,v_2)}{z_2\Delta^-(z_1,v_2,v_3)}$. Therefore

$$\begin{aligned}&\frac{z_2\Delta^-(z_1,v_0,v_1)}{z_1\Delta^-(z_2,v_1,v_3)}=\frac{z_1\Delta^-(z_2,v_0,v_2)}{z_2\Delta^-(z_1,v_2,v_3)},\\ &\frac{z_2\Delta^+(z_1,v_0,u_1)}{z_1\Delta^+(z_2,u_1,v_3)}=\frac{z_1\Delta^+(z_2,v_0,u_2)}{z_2\Delta^+(z_1,u_2,v_3)}\end{aligned}\tag{5.13}$$

are necessary algebraic conditions for the BPT (the second one is similarly obtained, when the rulings correspond to $x^0_{0u_0},\ x^1_{0v_1},\ x^2_{0v_2},\ x^3_{0u_3}$). Applying to (3.32) the symmetry $(0\leftrightarrow 1,2\leftrightarrow 3)$, multiplying the two relations and using (3.23) we get the first relation of (5.13) (the second one is similar has to do with the cross-ratio of the intersections of the rulings $x^1_{0v_1},\ R^0_3x^2_{2v_2}$ with the line $[x^0_{z_1}\ x^3_{z_2}]$ or we can use the last relation of (3.23)).

Conversely they are also sufficient for ruled surfaces (we shall prove the first one; the second one is similar). If we are in the conditions of the Permutability Theorem and we denote $\varphi_2(v_2):=\frac{2z_2}{\Delta^-(z_2,v_0,v_2)}\frac{dv_0}{dv_2}=\frac{2z_1}{\Delta^-(z_1,v_2,v_3)}\frac{dv_3}{dv_2}$, then the ruled surface x'^2 with the nonzero diagonal entry of the second fundamental form given by $\mathcal{B}_2\sqrt{g_0^2}\varphi_2(v_2)$ is a B_{z_2} (B_{z_1}) transform of $x'^1,\ (x'^3)$ (which is applicable to $x^0_0,\ (x^3_0)$).

Let $(x^j,dx^j)=(R_j,t_j)(x^j_0,dx^j_0),\ j=0,1,3,\ (x'^k,dx'^k)=(R'_k,t'_k)(x^k_0,dx^k_0),\ k=0,2,3$ be the rollings known so far. Then using (5.9) and (3.30) we have $(R_0,t_0)(R'_0,t'_0)^{-1}=$
$(R_1,t_1)(R^0_3,t^0_3)(R'_2,t'_2)^{-1}=(R_3,t_3)(R'_3,t'_3)^{-1}$. If $(R_0,t_0)(R'_0,t'_0)^{-1}$ is a rigid motion, then $x^2:=(R_0,t_0)(R'_0,t'_0)^{-1}x'^2$ will satisfy the conditions of the BPT. For this we need the second fundamental forms of $x^0,\ x'^0$ to be the same, or

$$v_0=v_0(v_1),\ v_3=v_3(v_1),\ \frac{dv_3}{dv_0}=\frac{z_1\Delta^-(z_2,v_1,v_3)}{z_2\Delta^-(z_1,v_0,v_1)}\Rightarrow\frac{dv_2}{dv_1}=\frac{z_1\Delta^-(z_2,v_0,v_2)}{z_2\Delta^-(z_1,v_0,v_1)}.\tag{5.14}$$

While it seems that we are finished if we apply the symmetry $(0\leftrightarrow 1,2\leftrightarrow 3)$ to the SITC in the previous conditions, a reduction to an algebraic identity is safer. If the lhs of (3.33)-(3.35) are $\phi=\phi(z_1,z_2,v_0,v_1,v_2,v_3)$ and we differentiate $\phi=0$ with the constraints $v_0=v_0(v_1),\ v_3=v_3(v_1),\ \frac{dv_3}{dv_0}=\frac{z_1\Delta^-(z_2,v_1,v_3)}{z_2\Delta^-(z_1,v_0,v_1)}$, then we get $\frac{dv_0}{dv_1}[\phi_{v_0}+\phi_{v_3}\frac{dv_3}{dv_0}]+\phi_{v_1}+\phi_{v_2}\frac{dv_2}{dv_1}=0$. If (5.14) were to be true, then this last relation becomes a linear function (with coefficients algebraic functions of $v_j,\ j=0,...,3,z_1,z_2$) of the a-priori arbitrary function $\frac{dv_0}{dv_1}(v_1)$ being 0, which is true if $\phi_{v_0}+\phi_{v_3}\frac{dv_3}{dv_0}=\phi_{v_1}+\phi_{v_2}\frac{dv_2}{dv_1}=0$. Therefore we need the algebraic relations $\frac{\phi_{v_0}}{\phi_{v_3}}=-\frac{z_1\Delta^-(z_2,v_1,v_3)}{z_2\Delta^-(z_1,v_0,v_1)},\ \frac{\phi_{v_1}}{\phi_{v_2}}=-\frac{z_1\Delta^-(z_2,v_0,v_2)}{z_2\Delta^-(z_1,v_0,v_1)}$. If either is proven, then by applying the symmetry $(0\leftrightarrow 1,2\leftrightarrow 3)$ (which preserves ϕ) we get the other one. The fact that the rhs of (3.41) is skew wrt $(0\leftrightarrow 3,z_1\leftrightarrow z_2)$ finishes the proof. Roughly the same argument works for changes $u_j\leftrightarrow v_j$ for some j's, so we also get $B_{z_1}\circ B'_{z_2}=B'_{z_2}\circ B_{z_1},\ B'_{z_1}\circ B'_{z_2}=B'_{z_2}\circ B'_{z_1}$.

An application of L'Hospital algebraically solves the problem of the iteration of the B transformation for $z_2=z_1$: let $v=v(u_0,v_0,z,b)$ be the solution of (5.3), $v_1=v_1(u_0,v_0,z_1,b_1)$ a particular solution, $v_3=v_3(u_0,v_0,z_1,z,b_1,b),\ z\neq z_1$ be given by (3.33)-(3.35) with $z_2,\ v_2$ replaced by $z,\ v$ and $b=b(z)$ a differentiable function with $b(z_1)=b_1,\ \frac{db}{dz}(z_1)=:b_0$. Letting now $z\to z_1$ in (3.33)-(3.35) and applying L'Hospital we get a definite value of $v_3=v_3(u_0,v_0,z_1,b_1,b_0)$, so the iteration of the B transformation with the same z_1 (done by means of algebraic computations and derivatives) introduces a new constant as expected. Note that in this case x^3 has the same rolling as $x^0:\ (x^3,dx^3)=(R_3,t_3)(x^3_0,dx^3_0),\ (R_3(u_3,v_3),t_3(u_3,v_3))=(R_0(u_0(u_3,v_3),v_0(u_3,v_3)),t_0(u_0(u_3,v_3),v_0(u_3,v_3)))$, so the rolling of a quadric on an applicable surface gives a 3-dimensional family of deformations of that quadric having the same rolling, a fact a-priori not evident. For the $B_z\circ B'_z$ transformation there is no need of L'Hospital; in fact these are the G transformations discovered by Guichard independently of Bianchi; they behave better at the analytic level and by use of permanent conjugate systems (especially for real deformations of real ellipsoids they give directly real deformations of the same real ellipsoid). Because of the BPT Möbius configurations are put in motion and become

moving Möbius configurations once the n leaves $\{x^{2^{k-1}}\}_{k=1,...,n}$ are B transforms of the seed x^0. The static n-soliton x^{2^n-1} of $\mathcal{M}_n$ becomes an actual (moving) soliton if the seed is the vacuum soliton; again one can apply the L'Hospital trick and even higher derivatives when some of $z_1, ...z_n$ are not distinct; again the rolling can be considered for a higher dimensional family of deformations of the considered quadric.

5.4. **Quadrics conjugate in deformation and the Hazzidakis transformation.**
Bianchi in ([4],§ 423) calls two non-homothetic non-flat surfaces conjugate in deformation if a pointwise correspondence is established between them such that the asymptotic coordinates and all the virtual asymptotic coordinates correspond. Because of this the coefficients of (4.7) on the conjugate in deformation $x,\ \tilde{x}$ must be the same, or if we choose (u^1, u^2) asymptotic coordinates on $x,\ \tilde{x}$:

$$\partial_{u^j} \log \frac{K(\tilde{x})}{K(x)} = 2(\Gamma^j_{jj} - \tilde{\Gamma}^j_{jj}) = 4(\Gamma^{j+1}_{12} - \tilde{\Gamma}^{j+1}_{12}),\ \Gamma^{j+1}_{jj} = \tilde{\Gamma}^{j+1}_{jj},\ j = 1, 2. \tag{5.15}$$

The geodesic equations $\frac{d^2u^l}{ds^2} + \Gamma^l_{jk}\frac{du^j}{ds}\frac{du^k}{ds} = 0,\ l = 1, 2$ can be put in implicit form ($u^2 = \phi(u^1)$) as $(\frac{du^1}{ds})^2(\phi'' - (\phi')^3\Gamma^1_{22} + (\phi')^2(\Gamma^1_{22} - 2\Gamma^1_{12}) - \phi'(\Gamma^1_{11} - 2\Gamma^2_{12}) + \Gamma^2_{11}) = 0$ and $\frac{du^1}{ds} = 0 \Rightarrow \Gamma^1_{22} = 0$ (Eisenhart ([19],§ 85)). Thus the geodesics on $x,\ \tilde{x}$ correspond. Conversely if on two non-flat non-homothetic surfaces $x,\ \tilde{x}$ the asymptotic coordinates and geodesics correspond, then the second fundamental forms are proportional with factor $\sqrt{\frac{K(\tilde{x})\tilde{g}}{K(x)g}}$; from the Codazzi-Mainardi equations we see that the equations of virtual asymptotic coordinates (4.7) on $x,\ \tilde{x}$ are the same. Thus conjugation in deformation is equivalent to correspondence of asymptotic coordinates and geodesics. Bianchi proved in ([5],Vol **5**,(87),(90)) for quadrics of revolution and Servant in [28] essentially for the remaining ones that only deformations of quadrics can be conjugate in deformation; these appear according to Bianchi II for $n = 3$. We shall reproduce below this proof.

Dini solved the problem of finding all the surfaces with geodesic correspondence (see Darboux ([13],§ 600-§ 603)). First one can choose on the two surfaces corresponding orthogonal systems. Take a particular tangent facet T of x; directions conjugate to isotropic directions in T and to the directions corresponding to isotropic directions in the corresponding tangent facet $\tilde{T}$ of $\tilde{x}$ are in general unique and distinct; if they are not unique, then the linear elements of $x,\ \tilde{x}$ are proportional; if they are not distinct (Lie pointed out this gap in Dini's proof), then $x,\ \tilde{x}$ are flat and do not present interest to us. Because (5.15) is coordinate invariant, it is valid for the common orthogonal system $(u, v):\ |dx|^2 = Edu^2 + Gdv^2,\ \ |d\tilde{x}|^2 = \tilde{E}du^2 + \tilde{G}dv^2$. From $\Gamma^j_{jj} - \tilde{\Gamma}^j_{jj} = 2(\Gamma^{j+1}_{12} - \tilde{\Gamma}^{j+1}_{12}),\ j = 1, 2$ we get $\tilde{E} = \frac{E}{VU^2},\ \tilde{G} = \frac{G}{UV^2},\ U = U(u),\ V = V(v)$. From $\Gamma^{j+1}_{jj} = \tilde{\Gamma}^{j+1}_{jj},\ j = 1, 2$ we get $E = U_1^2(U - V),\ G = V_1^2(U - V),\ U_1 = U_1(u),\ V_1 = V_1(v)$ (we exclude $U = V =$const which leads to homothetic $x,\ \tilde{x}$), so $|dx|^2 = (U - V)(U_1^2du^2 + V_1^2dv^2),\ |d\tilde{x}|^2 = (\frac{1}{V} - \frac{1}{U})(\frac{U_1^2}{U}du^2 + \frac{V_1^2}{V}dv^2)$. Thus both $x,\ \tilde{x}$ have Lioville linear element (the coordinate curves are orthogonal geodesic confocal ellipses and hyperbolas). This type of linear element generalizes the linear element of quadrics in the elliptic coordinates. Under the transformation $(U, V, U_1^2, V_1^2) \to (m(U + h), m(V + h), \frac{U_1^2}{m}, \frac{V_1^2}{m}),\ m,\ h$ constants $|dx|^2$ is preserved while $|d\tilde{x}|^2$ becomes up to the constant factor $\frac{1}{m^3}$: $(\frac{1}{V+h} - \frac{1}{U+h})(\frac{U_1^2}{U+h}du^2 + \frac{V_1^2}{V+h}dv^2)$. We can assume $|dx|^2 = (u - v)(U^2du^2 + V^2dv^2),\ |d\tilde{x}|^2 = (\frac{1}{v+h} - \frac{1}{u+h})(\frac{U^2}{u+h}du^2 + \frac{V^2}{v+h}dv^2)$. Since $\Gamma^j_{jj} + \Gamma^{j+1}_{12} = \frac{1}{2}\partial_{u^j}\log(g),\ j = 1, 2$, (5.15) are equivalent to correspondence of geodesics and (the coordinate invariant) $cK(x)^{\frac{3}{4}}da = K(\tilde{x})^{\frac{3}{4}}d\tilde{a}$ for some constant c (which we can assume to be 1); here da and $d\tilde{a}$ are the area forms. The last relation becomes: $\partial_u(\frac{V}{U}\frac{1}{u-v}) - \partial_v(\frac{U}{V}\frac{1}{u-v}) = \frac{1}{\sqrt{(u+h)(v+h)}}(\partial_u(\frac{V}{U}\frac{1}{u-v}\sqrt{\frac{v+h}{u+h}}) - \partial_v(\frac{U}{V}\frac{1}{u-v}\sqrt{\frac{u+h}{v+h}}))$, or $\frac{2}{u-v}(\frac{1}{U^2}(\frac{1}{u+h} - 1) + \frac{1}{V^2}(\frac{1}{v+h} - 1)) + \partial_v(\frac{1}{V^2}(\frac{1}{v+h} - 1)) - \partial_u(\frac{1}{U^2}(\frac{1}{u+h} - 1)) = 0$. Thus $\frac{1}{U^2}(\frac{1}{u+h} - 1) = au^2 + bu + c,\ \frac{1}{V^2}(\frac{1}{v+h} - 1) = -(av^2 + bv + c)$; after a normalization we have linear elements of quadrics in elliptic coordinates: $|dx|^2 = (u - v)(\frac{u}{(u-a)(u-b)(u-c)}du^2 - \frac{v}{(v-a)(v-b)(v-c)}dv^2),\ |d\tilde{x}|^2 = (\tilde{u} - \tilde{v})(\frac{\tilde{u}}{(\tilde{u}-\tilde{a})(\tilde{u}-\tilde{b})(\tilde{u}-\tilde{c})}d\tilde{u}^2 -$

$\frac{\tilde v}{(\tilde v-\tilde a)(\tilde v-\tilde b)(\tilde v-\tilde c)}d\tilde v^2)$, $(\tilde u,\tilde v):=\frac{1}{(b-a)(c-a)}(\frac{u}{u-a},\frac{v}{v-a})$, $(\tilde a,\tilde b,\tilde c):=\frac{1}{(b-a)(c-a)}(1,\frac{b}{b-a},\frac{c}{c-a})$. For a comprehensive discussion see Eisenhart ([20],§ 145).

Because of the virtual asymptotic coordinates correspondence, for any deformation of x we know infinitesimally a deformation of the H transform $\tilde x=H(x)$, so we need to integrate a Ricatti equation and a quadrature to recover it.

However the H transformation will turn out to commute with the B transformation ($B_{\tilde z}\circ H=H\circ B_z$) and $(B_{\tilde z}\circ H)(x^0)=\tilde x^1=(H\circ B_z)(x^0)$ can be algebraically recovered from $x^0,\ x^1=B_z(x^0),\ \tilde x^0=H(x^0)$.

In this vein the H transformation has the same involutory character wrt the B transformation as the Christoffel transformation wrt the D transformation of isothermic surfaces.

Assume first that $\tilde u_j,\tilde v_j$ depend respectively only on $u_j,v_j,\ j=0,1$. Since $\frac{\tilde\omega_0}{\tilde{\mathcal{B}}_0}=\frac{\tilde{\mathcal{N}}_0}{\mathcal{N}_0}((\omega^{u_0}_{0u_0}\tilde x^0_{0\tilde u_0}-(\omega^{v_0}_{0u_0}\frac{\tilde v_{0v_0}}{\tilde u_{0u_0}})\tilde x^0_{0\tilde v_0})d\tilde u_0+((\omega^{u_0}_{0v_0}\frac{\tilde u_{0u_0}}{\tilde v_{0v_0}})\tilde x^0_{0\tilde u_0}-\omega^{u_0}_{0u_0}\tilde x^0_{0\tilde v_0})d\tilde v_0)$, $\tilde v_1$ given by the homography of Bianchi II satisfies $(\tilde m^1_0)^T\tilde\omega_0+2\tilde z d\tilde v_1=0$ if the algebraic relations

$$\frac{\tilde\Delta^- d\tilde u_0}{\Delta^- du_0}=\frac{\tilde\Delta^+ d\tilde v_0}{\Delta^+ dv_0}=\frac{\tilde z d\tilde v_1/\tilde{\mathcal{N}}_0}{z dv_1/\mathcal{N}_0} \tag{5.16}$$

are valid. But since (3.26) and $\widetilde{(3.26)}$ are valid for independent $u_0,\ v_0,\ v_1$, (5.16) follows immediately. If $\tilde u_j,\tilde v_j$ do not depend respectively only on $u_j,v_j,\ j=0,1$, then the argument is similar.

Moreover there is an algebraic correspondence between infinitesimal deformations of surfaces conjugate in deformation and thus between W congruences having surfaces conjugate in deformation as one of their focal surfaces (see Bianchi ([4],§ 427)). If $s_0,\ s_3:=s_0+as_{0u}+bs_{0v},\ a=a(u,v),\ b=b(u,v)$ are the focal surfaces of a W congruence and $\tilde s_0$ is conjugate in deformation to s_0, then $\tilde s_0,\ \tilde s_3:=\tilde s_0+a\tilde s_{0u}+b\tilde s_{0v}$ are the focal surfaces of a W congruence. To prove this we can choose (u,v) common conjugate system for the focal surfaces of the W congruence, so $b=0$. Let $s_2:=rN_0$ be the infinitesimal deformation of s_3 from Darboux's 12 surfaces; using (5.15) $\tilde s_2:=r\sqrt[4]{\frac{K(s_0)}{K(\tilde s_0)}}\tilde N_0$ will be the infinitesimal deformation of $\tilde s_3$ from Darboux's 12 surfaces.

5.5. **(Singular) Bäcklund transformations of (singular) quadrics.**
If we let z approach the singular values $0,\ \infty$ or inverses a_j of nonzero eigenvalues of A (under restrictions as follows: we consider only ruled x^0 for $z=0$, only QC for $z=\infty$ and only a_j's such that the eigenspace of a_j^{-1} is not isotropic; these restriction will reveal themselves later), then we get singular B transformations. If we allow one of the a_j's tend to 0 (again under restrictions as above), then we get B transformations of singular quadrics. We shall call B_0 *infinitesimal B transformation*, B_{a_j} *finite singular B transformation* and B_∞ *infinite singular B transformation*.

For $z\to 0$ we have $\frac{\Delta^-}{z}\to\infty(v_1-v_0)^2$, $\frac{\Delta'^+}{z}\to\infty(u_1-u_0)^2$ and $\frac{\Delta^+}{z}$, $\frac{\Delta'^-}{z}$ have definite limits. If $v_1=v_0$, then $\omega^{v_0}_{0u_0}=\omega^{u_0}_{0u_0}=0$ and x^1 is a particular position of x^0_0 as it rolls on the ruled x^0 (with ruling x_{0u_0} corresponding to $x^0_{0u_0}$). If $v_1\neq v_0$, then $\omega^{u_0}_{0u_0}=\omega^{u_0}_{0v_0}=0,\ dv_1=\lim_{z\to0}\frac{\Delta^+}{2z}\omega^{v_0}_{0u_0}du_0$ and $x^0,\ x^1$ are ruled with rulings corresponding to $x^0_{0v_0},\ x^1_{0u_1}$. Note in particular

$$(v_1\to v_0)\circ\lim_{z\to0}\frac{\Delta^+}{z}=-\frac{1}{\mathcal{N}_0}. \tag{5.17}$$

In this case x^1 is obtained from x^0 by the application of Chieffi's result and finding solutions of (5.5) is equivalent to finding asymptotes of x^0. All these results have a simple geometric interpretation: as we roll a quadric on an applicable ruled surface, its ruling families describe two congruences decomposable in ruled surfaces applicable to that quadric. The first strata are the particular positions of the quadric as it rolls on its deformation; the other ones are obtained by assembling the tangents (to geodesics corresponding to the other ruling family on the quadric) along a curvilinear asymptote of the initial ruled surface. We can consider $z_1\to0,\ z_2\neq0$ or $z_1,z_2\to0$ in the BPT for x^0 ruled.

If we let z tend to a finite singular value, then all equations and quantities of interest, infinitesimal (fundamental forms) or finite (surfaces, rollings) have definite limits and enjoy the same properties; the only difference is that the conics (in the tangent spaces of x^0 and which are cut by four B transforms with constant cross ratio) become lines. Note that although the Ivory affinity degenerates, its action from x_0 to x_{a_j} is well defined (double cover), by means of which we can find a double valued inverse (or an inverse from the double cover of x_{a_j} as a set, considered as the tangent bundle of the conic $\mathcal{S}(x_{a_j})$) and complete the computations required to produce the RMPIA. For quadrics of revolution ($a_1 = a_2$ in QC (3.3) or in QWC (3.6)) B_{a_1} becomes the complementary transformation; therefore x^1 will roll on the axis of revolution as x^0 rolls on x_0^0. The BPT remains valid when either (or both) of z_1, z_2 take finite singular values. If the eigenspace of a_j^{-1} is isotropic, then one cannot define $\sqrt{R_{a_j}}$ and thus the Ivory affinity fails. One can try to replace the Ivory affinity with another degenerate affine transformation which should preserve as much as possible from the properties of the Ivory affinity, but most of these are lost and the program cannot be concluded (for example for QC one can preserve the symmetry of the TC by choosing a symmetric linear transformation linear combination of $J_3, J_3^2, e_3 e_3^T$). However limiting processes like the ones used in the discussion of totally real quadrics may provide the correct tools needed to extend the finite singular B transformation to isotropic eigenspaces.

For example for QC (3.3) we only consider the singular B_{a_1} transformation (the other B_{a_j}'s are similar). The singular quadric $x_{a_1}(u,v) := \sqrt{R_{a_1}}(\sqrt{A})^{-1}X(u,v)$ covers $\mathbb{C}^2$ twice: for any $\sqrt{R_{a_1}}(\sqrt{A})^{-1}[0\ a\ b]^T \in \mathbb{C}^2$ take v a solution of $[i\ \ -b\ a]Y(v) = 0$ and $u := -v\frac{1+b}{1-b}$. Through any point $x_{a_1}(u_0, v_0) \in \mathbb{C}^2$ we have two tangents $x_{a_1}(u, v_0),\ x_{a_1}(u_0, v),\ u, v \in \mathbb{C}$ at the conic $\frac{(x^2)^2}{a_2-a_1} + \frac{(x^3)^2}{a_3-a_1} = 1$ which touch the conic for $u = 2v_0^{-1},\ v = 2u_0^{-1}$. For $z \to a_1$ we have $\sqrt{R_z} \to \sqrt{R_{a_1}},\ \frac{\det\sqrt{R_z}(\sqrt{R_z})^{-1}\pm\sqrt{R_z}}{z\det\sqrt{A}} \to \mathrm{diag}[\pm\sqrt{\frac{a_2}{a_1}(a_3-a_1)}\ \ \pm\sqrt{\frac{a_3}{a_1}(a_2-a_1)}\ \ \pm\sqrt{\frac{1}{a_1}(a_2-a_1)(a_3-a_1)}]$ (here the $\pm$ signs depend in a certain fashion on $a_1,\ a_2,\ a_3$). If $\tilde{a}_2 := a_2^{-1},\ \tilde{a}_j := \tilde{a}_2 + (a_j - a_2)^{-1},\ j = 1, 3$, then $\frac{\tilde{a}_2}{\tilde{a}_1}(\tilde{a}_3 - \tilde{a}_1) = \frac{-1}{(a_3-a_2)(a_2-a_1)}\frac{1}{a_1}(a_2-a_1)(a_3-a_1),\ \frac{\tilde{a}_3}{\tilde{a}_1}(\tilde{a}_2 - \tilde{a}_1) = \frac{-1}{(a_3-a_2)(a_2-a_1)}\frac{a_3}{a_1}(a_2 - a_1),\ \frac{1}{\tilde{a}_1}(\tilde{a}_2 - \tilde{a}_1)(\tilde{a}_3 - \tilde{a}_1) = \frac{-1}{(a_3-a_2)(a_2-a_1)}\frac{a_2}{a_1}(a_3 - a_1)$, so $B_{\tilde{a}_1}$ is a conjugation of B_{a_1} with an H transformation (Bianchi). A natural question is what would be the corresponding B transformation conjugate of B_{a_2} with a Hazzidakis transformation; Calapso provided the natural answer: since $\tilde{a}_1$ corresponds to a_1 and ∞ corresponds to a_2, the B_∞ is the correct answer. If $\tilde{a}_1 := a_1^{-1},\ \tilde{a}_j := \tilde{a}_1 + (a_j - a_1)^{-1},\ j = 2, 3$, then $\frac{\tilde{a}_2}{\tilde{a}_1}(\tilde{a}_3 - \tilde{a}_1) = \frac{1}{(a_3-a_1)(a_2-a_1)}a_2,\ \frac{\tilde{a}_3}{\tilde{a}_1}(\tilde{a}_2 - \tilde{a}_1) = \frac{1}{(a_3-a_1)(a_2-a_1)}a_3,\ \frac{1}{\tilde{a}_1}(\tilde{a}_2 - \tilde{a}_1)(\tilde{a}_3 - \tilde{a}_1) = \frac{1}{(a_3-a_1)(a_2-a_1)}a_1$.

B_∞ transformations of QC (3.3) were introduced and studied by Calapso in [11]; he reduced the B_∞ transformation via a conjugation by an H transformation to the singular transformation B_{a_j}, thus naturally completing a formula of Bianchi [4] involving only finite singular transformations B_{a_j}'s.

Calapso studied B transformations of QC (3.3) from an intrinsic point of view; although Bianchi's simple geometric approach is replaced with complicated analytic computations, it is Calapso's approach which provides the correct setting in order to study B_∞ transformations, since as $z \to \infty$ finite quantities (leafs, rollings of leafs) do not have limit. For example B_z transforms of the seed x^0 tend to ∞, (R_0^1, t_0^1) does not have a rigid motion as limit and the ACPIA and the W congruence disappears.

However Calapso realized that for QC (3.3) the TC (at the analytic level), all infinitesimal quantities and all differential equations have a definite valid limit (after proper scaling) when $z \to \infty$. The direction fields $\hat{m}_0^1 := \lim_{z\to\infty} z^{-1}m_0^1 = iY(v_1),\ \hat{m}_0'^1 := \lim_{z\to\infty} z^{-1}m_0'^1 = iY(u_1)$ are well defined, finite and isotropic; they should be well defined finite isotropic direction fields since for $z = a_j$, the facets with normal fields $m_0^1, m_0'^1$ are tangent to the singular set (conic) of the singular quadric x_{a_j}, so $m_0^1,\ m_0'^1$ must be tangent to $C(\infty)$ when $z = \infty$. A simple argument in favor of the existence of the B_∞ transformation for all QC is that these are conjugate in deformation to certain (I)Q(W)C, so B_∞ transformations of QC are conjugation by an H transformation of finite singular

B transformations of certain (I)Q(W)C. Conversely B_∞ transformations for (I)QWC do not exist, since they should be conjugation by an H transformation of finite singular B_{a_j} transformations with a_j^{-1} having isotropic eigenspace. Taking the limit $z \to \infty$ in the scaled equations (5.3)-(5.8) we get the intrinsic characterization of the B_∞ transformation. It is natural to assume that the inversion of the B_∞ transformation is realized if in the inversion of the B_z transformation one makes $z \to \infty$ and one performs proper scalings, by means of which we can find the second fundamental form of x^1 (so we shall infinitesimally know $x^1 = B_\infty(x^0)$).

As $z \to \infty$ the TC becomes $X_0^T\sqrt{A}X_1 = 0$; (5.3) becomes the completely integrable Ricatti equation $Y(v_1)^T\omega_0 = 2idv_1$ and we have the similar equation $Y(u_1)^T\omega_0' = 2idu_1$. Since (3.24), (3.26), (3.27) (and its versions under symmetries) have a definite valid limit when $z \to \infty$, the analytic version of the inversion of the B_∞ transformation holds.

For finite z since x^0, x^1 are the focal surfaces of a W congruence, there exists $s_1 := r_1R_0\frac{m_0^1}{z}$ infinitesimal deformation of x^0; as $z \to \infty$ the limit of r_1 satisfies $d(\log r_1) - \frac{i}{2}Y'(v_1)^T\omega_0 = 0$ and $ds_1 = 0,\ s_1 := r_1R_0Y(v_1)$. Conversely if we have an isotropic vector s_1 and define r_1, v_1 by $r_1Y(v_1) := R_0^{-1}s_1$, then $0 = \frac{-i}{2}Y'(v_1)^Tr_1^{-1}R_0^{-1}ds_1 = Y(v_1)^T\omega_0 - 2idv_1,\ 0 = r_1^{-1}R_0^{-1}ds_1 = (d(\log r_1) - \frac{i}{2}Y'(v_1)^T\omega_1)Y(v_1)$ and we get a B_∞ transformation of x^0. Therefore all B_∞ transforms of x^0 are determined by isotropic translations (this property was communicated to Calapso by Bianchi). For the BPT if $z_1 \to \infty,\ z_2 \in \mathbb{C}$, then (5.13), (5.14), (3.32), (3.33) remain valid. Therefore if $x^1 = B_\infty(x^0),\ x^2 = B_{z_2}(x^0)$, then $x^3 = B_{z_2}(x^1) = B_\infty(x^2)$ can be algebraically recovered from $x^j,\ j = 0, 1, 2$. If $z_1,\ z_2 \to \infty$, then a-priori (3.33) has no definite limit. We can consider (5.13) in this case as the formula for the Permutability Theorem; it becomes $X(v_0, v_3)^T\sqrt{A}X(v_1, v_2) = 0$ and similar formulae for (u_1, u_2) replacing (v_1, v_2), but we only know x^3 infinitesimally from $x^j,\ j = 0, 1, 2$.

For an isotropic vector $c := mf_1 + n\bar{f}_1 + pe_3,\ 2mn + p^2 = 0,\ n \neq 0$ we have $c = \frac{n}{2}Y(\frac{p}{n})$. For $n = 0$ we can consider $c = mf_1 = -mt^2Y(t^{-1})$ for $t \to 0$. We have $c = rR_0Y(v)$, where $v := \frac{c^TR_0e_3}{c^TR_0f_1},\ r := \frac{c^TR_0f_1}{2}$. If $x^1,\ x^2$ are B_∞ transforms of x^0, then $c_1 := r_1R_0Y(v_1),\ c_2 := r_2R_0Y(v_2)$ are constant isotropic vectors, so $R_0X(v_1, v_2) = \frac{\pm c_1\times c_2}{|c_1\times c_2|}$ is a constant unit vector. Using this fact we can reduce the determination of the transforms $x^1 = B_\infty(x^0)$ and of their iterates to quadratures. If we fix $x^0,\ x^1$ and let $x^2,\ x^3$ vary in $X(v_0, v_3)^T\sqrt{A}X(v_1, v_2) = 0$, or equivalently $X(v_0, v_3) = \frac{-iY(v_0)\times\sqrt{A}X(v_1,v_2)}{Y(v_0)^T\sqrt{A}X(v_1,v_2)}$, then a homography will be established between $v_2,\ v_3$. R_1 can be algebraically recovered from $\frac{\pm c_0\times c_3}{|c_0\times c_3|} = R_1X(v_0, v_3) = R_1\frac{-iY(v_0)\times\sqrt{A}R_0^{-1}(c_1\times c_2)}{Y(v_0)^T\sqrt{A}R_0^{-1}(c_1\times c_2)}$, where $c_0 := r_0R_1Y(v_0),\ c_3 := r_3R_1Y(v_3)$ are constant isotropic vectors; then we can recover x^1 from $dx^1 = R_1dx_0^1$ with a quadrature. Note that while we do not know the constant unit vector $\frac{\pm c_0\times c_3}{|c_0\times c_3|}$, we can still find R_1 up to a constant rotation. As a consequence the B_∞ transforms can be found with a quadrature; thus the finite singular B_{a_j} transforms can be found with a quadrature once a certain H transform is known. Note again that the previous argument is the same for $(R_1, c_1) \leftrightarrow (R_2, c_2)$, so R_1 is the rotation of the rolling of all B_∞ transforms of x^0.

Note that while for QWC the B_∞ does not exist, we come close to having such a transformation: $\hat{u}_1 := \lim_{z\to\infty} z^{-\frac{1}{2}}u_1,\ \hat{v}_1 := \lim_{z\to\infty} z^{-\frac{1}{2}}v_1,\ \hat{m}_0^1 := \lim_{z\to\infty} z^{-\frac{3}{2}}m_0^1 = \hat{v}_1^2Y(-i\hat{v}_1^{-1}),\ \hat{m}_0'^1 := \lim_{z\to\infty} z^{-\frac{3}{2}}m_0'^1 = -\frac{1}{2}Y(-2i\hat{u}_1)$; $\hat{u}_1, \hat{v}_1$ satisfy the completely integrable Ricatti equations $Y(-i\hat{v}_1^{-1})^T\omega_0 = 2id(-i\hat{v}_1^{-1}),\ Y(-2i\hat{u}_1)^T\omega_0' = 2id(-2i\hat{u}_1)$ (scaled versions of (5.3)) and the (non-symmetric) scaled version of the TC $iZ_0^T\sqrt{A}\hat{Z}_1 - e_3^T\hat{Z}_1 - \frac{1}{2} = 0$. Again equations (3.24), (3.26), (3.27) (and its versions under symmetries), (5.3)-(5.8) have a definite valid limit when $z \to \infty$ and after proper scaling, by means of which one can prove the same equations with $(u_0, v_0, \hat{\Delta}^+, \hat{\Delta}'^-, \mathcal{N}_0) \leftrightarrow (\hat{u}_1, \hat{v}_1, \hat{\Delta}'^-, \hat{\Delta}^+, \widehat{\mathcal{N}_1})$, but since the scaled $\widehat{\mathcal{N}_1}$ is different than $\mathcal{N}_{\hat{1}}$ as a function of the scaled $\hat{u}_1, \hat{v}_1$, the version of (5.6) thus obtained loses its geometric significance. The generation by isotropic translation property remains, so we can find $\hat{u}_1, \hat{v}_1$ only by algebraic computations.

Similarly we have the case for $a_j \to 0$. For example for QC (3.3) and a_1 we replace X_0 with $[0\ \frac{i(2+u_0v_0)}{\sqrt{2}(u_0-v_0)}\ \frac{u_0+v_0}{u_0-v_0}]^T$ and x_0^0 becomes the tangent surface of a conic $\mathcal{X}_0^0$; thus x^0 becomes the tangent surface of a curve with the same curvature as that of $\mathcal{X}_0^0$ and the rolling of $\mathcal{X}_0^0$ on x^0 becomes the rolling of two developables with the osculating spaces of the lines of regression coinciding. In this case ω_0 depends only on $u_0(v_0)$ and (5.3) becomes a Ricatti ODE upon whose integration we find developables whose lines of regression have the same curvature as that of $\mathcal{X}_0^0$. Therefore we can state this geometrically: as the tangent surface x_0^0 of $\mathcal{X}_0^0$ rolls with correspondence of lines of regressions on an applicable developable the ruling families on a confocal quadric will describe two line congruences. The confocal conic (which is the intersection of the confocal quadric with the plane of the singular quadric) rigidly moved in this rolling will describe a focal surface for both these congruences. The developables of these congruences have lines of regression on this focal surface and will be applicable to x_0^0 (with correspondence of lines of regression).

5.6. **(Totally real) deformations of totally real quadrics.**
We are interested in deformations x^0 of surfaces $x_0^0 \subseteq x_0$ with real linear element $|dx_0^0|^2$.

Let $d\bar{x}_0^0 = Rdx_0^0,\ R \subset \mathbf{O}_3(\mathbb{C}),\ \bar{R}^T = R$. Since $R\hat{N}_0^0$ is a multiple of $\bar{\hat{N}}_0^0$ and keeping account of $K(x_0^0) \subset \mathbb{R}$, we get $R\hat{N}_0^0 = e^{ic}\bar{\hat{N}}_0^0,\ c \in \mathbb{R}$ constant. Applying to this last relation $R^{-1}d$ we get $\omega \times \hat{N}_0^0 = (e^{ic}R^{-1}\bar{A}R - A)dx_0^0$, so $(e^{ic}R^{-1}\bar{A}R - A)\hat{N}_0^0 = \lambda\hat{N}_0^0,\ \lambda \in \mathbb{C}$; by conjugation we obtain $\bar{\lambda} = -e^{-ic}\lambda$. Assume $e^{ic}R^{-1}\bar{A}R - A \neq 0$; if $v \in \mathbb{C}^3 \setminus \{0\}$ is one of the remaining eigenvectors of $e^{ic}R^{-1}\bar{A}R - A$ and having eigenvalue $\mu \in \mathbb{C}$, then $v^T\hat{N}_0^0 = 0$ and $R^{-1}\bar{v}$ is also eigenvector with eigenvalue $-e^{ic}\bar{\mu}$.

Note that $\bar{\hat{N}}_0^0 \times \bar{A}\bar{\hat{N}}_0^0 = e^{-2ic}R\hat{N}_0^0 \times \bar{A}R\hat{N}_0^0 = e^{-ic}\det(R)R^{-1}(\hat{N}_0^0 \times A\hat{N}_0^0)$ combined with (I)&(II) of § 3.2 give $e^{2ic} = \frac{\epsilon\bar{\mathcal{A}}}{\mathcal{A}}$ and $\mathcal{A}^2 \in \mathbb{R},\ e^{4ic} = 1$.

Since $\sqrt{\bar{a}} = \sigma_a\sqrt{a},\ \sigma_a = 1$ for a not negative real and -1 otherwise, we have $\overline{\sqrt{M}} = \sigma_M\sqrt{M}$ where M is SJ with non-isotropic kernel and the signature σ_M is diagonal with ± 1 entries (the sign is same for any SJ block). All computations are preserved by conjugation; for example for QC $(\bar{R}_0,\ \bar{t}_0)(\bar{x}_0^0, d\bar{x}_0^0) = (\bar{x}^0, d\bar{x}^0),\ \bar{x}_0^0 = \sigma_A(\sqrt{\bar{A}^{-1}})^{-1}\bar{X}_0 = (\sqrt{\bar{A}^{-1}})^{-1}X_0(\sigma_{X_0}(\bar{u}_0, \bar{v}_0)),\ \bar{x}_z^1 = ... = \sqrt{\bar{R}_{\bar{z}}}(\sqrt{\bar{A}^{-1}})^{-1}X_1(\sigma_{X_1}(\bar{u}_1, \bar{v}_1))$, where $\sigma_{X_0},\ \sigma_{X_1}$ are compositions of some of the involutions $\sigma_1^2, \sigma_2, \sigma_3$ of § 3.1; thus deforming the quadric x_0 is equivalent to deforming the quadric $\bar{x}_0$. Since the condition $(R_0m_0^1)^T dx^1 = 0$ holds for $x^1 = B_z(x^0)$, the condition $(\bar{R}_0\bar{m}_0^1)^T d\bar{x}^1 = 0$ holds for $\bar{x}^1 = \overline{B_z(x^0)} = B_{\bar{z}}(\bar{x}^0)$ (or $= B'_{\bar{z}}(\bar{x}^0)$), so the $B_{\bar{z}},\ B'_{\bar{z}}$ transforms of $\bar{x}^0$ are conjugates of the $B_z,\ B'_z$ transforms of x^0.

If x_0^0 is totally real (the seed x^0 may be non-totally real), then the B transformation behaves well wrt to these surfaces (due mainly to its algebraic character). For z and initial value of the Ricatti equation (5.3) chosen to satisfy certain totally real conditions the corresponding surface $x_0^1 \subseteq x_0$ of the leaf x^1 is totally real (not necessarily of the same signature as that of x_0^0). However most B_z transforms of seeds x^0 with totally real linear element do not obey these totally real requirements, in which case the leafs x^1 and their corresponding surfaces $x_0^1 \subseteq x_0$ although being real 2-dimensional may have complex linear element. In this case the leaves are not interesting from a totally real point of view, but they are still useful to generate via conjugation and an application of the BPT surfaces with x_0^3 or both x_0^3, x^3 being totally real respectively of the same signature as that of x^0, x_0^0 and x^3 in the same surrounding space as that of x^0.

If $\bar{x}^0 = \epsilon x^0,\ \epsilon := \mathrm{diag}[\epsilon_1\ \epsilon_2\ \epsilon_3]$ and $x^1 = B_z(x^0)$, then $\bar{x}^1$ is a $B_{\bar{z}}$ $(B'_{\bar{z}})$ transform of $\bar{x}^0$. We would like to have $x^2 := \epsilon\bar{x}^1$ a $B_{\bar{z}}$ $(B'_{\bar{z}})$ transform of x^0. Once this is accomplished, if $x^2 \neq x^1$, then one can find x^3 from the BPT applied to $(x^0, (x^1, z), (x^2, \bar{z}))$. Since the first three vertices of the quadrilateral $(x^0, (x^1, z), (x^2, \bar{z})), x^3)$ are preserved under $((x^1, z) \leftrightarrow (x^2, \bar{z})) \circ \epsilon \circ \bar{\cdot}$, we conclude that also the fourth one has to be preserved: $\bar{x}^3 = \epsilon x^3$. If x_0^0 is given by $\begin{bmatrix} x_0^0 \\ 1 \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} x_0^0 \\ 1 \end{bmatrix} = 0$, then, since with $R := \bar{R}_0^T\epsilon R_0,\ t := \epsilon t_0 - \bar{R}_0^T\bar{t}_0$ we have $\begin{bmatrix} \bar{x}_0^0 \\ 1 \end{bmatrix} = \begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_0^0 \\ 1 \end{bmatrix}$, we have

$\begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix}^T \begin{bmatrix} \bar{A} & \bar{B} \\ \bar{B}^T & \bar{C} \end{bmatrix} \begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$ (a quadric is uniquely determined by the requirement that 3 facets in general position are among its tangent planes among the tangent planes of x_0^0 one can certainly find such facets; thus the above equality is true, a-priori, modulo a constant e^{ic}, but it turns out that $c \in \mathbb{R}$ and one can absorb it in A, B, C). This ensures that $\begin{bmatrix} x_{\bar{z}}^2 \\ 1 \end{bmatrix} := \begin{bmatrix} R_0 & t_0 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} x^2 \\ 1 \end{bmatrix} =$ $= \begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} \bar{x}_z^1 \\ 1 \end{bmatrix}$ satisfies $\begin{bmatrix} x_{\bar{z}}^2 \\ 1 \end{bmatrix}^T (\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} - \bar{z} \begin{bmatrix} I_3 & 0 \\ 0^T & 0 \end{bmatrix})^{-1} \begin{bmatrix} x_{\bar{z}}^2 \\ 1 \end{bmatrix} = 0$. We need now only prove that if w_0^1 is a ruling of x_0 at x_0^1 (that is $(w_0^1)^T A w_0^1 = (w_0^1)^T \hat{N}_0^1 = 0$), then $w_0^2 := (\sqrt{R_{\bar{z}}})^{-1} R^T \overline{\sqrt{R_z}} \bar{w}_0^1$ is a ruling of x_0 at x_0^2. This is certainly true, since $RAR^T = \bar{A}$ implies $((\sqrt{R_{\bar{z}}})^{-1} R^T \overline{\sqrt{R_z}})^T A (\sqrt{R_{\bar{z}}})^{-1} R^T \overline{\sqrt{R_z}} = \bar{A}$; also $((\sqrt{R_{\bar{z}}})^{-1} R^T \overline{\sqrt{R_z}})^T \hat{N}_0^2 = \bar{\hat{N}}_0^1$ boils down to the above, $\bar{A}t + \bar{B} = RB$ and $AC(\bar{z}) + (I_3 - \sqrt{R_{\bar{z}}})B = \bar{A}\overline{C(z)} + (I_3 - \overline{\sqrt{R_z}})\bar{B} = 0$.

If $\bar{x}_0^0 = \epsilon x_0^0,\ x^1 = B_z(x^0)$, then as above $x_{\bar{z}}^2 := \epsilon \bar{x}_z^1 \subseteq x_{\bar{z}}$ satisfies the TC requirement $(N_0^0)^T (x_{\bar{z}}^2 - x_0^0) = 0$. If $x_z^1 \neq x_{\bar{z}}^2$, then one can complete $(x_0^0, (x_z^1, z), (x_{\bar{z}}^2, \bar{z}))$ to an SITC quadrilateral $(x_0^0, (x_z^1, z), (x_{\bar{z}}^2, \bar{z}), x_0^3)$. As above we conclude that $\bar{x}_0^3 = \epsilon x_0^3$ we only need to show that $x^2 := (R_0, t_0) x_{\bar{z}}^2$ is a $B_{\bar{z}}$ ($B'_{\bar{z}}$) transform of x^0.

If $|dx_0^0|^2$ is real, then $d\bar{x}_0^0 = R dx_0^0,\ R \in \mathbf{O}_3(\mathbb{C}),\ \bar{R}^T = R$ and $d\bar{x}^0 = \bar{R}_0 R R_0^T dx^0$

Sometimes one gets deformations for free: if $|dx^0|^2$ is real, the conjugation becomes a rigid motion for x^0, but $|dx^1|^2$ may not be real, in which case we get a new deformation $\bar{x}^1$. If x^0 is real, then the composition of two distinct $B_z,\ B_{\bar{z}}$ transformations (that is x^1 not real; we always assume that $B_{\bar{z}} = \overline{B_z}$ is the conjugate of the leaf in discussion) gives, by the BPT, a surface x^3; applying the conjugation to the quadrilateral $(x^0,\ x^1,\ \bar{x}^1,\ x^3)$ we get $\bar{x}^3 = x^3$, so x^3 is real. We shall prove later that if x_0^0 is totally real, then x^3 is applicable to x^0.

This result can be extended to the case when there exists a rigid motion (R, t) such that $x^0 = (R, t)\bar{x}^0$; then, if $x^1 \neq (R, t)\bar{x}^1$, the surface x^3 given by the BPT satisfies $x^3 = (R, t)\bar{x}^3$. In particular, if x^0 is totally real of signature $\mathrm{diag}[\epsilon_1\ \epsilon_2\ \epsilon_3],\ \epsilon_j = \pm 1$, we can take $(R, t) := (\mathrm{diag}[\epsilon_1\ \epsilon_2\ \epsilon_3], 0)$; if x^1 is not totally real of the same signature as x^0, then x^3 is totally real with the same signature as x^0; if furthermore x_0^0 is totally real then x^3 is applicable to x^0. By iteration we obtain Möbius configurations with $z_1,\ z_2,\ ..., z_n$ and their conjugates; the composition $x^{2^{2n}-1}$ of all the transformations $B_{z_j},\ B_{\bar{z}_j},\ j = 1, ..., n$ will have the same character as x^0; moreover $x^{2^{2n}-1}$ will be applicable to x^0 if x_0^0 is totally real.

If we are in the previous conditions ($x^0 = (R, t)\bar{x}^0,\ x^1 \neq (R, t)\bar{x}^1,\ x^3 = (R, t)\bar{x}^3$) and there is a rigid motion (R', t') such that $x_0^0 = (R', t')\bar{x}_0^0$, then $(R_0, t_0)(R', t') = (R, t)(\bar{R}_0, \bar{t}_0)$. We have $x_0^3 = (R_3, t_3)^{-1}(R, t)(\bar{R}_3, \bar{t}_3)^{-1}\bar{x}_0^3$; if $(R_3, t_3)^{-1}(R, t)(\bar{R}_3, \bar{t}_3)^{-1}$ is a rigid motion, then x_0^3 has a character similar to x_0^0. This is the case for x_0^0 totally real: we shall see later that we have $(\bar{u}_0, \bar{v}_0) = \sigma(u_0, v_0)$, where σ is the composition of some of $\sigma_1,\ \sigma_2,\ \sigma_3$ and uniquely describes the totally real quadric x_0^0. Note that the SITC provides homographies between v_0, v_1, v_2, v_3 and u_0, v_1, v_2, u_3; using these two homographies and their conjugates we get $(\bar{u}_3, \bar{v}_3) = \sigma(u_3, v_3)$, so x_0^3 coincides with x_0^0. For example for the real ellipsoid $\sigma_A = I_3$ and we have $u_0\bar{v}_0 = u_3\bar{v}_3 = -2$.

The composition of complex conjugate B transformations (with certain rationality conditions) of the axis of the tractrix (vacuum soliton) gives a surface called a breather (see Rogers-Schief [26]).

With $\sigma_1^2,\ \sigma_2,\ \sigma_3,\ \sigma_4$ the involutions of § 2.1 and $\epsilon =: (-1)^{s(\epsilon)}$ the totally real unit spheres $X(u, v)$ have signatures as follows: $\mathrm{diag}[\epsilon\ \epsilon\ 1]$, $\mathrm{diag}[\epsilon\ -\epsilon\ 1]$, $\mathrm{diag}[\epsilon\ -\epsilon\ -1]$, $\mathrm{diag}[1\ 1\ -1]$ respectively for $\sigma_4 = \sigma_1^{2(s(\epsilon)+1)}\sigma_2\sigma_3,\ \sigma_1^{2s(\epsilon)},\ \sigma_1^{2(s(\epsilon)+1)}\sigma_2,\ \sigma_3$. The totally real equilateral paraboloids $Z(u, v)$ have signatures $\mathrm{diag}[\epsilon\ \epsilon\ 1]$, $\mathrm{diag}[\epsilon\ -\epsilon\ 1]$ for $\sigma_4 = \sigma_1^{2s(\epsilon)}\sigma_2,\ \sigma_1^{2s(\epsilon)}$. Thus all totally real QC (3.3) and QWC (3.6) with $a_j \in \mathbb{R}$ are obtained respectively from totally real spheres and equilateral paraboloids by changing the signature ot the j^{th} coordinate if $a_j < 0$. We need now only describe the totally real QC (3.3), QWC (3.6) with $\sqrt{a_1^{-1}} = \sqrt{\bar{a}_2^{-1}} = b_1 + ib_2 \notin \mathbb{R} \cup i\mathbb{R}$, the remaining

quadrics (for which only $a_j \in \mathbb{R}$ is possible) and some (I)QWC for which the canonical form does not admit totally real realizations with the coordinates being real or purely imaginary.

5.7. **Discrete deformations of quadrics.**
The study of Möbius configurations by Bianchi ([5],Vol **5**,(117)) and discrete CGC and isothermic surfaces by Bobenko-Pinkall [6],[7] motivates the study of DDQ.

A $\mathbb{Z} \times \mathbb{Z}$ lattice of B transforms $x^{(j,k)}$ of a deformation $x^{(0,0)}$ of a surface $x_0^{(0,0)} \subset x_0$, x_0 being a quadric ($x^{(j+1,k+1)} = B_{z_j}(x^{(j,k+1)}) = B_{z'_k}(x^{(j+1,k)})$, $z_j, z'_k \in \mathbb{C}$, $j,k \in \mathbb{Z}$) forms a special 2-dimensional family of DDQ; a DDQ is formed by points corresponding under the B transformations. All surfaces $x^{(j,k)}$ can be algebraically obtained once we know the surfaces on a connected *initial value tree* (IVT) which contains once each transformation B_{z_j}, $B_{z'_k}$, $j,k \in \mathbb{Z}$ (for example the cross $x^{(j,0)}$, $x^{(0,k)}$, $j,k \in \mathbb{Z}$ or the staircase $x^{(j,j)}$, $x^{(j+1,j)}$, $j \in \mathbb{Z}$); more generally from a connected tree containing once each transformation B_{z_j}, $B_{z'_k}$, $j,k \in \mathbb{Z}$ one can recover the corresponding double infinite Möbius moving configuration and the DDQ as a subset of this (for example the connected tree $x^{(0,0)}$, $B_{z_j}(x^{(0,0)})$, $B_{z'_k}(x_{(0,0)})$, $j,k \in \mathbb{Z}$). The four segments joining a point of a DDQ with its neighbors reside in the tangent plane of the subjacent deformation, so a DDQ naturally admits a discrete Gauß map.

From (5.9) we get:

$$(R_{(j+1,k)}, t_{(j+1,k)})^{-1}(R_{(j,k)}, t_{(j,k)}) = (R_{(j,k)}^{(j+1,k)}, t_{(j,k)}^{(j+1,k)}),$$
$$(R_{(j,k+1)}, t_{(j,k+1)})^{-1}(R_{(j,k)}, t_{(j,k)}) = (R_{(j,k)}^{(j,k+1)}, t_{(j,k)}^{(j,k+1)}) \tag{5.18}$$

and the permutability theorem allows us to recover all the surfaces $x_{(j,k)}$ from an IVT using only algebraic computations:

$$(R_{(j+1,k+1)}, t_{(j+1,k+1)})^{-1}(R_{(j,k)}, t_{(j,k)}) = (R_{(j+1,k)}^{(j,k+1)}, t_{(j+1,k)}^{(j,k+1)}). \tag{5.19}$$

One can remove the scaffolding of subjacent surfaces; thus the notion of DDQ can be parted with the notion of deformations of quadrics.

A DDQ is a $\mathbb{Z} \times \mathbb{Z}$ lattice of facets $(x^{(j,k)}, T^{(j,k)})$ such that

(i) $T^{(j,k)}$ contains the points of its 4 neighboring facets.

(ii) There is a $\mathbb{Z} \times \mathbb{Z}$ lattice of facets $(x_0^{(j,k)}, T_{x_0^{(j,k)}} x_0)$ in the quadric x_0 and a $\mathbb{Z} \times \mathbb{Z}$ lattice of rigid motions $(R_{(j,k)}, t_{(j,k)})$ (discrete rolling) satisfying (5.18) and $(R_{(j,k)}, t_{(j,k)})(x_0^{(j,k)}, T_{x_0^{(j,k)}} x_0) = (x^{(j,k)}, T^{(j,k)})$.

In requiring that (5.18) is satisfied we assume that $(R_{(j,k)}, t_{(j,k)})^{-1}(x^{(j,k+1)}, T^{(j,k+1)})$ is the facet centered at $x_{z'_k}^{(j,k+1)}$ and spanned by $V_{(j,k)}^{(j,k+1)}$ and one of the rulings of $x_{z'_k}$ at $x_{z'_k}^{(j,k+1)}$, etc.

The discrete rolling $(x^{(j,k)}, T^{(j,k)}) = (R_{(j,k)}, t_{(j,k)})(x_0^{(j,k)}, T_{x_0^{(j,k)}} x_0)$ is a natural discrete version of the rolling $(x, dx) = (R, t)(x_0, dx_0)$.

As the rotation of the rolling must have constant determinant, all discrete rotations $R_{(j,k)}$ must have same determinant (we can take it to be 1), so the rulings involved in a homography of a Bianchi quadrilateral must be the same; we can take them to be v's. However DDQ can have discrete rollings with discrete rotations of different determinants, as long as the co-cycle requirements are satisfied.

The discrete version of the compatibility condition $d \wedge \omega_0 + \frac{1}{2}\omega_0 \times \wedge\omega_0 = 0$ is
$R_{(j,k)}^{-1} R_{(j+1,k)} R_{(j+1,k)}^{-1} R_{(j+1,k+1)} = R_{(j,k)}^{-1} R_{(j,k+1)} R_{(j,k+1)}^{-1} R_{(j+1,k+1)}$, or the co-cycle relation
$(R_{(j+1,k)}^{(j,k)}, t_{(j+1,k)}^{(j,k)})(R_{(j+1,k+1)}^{(j+1,k)}, t_{(j+1,k+1)}^{(j+1,k)}) = (R_{(j,k+1)}^{(j,k)}, t_{(j,k+1)}^{(j,k)})(R_{(j+1,k+1)}^{(j,k+1)}, t_{(j+1,k+1)}^{(j,k+1)})$.

A DDQ is recovered only by algebraic computations from the knowledge of the facets $(x^{(j,k)}, T^{(j,k)})$ along an IVT and a particular point $x_0^{(j_0,k_0)}$ on x_0 corresponding to a point $x^{(j_0,k_0)}$ of a facet on that tree. That is so because we know a rigid motion $(R_{(j_0,k_0)}, t_{(j_0,k_0)})$ (modulo a rotation of the facet into itself), then the confocal quadrics upon which the points of the neighboring facets are taken by this rigid motion, so by use of the RMPIA's we know the facets on x_0 corresponding

to the neighboring facets (of the initial facet), etc. Note that the facets along an IVT cannot be arbitrarily prescribed: if we roll a facet to its corresponding point on x_0 by a rigid motion, then the neighboring rolled facets must contain one of the rulings of the confocal quadric they are situated on (there are three choices of such quadrics so 6 choices of such facets), so if we prescribe the lengths of the segments of an IVT and the angles between neighboring segments of an IVT, then the DDQ is uniquely determined, modulo countable many choices.

While all discrete versions of the fundamental forms, Gauß curvature, etc are good as long as one replaces each derivative with an edge, the correct discrete version is the one where computations are exact and not approximate to order two. The fundamental forms can be found at the level of a single Bianchi quadrilateral; we consider the one from the discussion of the SITC with the edges and the discrete projective Gauß map highlighted:

$$\begin{array}{ccc} R_2^0\hat{N}_0^2 & \xrightarrow{\mathcal{A}(R_2^1\hat{N}_0^3)\times R_2^0\hat{N}_0^2} & R_2^1\hat{N}_0^3 \\ {\scriptstyle \mathcal{A}(R_2^0\hat{N}_0^2)\times\hat{N}_0^0}\uparrow & & \uparrow{\scriptstyle \mathcal{A}(R_2^1\hat{N}_0^3)\times R_1^0\hat{N}_0^1} \\ \hat{N}_0^0 & \xrightarrow{\mathcal{A}(R_1^0\hat{N}_0^1)\times\hat{N}_0^0} & R_1^0\hat{N}_0^1 \end{array}$$

From (3.22) we have $x_u^0 \rightsquigarrow V_0^1 = \mathcal{A}R_1^0\hat{N}_0^1 \times \hat{N}_0^0$, $x_v^0 \rightsquigarrow V_0^2 = \mathcal{A}R_2^0\hat{N}_0^2 \times \hat{N}_0^0$. It is convenient to work with the projective Gauß map $\hat{N}$ instead of N: $\hat{N}_u^0 \rightsquigarrow R_1^0\hat{N}_0^1 - \hat{N}_0^0$, $\hat{N}_v^0 \rightsquigarrow R_2^0\hat{N}_0^2 - \hat{N}_0^0$. From the existence of the Bianchi quadrilateral we get $\mathcal{A}(R_2^0\hat{N}_0^2) \times \hat{N}_0^0 + \mathcal{A}(R_2^1\hat{N}_0^3) \times R_2^0\hat{N}_0^2 = \mathcal{A}(R_1^0\hat{N}_0^1) \times \hat{N}_0^0 + \mathcal{A}(R_2^1\hat{N}_0^3) \times R_1^0\hat{N}_0^1$; equivalently

$$(R_2^1\hat{N}_0^3 - \hat{N}_0^0) \times (R_1^0\hat{N}_0^1 - R_2^0\hat{N}_0^2) = 0. \tag{5.20}$$

A similar relation for CGC -1 appears in Bobenko-Pinkall [7]. Thus the existence of the Bianchi quadrilateral is the discrete version of $x_{uv}^0 = x_{vu}^0$ (equivalently of the compatibility condition $\omega_0 \times \wedge dx_0^0 = 0$).

Since $(x_u^0)^T\hat{N}_u^0 \rightsquigarrow \mathcal{A}(R_1^0\hat{N}_0^1 \times \hat{N}_0^0)^T(R_1^0\hat{N}_0^1 - \hat{N}_0^0) = 0$, it is clear that the piecewise linear curves j, k =const on the DDQ are discrete versions of asymptotes. The discrete second fundamental form is $(x_{uv}^0)^T\frac{\hat{N}^0}{|\hat{N}^0|} \rightsquigarrow \mathcal{A}\frac{(\hat{N}_0^0)^T}{|\hat{N}_0^0|}((R_2^1\hat{N}_0^3) \times R_2^0\hat{N}_0^2) = \mathcal{A}\frac{(\hat{N}_0^0)^T}{|\hat{N}_0^0|}((R_2^1\hat{N}_0^3) \times R_1^0\hat{N}_0^1)$. Using (5.20) the condition that the discrete Gauß curvature coincides with the Gauß curvature $K(x_0^0) = \frac{-1}{\mathcal{A}^2|\hat{N}_0^0|^4}$ becomes $(\hat{N}_0^0)^T((R_1^0\hat{N}_0^1) \times R_2^0\hat{N}_0^2) = 0$, which is straightforward.

Bobenko-Pinkall [7] discuss many other issues, including discrete B transformations and the corresponding discrete permutability theorem. Most of their results should remain valid for DDQ as long as the correct metric-projective equivalents are found.

Now the existence of the $\mathcal{M}_3$ configuration has to do with commuting of third order derivatives, so the algebraic relations needed for the existence of this configuration may be the correct discretization of the GCM equations.

5.8. **Delaunay's, Guichard's and Weingarten's results.**

5.8.1. *Delaunay's result and its converse.* All diagonal parabolas are homothetic to $p_0 = [2\sinh(u)\ \ \sinh^2(u)]^T$ with focus $f = [0\ \ 1]^T$ we have the rolling of p_0 on the line $p = [0\ \ \cosh(u)\sinh(u) + u)]^T$: $(p, dp) = (R, t)(p_0, dp_0)$, where R is the rotation $\begin{bmatrix} \tanh(u) & -\frac{1}{\cosh(u)} \\ \frac{1}{\cosh(u)} & \tanh(u) \end{bmatrix}$.

The curve described by the focus through the rolling of p_0 on p is the catenary $p + R(f - p_0) = [\cosh(u), u]^T$, so we get the statement for minimal surfaces of revolution.

All diagonal conics with center are homothetic to $e_0 = [\cos(u)\ b\sin(u)]^T$ with excess E, $E^2 := b^2 - 1$ and foci $f = [0\ E]^T$ (there are two choices for the sign of E). We have the rolling of e_0 on the line $e = [0\ \int\sqrt{1 + E^2\cos^2(u)}du]^T$: $(e, de) = (R, t)(e_0, de_0)$, where R is the rotation $\frac{1}{\sqrt{1+E^2\cos^2(u)}}\begin{bmatrix} b\cos(u) & \sin(u) \\ -\sin(u) & b\cos(u) \end{bmatrix}$. The curves described by the foci through the rolling of e_0 on

e are $c = e + R(f - e_0) = e + \frac{E\sin(u)-b}{\sqrt{1+E^2\cos^2(u)}}[1 \ -E\cos(u)]^T$ and if we denote $[\cos(r) \ \sin(r)]^T := \frac{1}{\sqrt{1+E^2\cos^2(u)}}[1 \ -E\cos(u)]^T$, then we have $dr = -\cot(r)d\log(\cos(r)) = \frac{E\sin(u)du}{1+E^2\cos^2(u)}$ and $dc = \frac{1}{\sqrt{1+E^2\cos^2(u)}}[E\cos(u) \ 1]^T du + (E\sin(u)-b)d(\frac{1}{\sqrt{1+E^2\cos^2(u)}}[1 \ -E\cos(u)]^T) = (du + (E\sin(u)-b)dr)[-\sin(r) \ \cos(r)]^T = \frac{b(b-E\sin(u))du}{1+E^2\cos^2(u)}[-\sin(r) \ \cos(r)]^T$.

We have $dc \times d^2c = \frac{E\sin(u)[b(b-E\sin(u))]^2du^3}{[1+E^2\cos^2(u)]^3}[0 \ 0 \ 1]^T$ and the curvature k_c of c is $\frac{|dc\times d^2c|}{|dc|^3} = \frac{E\sin(u)}{b(b-E\sin(u))}$. The mean curvature of the surface obtained by the rotation of c around the y axis is the sum of k_c and $\cos(r)$ of the inverse of the x coordinate (the projection of the curvature of the parallel of the surface on the normal to the surface), that is $\frac{E\sin(u)}{b(b-E\sin(u))} + \frac{1}{\sqrt{1+E^2\cos^2(u)}}\frac{\sqrt{1+E^2\cos^2(u)}}{E\sin(u)-b} = -\frac{1}{b}$, so we obtain CMC surface of revolution.

For the general conic with center ($e_0 = [a\cos(u) \ b\sin(u)]^T$) we get the mean curvature $-\frac{1}{b}$ since the mean curvature scales inversely. Also because CMC surfaces of revolution are described in arc-length parametrization of the meridian by the 2^{nd} order ODE $H = \sqrt{1-x'^2}(\frac{1}{x} - \frac{x''}{1-x'^2})$ all CMC surface of revolution are obtained in this way (pick initial conditions at a point where the x coordinate is critical, so we have only two parameters).

For $u, a, b \in \mathbb{R}, \ b > a > 0$ ($i(u-\frac{\pi}{2}), ia, b \in \mathbb{R}$) we have real ellipse (hyperbola), real foci and real CMC (the real ellipse $[a\cos(u) \ b\sin(u)]^T, \ a, b, u \in \mathbb{R}, \ b > a > 0$ (hyperbola $[a\sinh(u) \ b\cosh(u)]^T$, $a, b, u \in \mathbb{R}$) has also imaginary foci $[\pm i\sqrt{b^2-a^2} \ 0)]^T$ ($[\pm i\sqrt{a^2+b^2} \ 0]^T$)).

Conversely we have the rolling of the curve c_0 on the line $c = [0 \ \int |dc_0|]^T : \ (c, dc) = (R, t)(c_0, dc_0)$, $R^T = [N_0 \ \frac{dc_0}{|dc_0|}], \ t = c - Rc_0$. Then t is the curve described by a point (which can be considered the origin) rigidly moved with c_0 as it rolls on c. We have: $R^{-1}dt = \frac{N_0^T R^{-1}dRdc_0}{|dc_0|^2}(N_0 \times dc_0) \times c_0, \ |dt|^2 = k_{c_0}^2|c_0|^2|dc_0|^2, \ N_t = \frac{1}{|c_0|}Rc_0, \ k_t = \frac{N_t^T d^2t}{|dt|^2} = \frac{1}{|c_0||dt|^2}(|dt|^2 - N_0^T c_0 N_0^T R^{-1}dRdc_0) = \frac{1}{|c_0|} + \frac{N_0^T c_0}{k_{c_0}|c_0|^3}$. Since the surface of revolution described by the rotation of t around the y axis has CMC, t coincides with the curve described by a (the) focus of a conic as it rolls on the y axis is uniquely determined modulo rigid motions by $|dt|, \ k_t$. But these quantities uniquely determine $|c_0|, \ |dc_0|$, which in turn uniquely determine modulo rigid motions c_0 (since $(N_0^T c_0)^2 = |c_0|^2 - \frac{1}{4|dc_0|^2}(d|c_0|^2)^2$ and $k_{c_0}N_0^t c_0|dc_0| = c_0^T d\frac{dc_0}{|dc_0|} = d\frac{d|c_0|^2}{2|dc_0|} - |dc_0|$). Thus c_0 is uniquely determined and must be a conic.

5.8.2. *Guichard's result and its converse.* (see Bianchi ([4], ch XX))

For this subsubsection we use the invariant notation established at the beginning of 4.2.

Bianchi proposed and solved the inversion of the Guichard result: find all surfaces x_0 and fixed points (which can be considered the origin) such that all the surfaces t described by this point through the rolling of x_0 on an applicable surface x have the same CMC $H(t)$ (including the statement in space forms and CGC $K(t)$).

From (4.9) we thus need:

$$x_{0(j+1)(k+1)} = (-1)^{j+k+1}n^{-1}B^{jk}g, \ 2K(x_0)n + x_{0jk}B^{jk} = 0; \tag{5.21}$$

using Gauß theorem for x_0 we get from the first three relations $K(x_0)n^2 = (N_0^T x_0)^2|x_0|^2$, so the last relation is a consequence of the first three.

We are interested in the Gauß equation of x_0 for the square of the distance from a point and distance from a plane (Darboux equations).

Let $\rho := |x_0|^2, \ x_0 = (N_0N_0^T + dx_0 \llcorner \nabla x_0^T)x_0$, so $4(N_0^T x_0)^2 = 4\rho - |\nabla\rho|^2$.

Also $\nabla^2\rho = 2(x_0^T\nabla^2 x_0 + |dx_0|^2) = 2(x_0^T N_0 N_0^T d^2x_0 + |dx_0|^2)$ and thus the Gauß equation becomes, with the area form $da := \sqrt{\det(|dx_0|^2)} = \sqrt{g}du^1 \wedge du^2$:

$$\det(\nabla^2\rho - 2|dx_0|^2) = (4\rho - |\nabla\rho|^2)K(x_0)da^2. \tag{5.22}$$

For e constant unit vector let $\tau := e^T x_0$; then $e = (N_0 N_0^T + dx_0 \llcorner \nabla x_0^T)e$, so $(e^T N_0)^2 = 1 - |\nabla\tau|^2$, $\nabla^2\tau = e^T\nabla^2 x_0 = e^T N_0 N_0^T d^2 x_0$, so the Gauß equation becomes:

$$\det(\nabla^2\tau) = (1 - |\nabla\tau|^2)K(x_0)da^2. \tag{5.23}$$

Once either version of the Gauß equation is satisfied the Codazzi-Mainardi equations are automatically satisfied.

Also from (5.22) one can easily read Darboux's result about cyclic systems: circles of cyclic systems have centers at $x_0 + \frac{\nabla x_0(\mu)}{2}$ and radii $\mu + \frac{|\nabla\mu|^2}{4}$ where $-\mu$ satisfies the Gauß equation for the square of the distance from a point. With $\mu^2 := \rho$ we have from (5.21) $\frac{\nabla^2(\mu^2) - 2|dx_0|^2}{2\mu\sqrt{1-|\nabla\mu|^2}} = N_0^T d^2 x_0 = -\frac{(da\llcorner\nabla\mu)^2 + (1-|\nabla\mu|^2)|dx_0|^2}{2(H(t)\mu+1)\mu\sqrt{1-|\nabla\mu|^2}}$, so we get

$$\mu\nabla^2\mu + d\mu^2 - |dx_0|^2 = -\frac{(da\llcorner\nabla\mu)^2 + (1-|\nabla\mu|^2)|dx_0|^2}{2(H(t)\mu+1)}. \tag{5.24}$$

Applying to this $2\llcorner\nabla\mu$ and using $d(|\nabla\mu|^2) = 2\nabla^2\mu\llcorner\nabla\mu$ we get $\mu d(|\nabla\mu|^2) + 2|\nabla\mu|^2 d\mu - 2d\mu = -\frac{(1-|\nabla\mu|^2)d\mu}{H(t)\mu+1}$, or $\frac{-d(|\nabla\mu|^2)}{1-|\nabla\mu|^2} = \frac{(2H(t)\mu+1)d\mu}{\mu(H(t)\mu+1)}$, so $d\log(1-|\nabla\mu|^2) = d\log(\frac{1}{\mu(H(t)\mu+1)})$; thus $|\nabla\mu|^2 = 1 - \frac{c}{\mu(H(t)\mu+1)}$ for some constant c. Applying $\llcorner\llcorner\partial x_0^2$ to (5.24) we get $\mu\Delta\mu + |\nabla\mu|^2 - 2 = -\frac{|\nabla\mu|^2 + 2(1-|\nabla\mu|^2)}{2(H(t)\mu+1)}$, $\Delta\mu := \nabla^2\mu\llcorner\llcorner\partial x_0^2$. Applying again $\llcorner\llcorner(d\mu\llcorner\partial a)^2$, $\partial a := (\sqrt{g})^{-1}\partial_{u^1}\wedge\partial_{u^2}$ to (5.24) we get $\mu\nabla^2\mu\llcorner\llcorner(d\mu\llcorner\partial a)^2 - |\nabla\mu|^2 = -\frac{|\nabla\mu|^4 + (1-|\nabla\mu|^2)|\nabla\mu|^2}{2(H(t)\mu+1)}$. Equation (5.22) can be rewritten as: $\mu^2\det(\nabla^2\mu) + (\mu\nabla^2\mu\llcorner\llcorner(d\mu\llcorner\partial a)^2 - \mu\Delta\mu - |\nabla\mu|^2 + 1)da^2 = \mu^2(1-|\nabla\mu|^2)K(x_0)da^2$, or $\det(\nabla^2\mu) = (1-|\nabla\mu|^2)(K(x_0) + \frac{H(t)}{\mu(H(t)\mu+1)})da^2$. If we set $\tau := k\mu$, $k := \frac{1}{\sqrt{1+4cH(t)}}$, then $1 - |\nabla\tau|^2 = k^2(1-|\nabla\mu|^2)(1+2H(t)\mu)^2$, $K(x_0) = \frac{1}{4\mu^2(H(t)\mu+1)^2}$ and τ satisfies (5.23).

Thus x_0 has the property that the ratio of the distances to a point and to a plane is a constant, so x_0 is a quadric of revolution around the focal axis (with center if $k \neq 1$ and without center if $k = 1$).

Bianchi ([4], ch XX) proved that all CMC surfaces occur as translations of rolling of quadrics of revolution (around the focal axis) in a 2-dimensional fashion for a fixed quadric; this is how the 2-dimensionality of the B transformation for quadrics of revolution was first established. In particular all real surfaces applicable to real quadrics of revolution around the focal axis are real analytic in the common conjugate system of coordinates (a conformal parametrization of the second fundamental form), which corresponds to the real analytic lines of curvature parametrization on the real CMC surface.

The reality conditions are not necessary for the validity of the computations, so we get statements about surfaces in $\mathbb{C}^3$ applicable to quadrics of revolution and CMC surfaces in $\mathbb{C}^3$.

5.8.3. *Darboux's geometrization of Weingarten's result.* (see Bianchi ([5],Vol **4**,(202)))

5.9. **Conic systems.**
Motivated by Darboux's and Ribacour's result about cyclic systems, consider the integrable 3-dimensional distribution of facets obtained as follows: intersect the tangent planes of a surface y_0 with a quadric x_z (thus along conics); for each point of the conic take the facet determined by the segment joining the point of the conic with the origin of the tangent plane and one of the rulings passing through that point. As we roll y_0 on any applicable surface y : $(y, dy) = (R, t)(y_0, dy_0)$ while rigidly moving this distribution in the process, does this distribution always remain integrable? The case when x_z is non-isotropic singular quadric (tangent bundle of a conic) was classified by Bianchi ([5],Vol **4**,(173)): y_0 must be a quadric confocal to x_z and we recover the finite singular B transformations of x_0 (Bianchi in fact showed more: a non-isotropic plane must also have the projective structure of a singular quadric, as a-priori the choice of rulings is not well defined in a plane).

Again it is convenient to derive certain identities before imposing the TC: consider $x_z = x_z(u,v),\ |N_0| = 1,\ N_0^T dy_0 = 0,\ V := x_z - y_0$ and (3.13) becomes

$$(V_v V_u^T + V_u V_v^T)N_0 = V_{uv}V^T N_0 + \frac{2}{\mathcal{B}}(A^{\circ -1} - zI_3 + \Gamma)N_0,$$

$$\Gamma := -(V+y_0)y_0^T \text{ for QC, } := (V+y_0)(Le_3)^T + Le_3 y_0^T \text{ for (I)QWC.} \tag{5.25}$$

Again multiplying (5.25) on the left respectively with $N_0^T, (V\times N_0)^T$ in the TC assumption $V^T N_0 = 0$ and with $n := (A^{\circ -1} - y_0 y_0^T)N_0$ for QC, $:= (A^{\circ -1} + y_0(Le_3)^T + Le_3 y_0^T)N_0$ for (I)QWC, $m := \mathcal{B}x_{zu}\times V,\ m' := \mathcal{B}x_{zv}\times V$ we get the analogous of (3.12):

$$N_0^T x_{zu} N_0^T x_{zv}\mathcal{B} = n^T N_0 - z,$$

$$x_{zv}^T(I_3 - 2N_0N_0^T)m = -2V^T(N_0\times n). \tag{5.26}$$

The condition (5.1) remains the same; equation (5.3) becomes

$$m^T\omega m^T N_0 + 2m^T(zN_0 - n)dv = 0,\ \omega := \alpha^{-1}(R^{-1}dR). \tag{5.27}$$

Since applying directly (4.12) is complicated (we have $V = \frac{m\times N_0}{N_0^T\mathcal{B}x_{zu}}$), direct computations again must be applied as in the B transformation case. If we let $z\to\infty$ for x_z QC, then we get $N_0^T X = 0,\ Y(v)^T\omega = 2idv$ again isotropic translations, just as for B_∞. This can also be considered as infinite cyclic systems: we intersect the tangent spaces of a surface with a null cone with center at ∞; the Darboux equation for the distance from a point to a surface becomes the equation for the distance from a point at ∞ to the surface, that is the distance from a plane to the surface. This can be easier seen by considering the surrounding space of constant curvature k and letting $k\to 0$.

Thus applying $d\wedge$ to (5.27), using the equation itself, $dm = m_v dv + dy_0\times\mathcal{B}x_{zu}$, (4.6) and $dm^T\wedge\omega = dv\wedge m_v^T\omega$ we get $(m^TN_0)^2(N_0^T(m\times m_v + 2zm) - 2m^Tn)N_0^T(\frac{1}{2}\omega\times\wedge\omega) + 2m^T\omega\wedge(-(dy_0\times\mathcal{B}x_{zu})^T(N_0\times n)\times m + m^TN_0m^T(zdN_0 - dn) - m^T(zN_0 - n)m^TdN_0) = 0$. Note $((N_0\times n)\times m)\times\mathcal{B}x_{zu} = -(N_0\times n)^T\mathcal{B}x_{zu}m$ and the remaining two terms in the second parenthesis give $m^T(ndN_0^T - N_0dn^T)m$. Similarly to (3.18) we can use (5.26) to get $N_0^T(m\times m_v + 2zm) = \frac{2zV^T(N_0\times n) - n^TN_0V^Tm_v}{N_0^Tx_{zv}} = 2m^Tn$ (use $V^Tm_v = -m^Tx_{zv}$ and further (5.26) again). Thus we get the restriction $2m^T\omega\wedge(dy_0^Tm(\mathcal{B}x_{zu})^T(N_0\times n) + m^T(ndN_0^T - N_0dn^T)m) = 0$; this is a polynomial of degree ≤ 6 in v which must be identically 0 for any v and choice of ω satisfying (4.6) and thus the restrictions on the geometry of y_0 revealed themselves: $m^Tdy_0 = 0$ (in general this is over-determined; the infinitesimal B transformation may be considered an example) or $dy_0^Tm(\mathcal{B}x_{zu})^T(N_0\times n) + m^T(ndN_0^T - N_0dn^T)m = 0$.

6. Particular deformations in $\mathbb{C}^{n+p}$ of quadrics in $\mathbb{C}^{n+1}$

6.1. Overview and particular deformations in $\mathbb{C}^{2+p}$ of quadrics in $\mathbb{C}^3$.

Some results about surfaces do not generalize for sub-manifolds in scalar product spaces. If the co-dimension is lower, the GCMR equation will impose higher restrictions. However the Ricci equations (partially) disappear if we only consider deformations with (partially) flat normal bundle; further the existence of imaginary numbers relieves part of restrictions (for example according to Cartan [12] $\mathbf{H}^n(\mathbb{R})$ cannot be locally immersed in $\mathbb{R}^{n+p},\ p < n-1$, but it can be embedded in $\mathbb{R}^n\times i\mathbb{R}$ as a pseudo-sphere; also Tenenblat and Terng in [32] have developed the B transformation for local immersions of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$; more generally Tenenblat, Terng, Uhlenbeck and collaborators for space forms in space forms). Beginning with a seed sub-manifold the same formalism of B transformation can be applied and we can replace the GCRM equations of the B transform with a Ricatti equation.

Note that (4.5) is true for sub-manifolds in $\mathbb{C}^{n+p}$: if $(x,dx) = (R,t)(x_0,dx_0),\ (R,t)\subset \mathbb{O}_{n+p}(\mathbb{C}^{n+p})\ltimes\mathbb{C}^{n+p},\ a_{jkl} := \partial_{u^j}x_0^TR^{-1}\partial_{u^k}R\partial_{u^l}x_0$, then $a_{jkl} = a_{jlk},\ a_{jkl} = -a_{lkj}$ so $a_{jlk} = -a_{klj} = -a_{kjl} = a_{ljk} = a_{lkj} = -a_{jkl} = -a_{jlk}$. Let N_0 be orthonormal frame of $x_0,\ N := RN_0,\ \mathbb{C}^{n+p}\ni a = a^\top + a^\perp = (I - N_0N_0^T)a + N_0N_0^Ta,\ R^{-1}dRa = (N_0N_0^TR^{-1}dR(I - N_0N_0^T) + R^{-1}dRN_0N_0^T)a = ([N_0, R^{-1}dRN_0] - N_0N_0^TR^{-1}dRN_0N_0^T)a$, where $[a,b] := ba^T - ab^T$. The part $[N_0, R^{-1}dRN_0]$

will encode the difference of the second fundamental forms of x, x_0 : $[N_0, R^{-1}dRN_0]dx_0 = R^{-1}NN^Td^2x - N_0N_0^Td^2x_0$, while the part $N_0N_0^TR^{-1}dRN_0N_0^T$ the difference of the normal connections of x, x_0 : $N_0N_0^TR^{-1}dRN_0 = R^{-1}NN^TdN - N_0N_0^TdN_0$. The rotation of the rolling can be changed so as to take N_0 to any other normal frame N_a of x : $N_a := aN,\ a \subset \mathbb{O}_{n+p}(\mathbb{C}^{n+p}),\ adx = dx,\ R_a := aR$.

We can now complete the only remaining part of the proof when $n = 2$: the complete integrability of the Ricatti equation; in doing so the necessary conditions on the seed x^0 reveal themselves. Let $N^0 := R_0N_0^0$ be the first vector of a normal frame of x^0.

We have $\omega_0 := N_0^0 \times (R_0^{-1}dR_0N_0^0)^{\perp_{\mathbb{C}^3}}$ and we need only prove $0 = d\wedge\omega_0 + \frac{1}{2}\omega_0 \times \wedge\omega_0 = \frac{1}{2}((R_0^{-1}dN^0)^{\perp_{\mathbb{C}^3}} \times \wedge(R_0^{-1}dN^0)^{\perp_{\mathbb{C}^3}} - dN_0^0 \times \wedge dN_0^0) + N_0^0 \times d\wedge(R_0^{-1}dR_0N_0^0)^{\perp_{\mathbb{C}^3}}$. The first term is normal, while the second one is tangent, so both must be 0. We have $(R_0^{-1}dN^0)^{\perp_{\mathbb{C}^3}} = (dx_0^0)^TR_0^{-1}dN^0\llcorner\nabla x_0^0 = -(d^2x^0)^TN^0\llcorner\nabla x_0^0,\ dN_0^0 = -(d^2x_0^0)^TN_0^0\llcorner\nabla x_0^0,\ -d\wedge(R_0^{-1}dR_0N_0^0)^{\perp_{\mathbb{C}^3}} = d\wedge((N_0^0)^TR_0^{-1}dR_0(dx_0^0\llcorner\nabla x_0^0)) = d\wedge(((N^0)^Td^2x^0 - (N_0^0)^Td^2x_0^0)\llcorner\nabla x_0^0)$. But $0 = d\wedge((N_0^0)^Td^2x_0^0\llcorner\nabla x_0^0)$ are the Codazzi-Mainardi equations of x_0^0, so the two terms being 0 is equivalent to the statement that x^0 is a sub-manifold of a ruled 3-dimensional flat $x^0 + wR_0N_0^0,\ w \in \mathbb{C}$ in $\mathbb{C}^{p+2}$. The B transformation thus obtained is more general than the one for $p = 1$ because the flats of the seed and the leaf are in general distinct (there is the question if this B transformation can be extended to B transformation of the corresponding flats, just as the B transformation for quadrics in $\mathbb{C}^3$ can be extended the the B transformation of triply conjugate systems). This result is similar to the B transformation of singular 2-dimensional quadrics, but is not in the spirit of Tenenblat-Terng B transformation, so to get a higher dimensional generalization of deformation of quadrics a different point of view with $p = n - 1$ must be established. We reproduce in the next subsection the Tenenblat-Terng B transformation, in the hope that it will provide us with insight as to what is the correct setting.

6.2. Tenenblat-Terng's Bäcklund transformation of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$.

We need the notion of two $(n-1)$-dimensional subspaces $V_1,\ V_2$ of $(\mathbb{R}^{2n-2},\ <,>)$ being *isoclinic* (having constant inclination σ). Define a symmetric bilinear inner product $<,>'$ on V_2 by $< v_1, v_2 >' := < (v_1)^{\perp_{V_1}}, (v_2)^{\perp_{V_1}} >$. There is a self-adjoint operator A on V_2 such that $< v_1, v_2 >' = < Av_1, v_2 >$; the angles $\sigma_1, ..., \sigma_{n-1}$ between $V_1,\ V_2$ are defined such that $\cos^2(\sigma_j),\ j = 1, ..., n-1$ are the eigenvalues of A. Geometrically these are obtained as follows: $\cos\sigma_1 = < v_1, (v_1)^{\perp_{V_1}} > := \max_{v\in V_2,\ |v|=1} < v, (v)^{\perp_{V_1}} >$, $\cos\sigma_2 = < v_2, (v_2)^{\perp_{V_1}} > := \max_{v\in V_2,\ |v|=1,\ <v,v_1>=0} < v, (v)^{\perp_{V_1}} >$ and so on. This accounts for orthonormal bases $E^j = [E'^j\ E''^j],\ E'^j = [e_1^j ... e_{n-1}^j],\ E''^j = [e_n^j ... e_{2n-2}^j],\ j = 1, 2$ of $\mathbb{R}^{2n-2}$ such that E'^j is basis of $V_j,\ j = 1, 2,\ AE'^2 = E'^2D'^2$ and

$$E^2 = E^1 \begin{bmatrix} D' & -D'' \\ D'' & D' \end{bmatrix}, \tag{6.1}$$

where $D' := \operatorname{diag}[\cos\sigma_1 ... \cos\sigma_{n-1}],\ D'' := \operatorname{diag}[\sin\sigma_1 ... \sin\sigma_{n-1}]$. If $\sigma_1 = ... = \sigma_{n-1} =: \sigma$, then $V_1,\ V_2$ have constant inclination σ.

Consider x an n-dimensional sub-manifold of $\mathbb{R}^N$ and $X = [X'\ X''],\ X' = [e_1 ... e_n],\ X'' = [e_{n+1} ... e_N]$ a moving orthonormal frame along x such that $\{e_j\}_{j=1,...,n}$ are tangent and $\{e_\alpha\}_{\alpha=n+1,...,N}$ normal. Then $\omega' := X'^TdX',\ \omega'' := X'^TdX'' = -(X''^TdX')^T,\ \tilde{\omega} := X''^TdX''$ will respectively encode the Levi-Civita connection on x, the second fundamental form of x and the normal connection of x. We have $\omega'^T = -\omega',\ \tilde{\omega}^T = -\tilde{\omega},\ dx = X'\omega,\ \omega := X'^Tdx$, so $d\wedge\omega = dX'^T\wedge X'\omega = -\omega'\wedge\omega$. Since $0 = X''^Tdx$ we get $0 = d\wedge(X''^Tdx) = dX''^T\wedge X'\omega = \omega''^T\wedge\omega$. Using Cartan's lemma we find $\omega''^T = H(\omega)$, where H is linear symmetric (if $\omega = \{\omega^j\}_{j=1,...,n},\ \omega'' = \{\omega''^j{}_\alpha\}_{j=1,...,n,\ \alpha=n+1,...,N},\ H = \{h^j_{\alpha i}\}_{i,j=1,...,n,\ \alpha=n+1,...,N}$, then $\omega''^i{}_\alpha = \sum_j h^i_{\alpha j}\omega^j,\ h^i_{\alpha j} = h^j_{\alpha i}$). If $\Omega' := d\wedge\omega' + \omega'\wedge\omega',\ \tilde{\Omega} := d\wedge\tilde{\omega} + \tilde{\omega}\wedge\tilde{\omega}$ are the Gauß and normal curvature forms, then imposing the compatibility condition $d\wedge(X^TdX) + (X^TdX)\wedge(X^TdX) = 0,\ X^TdX = \begin{bmatrix} \omega' & \omega'' \\ -\omega''^T & \tilde{\omega} \end{bmatrix}$ we obtain the GCMR equations:

$$\Omega' - \omega'' \wedge \omega''^T = 0,\ d \wedge \omega'' + \omega' \wedge \omega'' + \omega'' \wedge \tilde{\omega} = 0,\ \tilde{\Omega} - \omega''^T \wedge \omega'' = 0.$$

The first fundamental form of x is $|dx|^2 = |\omega|^2 := \sum_j (\omega^j)^2$ and the second one is $X''X''^T d^2x = X''\omega''^T\omega = \sum_{i,\alpha} \omega^i \omega''^i_\alpha e_\alpha = \sum_{i,j,\alpha} h^i_{\alpha j}\omega^i\omega^j e_\alpha$. x has CGC -1 iff $\Omega' = -\omega \wedge \omega^T$; if it has flat normal bundle $\tilde{\Omega} = 0$, then we can choose X' formed by principal directions $h^i_{\alpha j} = \delta^i_j b^i_\alpha$ and X'' giving normal connection $\tilde{\omega} = 0$. We have the next theorem due to Cartan and Moore:

(Cartan-Moore) If x is an n-dimensional CGC -1 in $\mathbb{R}^{2n-1}$, then it has flat normal bundle. Let X'' be a normal frame such that $\tilde{\omega} = 0$. Then x admits a parametrization by its lines of curvature (Tchebyshef coordinates):

$$|dx|^2 = \sum_i a_i^2 (du^i)^2,\ a_i \neq 0,\ \sum_i a_i^2 = 1,\ X''X''^T d^2x = \sum_{i,\alpha} b^i_\alpha a_i^2 (du^i)^2 e_\alpha.$$

Moreover $\sum_i \partial_{u_i}$ is the unique asymptotic vector in the first orthant.

For example if we have $\{e_j\}_{j=1,...,2n-1}$ the standard basis of $\mathbb{R}^{2n-1}$ and 'rotate' the tractrix $0 < t \to \int_0^t \sqrt{1-e^{-2u}}du e_1 + e^{-t}\sum_{i=2}^n \lambda^i e_{2i-2},\ \lambda^i > 0,\ \sum_{i=2}^n (\lambda^i)^2 = 1$, then we obtain $x = \int_0^{u^1} \sqrt{1-e^{-2u}}du e_1 + e^{-u^1}\sum_{i=2}^n \lambda^i(\cos\frac{u^i}{\lambda^i}e_{2i-2} + \sin\frac{u^i}{\lambda^i}e_{2i-1})$ and $X' = [\frac{\partial_{u^1}x}{|\partial_{u^1}x|}...\frac{\partial_{u^n}x}{|\partial_{u^n}x|}]$, $\omega'^j_i = 0,\ i,j > 1,\ \omega'^i_1 = -e^{-u^1}du^i,\ i > 1,\ \omega = [du^1\ e^{-u^1}du^2...e^{-u^1}du^n]^T$. One can easily check that $\Omega' = -\omega \wedge \omega^T$, so x has CGC -1. For $u^1 = \text{const}$ we obtain Clifford tori (deformations of $\mathbb{R}^{n-1}$ in $\mathbb{S}^{2n-3}$), so the correct generalization of the 2-dimensional real pseudo-sphere in $\mathbb{R}^3$ is to replace the homothetic parallels with homothetic Clifford tori (under the same homothety). These flats are parallel horospheres in $\mathbf{H}^n(\mathbb{R})$, so the n-dimensional real pseudo-sphere is applicable to a region bounded by a horosphere of radius half of the Poincaré's unit ball model of $\mathbf{H}^n(\mathbb{R})$; the orthogonal trajectories of these flats are meridians and thus one can establish the applicability correspondence at the level of linear elements without need of further computing the curvature Ω' (the parallel horospheres on the pseudo-sphere $\subset \mathbb{R}^n \times (i\mathbb{R})$ are intersections of this with the pencil of n-dimensional isotropic spaces tangent to $C(\infty)$ at one point). To get the parametrization as in the theorem further apply the change $u^1 = \ln(\cosh(v^1)),\ v^1 > 0$; the normal frame is $\frac{1}{\sqrt{1-e^{-2u^1}(1-(\lambda^i)^2)}}\{e^{-u^1}\lambda^i e_1 + \sqrt{1-e^{-2u^1}}(\cos\frac{u^i}{\lambda^i}e_{2i-2} + \sin\frac{u^i}{\lambda^i}e_{2i-1})\}_{i=2,...,n}$. Note that we never essentially used the fact that the particular deformation of $\mathbb{R}^{n-1}$ in $\mathbb{S}^{2n-3}$ is a Clifford torus, so from any such deformation we get one of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$.

Since $-\{\omega^i \wedge \omega^j\}_{i,j=1,...,n} = -\omega \wedge \omega^T = \Omega' = \omega'' \wedge \omega''^T = \{\sum_\alpha \omega''^i_\alpha \wedge \omega''^j_\alpha\}_{i,j=1,...,n} = \{\sum_\alpha b^i_\alpha b^j_\alpha \omega^i \wedge \omega^j\}_{i,j=1,...,n}$ we get

$$\sum_\alpha b^i_\alpha b^j_\alpha = -1,\ i \neq j. \tag{6.2}$$

Since $\sum_i \partial_{u^i}$ is an asymptotic vector we get $\sum_{i,\alpha} b^i_\alpha a_i^2 e_\alpha = 0$, or

$$\sum_i b^i_\alpha a_i^2 = 0\ \forall\ \alpha = n+1, ..., 2n-1. \tag{6.3}$$

Multiplying this by b^j_α and summing after α we get $\sum_{i,\alpha} b^j_\alpha b^i_\alpha a_i^2 = 0$ using (6.2) we finally get $(\sum_\alpha a_j b^j_\alpha)^2 + a_j^2 = 1$. Thus the n-dimensional sub-manifold $A := \{a^i_j\}_{i,j=1,...,n},\ a^1_i := a_i,\ i = 1, ..., n,\ a^i_j := a_j b^j_{n+i-1},\ i = 2, ..., n,\ j = 1, ..., n$ is orthogonal and has nonzero entries on the first row. The GCMR equations for x become the *generalized sine-Gordon equation* (GSGE) for A. Consider on x the frame X' formed by the principal vectors $v_i := \frac{1}{a^1_i}\partial_{u^i}x$ and X'' for which $\tilde{\omega} = 0$. Then $\omega^i = a^1_i du^i,\ \omega'^i_j = \frac{\partial_{u^j}a^1_i}{a^1_j}du^i - \frac{\partial_{u^i}a^1_j}{a^1_i}du^j,\ \omega''^i_{n+j-1} = a^j_i du^i,\ i = 2, ..., n,\ j = 2, ..., n$; with $\delta := \text{diag}[du^1...du^n]$ we have $\omega = \delta A^T e_1$ (here $e_1 \in \mathbb{R}^n$ is the first vector of the standard basis and in the future it may be in $\mathbb{R}^m$ for different values of m, as suitable) and the GSGE becomes the

completely integrable

$$d\wedge\omega'+\omega'\wedge\omega'=-\delta A^Te_1\wedge(\delta A^Te_1)^T,\ \delta\wedge\omega'+A^TdA\wedge\delta=0. \tag{6.4}$$

Conversely, if the orthogonal n-dimensional sub-manifold A satisfies the GSGE and has nonzero entries on the first row, then we know the first and second fundamental forms of a CGC -1 n-dimensional sub-manifold in the lines of curvature parametrization and we can recover it by the integration of a Ricatti equation and a quadrature. Note however that the GSGE cannot apply to the pseudo-sphere itself since the shape is imaginary degenerate.

Note that in order to get Terng's orthogonal A of the GSGE we only used the shape (extrinsic) part of the Gauß equation; the remaining intrinsic part of the Gauß equation and the Codazzi-Mainardi-Ricci equations followed immediately and did not further impose functional restrictions on A; in fact if one d-wedges the intrinsic Gauß equation (the first equation of (6.4)), then one naturally gets the second equation of (6.4) (namely the Codazzi-Mainardi equations) and conversely. In fact this is the method employed by Berger, Bryant and Griffiths [3] by using only the Gauß equation and its consequences obtained by d-wedging (because of its algebraic character one does mostly Algebraic Geometry); once the obstructions imposed by the Gauß equation and its higher d-wedges are satisfied, everything else falls into place. If one makes a detailed analysis of (6.4) and à la Cartan-Kähler, then one gets the Berger, Bryant and Griffiths [3] dependence of deformations of $\mathbf{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$ on $n(n-1)$ functions of one variable. Note that Cartan's argument according to which the co-dimension $n-1$ cannot be lowered follows from linear algebra and his exteriorly orthogonal symmetric 2-forms: basically from the extrinsic part of the Gauß equation we have $\sum_{\alpha=1}^r H_\alpha[e_i,e_j]H_\alpha+[e_i,e_j]=0$, where $\{e_i\}_{i=1,...,n}$ is the standard basis of $\mathbb{R}^n$, $[e_i,e_j]:=e_je_i^T-e_ie_j^T$ and the symmetric shape operator is $\{H_\alpha\}_{\alpha=1,...,r}=\{h^j_{\alpha i}\}_{i,j=1,...,n,\alpha=1,...,r}$, so $H_1,...,H_r,I_n$ are exteriorly orthogonal symmetric matrices. This exteriorly orthogonal property remains invariant under linear transformations of $\mathbb{R}^n$ (the exteriorly part) and under orthogonal transformations of $\mathbb{R}^{r+1}$ (the orthogonal part). Cartan proves in ([12], ch II, § 20) that if $r\leq n-1$ and one completes $H_1,...,H_r,I_n$ with 0's up to n exteriorly orthogonal symmetric 2-forms, then the resulting forms are linearly independent (so $r=n-1$) and can be found from an orthogonal substitution of 2-forms which are perfect squares: this is exactly the above theorem if the squares are $(\sqrt{|a_i|}du^i)^2$ and Terng's completely integrable GSGE shows that the infinitesimal picture (purely algebraic in character) integrates to the finite picture with no other restrictions.

In the construction of the B transformation and the BPT two types of frames will be useful: principal vectors completed with a normal frame with 0 normal connection and frames in which the normal spaces of x_0 and $x_1:=B_\sigma(x_0)$ clearly appear isoclinic. The change of tangent frames of x_0 will be an orthogonal n-dimensional sub-manifold. An important observation is that this is A_1; by symmetry the change of tangent frames of x_1 will be $JA_0,\ J:=\mathrm{diag}[-1\ 1...1]$.

The orthogonal A_1 (and thus the B transform $x_1=B_\sigma(x_0)$) will be given by the integration of the Ricatti equation:

$$dA_1=A_1\omega_0'+A_1\delta A_0^TDA_1-DA_0\delta, \tag{6.5}$$

where $D:=\mathrm{diag}[\csc\sigma\ \cot\sigma...\cot\sigma]$. Imposing the compatibility condition $d\wedge$ on (6.5) and using the equation itself we thus need: $0=(A_1\omega_0'+A_1\delta A_0^TDA_1-DA_0\delta)\wedge\omega_0'+A_1d\wedge\omega_0'+(A_1\omega_0'+A_1\delta A_0^TDA_1-DA_0\delta)\wedge\delta A_0^TDA_1-A_1\delta\wedge dA_0^TDA_1-A_1\delta A_0^TD\wedge(A_1\omega_0'+A_1\delta A_0^TDA_1-DA_0\delta)-DdA_0\wedge\delta=A_1(\omega_0'\wedge\omega_0'+d\wedge\omega_0'+(DA_0\delta)^T\wedge(DA_0\delta))-DA_0(\delta\wedge\omega_0'+A_0^TdA_0\wedge\delta)+A_1(\omega_0'\wedge\delta+\delta\wedge A_0^TdA_0)A_0^TDA_1$. But because A_0 satisfies GSGE, this becomes $\delta A_0^Te_1\wedge e_1^TA_0\delta=(DA_0\delta)^T\wedge(DA_0\delta)$, which is straightforward (use $\csc^2\sigma-\cot^2\sigma=1$ and $D^2=\cot^2\sigma I_n+e_1e_1^T$). Therefore (6.5) is completely integrable and admits solution for any initial value of A_1. If the initial value is orthogonal, we would like the solution to remain orthogonal, or $0=dA_1^TA_1+A_1^TdA_1=(-\omega_0'A_1^T+A_1^TDA_0\delta A_1^T-\delta A_0^TD)A_1+A_1^T(A_1\omega_0'+A_1\delta A_0^TDA_1-DA_0\delta)$ which is straightforward.

It is useful for the simplicity of the computations to add to the normal frames with $n-1$ columns the zero column vector as the first column and still call the extended 'frames' thus obtained frames; all computations remain valid; for example $\omega_0'':=\delta A_0^T\frac{J+I}{2}$ is augmented by a first column 0 ($\omega''_{0in}:=$

0), which does not change the GCMR equations or any other equation. Putting together $d\wedge\omega_0 = -\omega_0'\wedge\omega_0,\ d\wedge\omega_0''+\omega_0'\wedge\omega_0''=0$, or $-\delta\wedge dA_0^Te_1 = -\omega_0'\wedge\delta A_0^Te_1,\ -\delta\wedge dA_0^T\frac{J+I}{2}+\omega_0'\wedge\delta A_0^T\frac{J+I}{2}=0$ we get the second equation of (6.4). With $i(A) := \begin{bmatrix} A & 0\\ 0 & I\end{bmatrix}$ the change of tangent frames extends to frames as $[X_0'A_1^T\ \ X_0''] = X_0i(A_1^T)$. Since the normal spaces of $x_0,\ x_1$ in the frames $X_0i(A_1^T),\ X_1i(A_0^TJ)$ must have equal inclination σ and $x_1 := x_0+\sin\sigma X_0'A_1^Te_1$ is the formula for the B transform, we must extend (6.1) by inserting a row and a column of zeros in the middle and a block diagonal matrix of dimension 1 to the upper left corner and having entry -1. Thus

$$X_1i(A_0^TJ) = X_0i(A_1^T)R_\sigma, \tag{6.6}$$

where $R_\sigma := \left[\begin{array}{cc|cc} -1 & 0 & 0 & 0\\ 0 & \cos\sigma I_{n-1} & 0 & \sin\sigma I_{n-1}\\ \hline 0 & 0 & 0_{1,1} & 0\\ 0 & -\sin\sigma I_{n-1} & 0 & \cos\sigma I_{n-1}\end{array}\right] = \sin\sigma\begin{bmatrix} DJ & \frac{J+I}{2}\\ -\frac{J+I}{2} & D\frac{J+I}{2}\end{bmatrix}$ (multiplication on the right (left) by $\frac{J+I}{2}$ (respectively $\frac{I-J}{2}$) replaces the first column (row) with 0 or respectively the remaining columns (rows) with 0). Therefore $X_1' = \sin\sigma(X_0'A_1^TD - X_0''\frac{J+I}{2})A_0,\ X_1'' = \sin\sigma(X_0'A_1^T+X_0''D)\frac{J+I}{2}$. Since $\omega_0' = {X_0'}^TdX_0'$, (6.5) can be written as $A_1{X_0'}^Td(X_0'A_1^T) = DA_0\delta A_1^T - A_1\delta A_0^TD$. We have $dx_1 = X_0'\omega_0+\sin\sigma d(X_0'A_1^T)e_1 = (X_0'\delta A_0^T+\sin\sigma d(X_0'A_1^T))e_1$, so ${X_1''}^Tdx_1 = \sin\sigma\frac{J+I}{2}(A_1\delta A_0^T+\sin\sigma A_1{X_0'}^Td(X_0'A_1^T)-\sin\sigma D{\omega_0''}^TA_1^T)e_1 = \sin\sigma\frac{J+I}{2}(A_1\delta A_0^T+\sin\sigma(DA_0\delta A_1^T - A_1\delta A_0^TD)-\sin\sigma\frac{J+I}{2}DA_0\delta A_1^T)e_1 = 0$, since $\sin\sigma De_1 = e_1$. Also ${X_1'}^Tdx_1 =$
$= \sin\sigma A_0^T(\sin\sigma\frac{J+I}{2}{\omega_0''}^TA_1^T + DA_1\delta A_0^T+\sin\sigma DA_1{X_0'}^Td(X_0'A_1^T))e_1 = \sin\sigma A_0^T(\sin\sigma\frac{J+I}{2}A_0\delta A_1^T + DA_1\delta A_0^T+\sin\sigma D(DA_0\delta A_1^T - A_1\delta A_0^TD))e_1 = \sin^2\sigma A_0^T(\frac{J+I}{2}+D^2)A_0\delta A_1^Te_1 = \delta A_1^Te_1$, so $\omega_1 := {X_1'}^Tdx_1 = \delta A_1^Te_1$ and thus x_1 is n-dimensional if A_1 has nonzero entries on the first row. We have ${X_1'}^TdX_1'' = \sin^2\sigma A_0^T(\frac{J+I}{2}{\omega_0''}^TA_1^T + DA_1{X_0'}^Td(X_0'A_1^T)+DA_1\omega_0''D)\frac{J+I}{2} = \sin^2\sigma A_0^T(\frac{J+I}{2}A_0\delta A_1^T + D(DA_0\delta A_1^T - A_1\delta A_0^TD)+DA_1\delta A_0^TD)\frac{J+I}{2} = \delta A_1^T\frac{J+I}{2}$, so $\omega_1'' := {X_1'}^TdX_1'' = \delta A_1^T\frac{J+I}{2}$. Therefore x_1 has CGC -1: $\Omega_1' = \omega_1''\wedge{\omega_1''}^T = -\omega_1\wedge\omega_1^T$ and zero normal connection: ${X_1''}^TdX_1'' = \frac{J+I}{2}\sin^2\sigma(A_1{X_0'}^Td(X_0'A_1^T)+A_1\omega_0''D-D{\omega_0''}^TA_1)\frac{J+I}{2} = \frac{J+I}{2}\sin^2\sigma(DA_0\delta A_1^T-A_1\delta A_0^TD+A_1\delta A_0^TD-DA_0\delta A_1^T)\frac{J+I}{2} = 0$. A_1 satisfies GSGE, all computations admit the symmetry $(0,\sigma)\leftrightarrow(1,-\sigma)$ and lines of curvature (asymptotic directions) correspond on $x_0,\ x_1$.

Note that from the isoclinic spaces property we always assumed that all angles σ are positive; in fact some of them can be negative, so all computations involving diagonal matrices with entries $\sin\sigma$ are valid if one attaches a sign to these entries (which may differ from entry to entry). The signs can be absorbed in the rows of A_1; since these changes do not change the linear element and the shape of x_1, they do not change x_1. For example in equation (6.5) the signs of the entries in the diagonal D are transferred by multiplication on the left to the rows of A_1 and in $x_1 = x_0+\sin\sigma X_0'A_1^Te_1$ one can multiply the rows of A_1 with same signs and keep $\sin\sigma>0$ for all frames in the isoclinic picture.

6.2.1. *Bianchi Permutability Theorem.* Let $x_i = B_{\sigma_i}(x_0),\ i=1,2$. We need to find just by algebraic computations $B_{\sigma_2}(x_1) = x_3 = B_{\sigma_1}(x_2)$. We assume the theorem to be true, derive an algebraic formula and post-priori show that the formula works. From (6.6) we thus have: $X_2i(A_0^T)R_{-\sigma_2}i(JA_2)i(A_1^T)R_{\sigma_1}i(JA_0) = X_0i(A_1^T)R_{\sigma_1}i(JA_0) = X_1 = X_3i(A_1^T)R_{-\sigma_2}i(JA_3) =$
$X_2i(A_3^T)R_{\sigma_1}i(JA_2)i(A_1^T)R_{-\sigma_2}i(JA_3)$, or with $X := A_3A_0^T,\ C := A_2A_1^T$: $i(X)R_{-\sigma_2}i(JC)R_{\sigma_1}i(J) = R_{\sigma_1}i(JC)R_{-\sigma_2}i(JX)$, or $X(D_2CD_1+\frac{J+I}{2}) = (D_1CD_2+\frac{J+I}{2})X,\ (X(D_1-D_2C)-(D_1C-D_2))\frac{J+I}{2} = \frac{J+I}{2}((D_1-CD_2)X-(CD_1-D_2)) = 0$. Since $\sin\sigma_1(X_0'A_1^T+\sin\sigma_2(X_0'A_1^TD_1 - X_0''\frac{J+I}{2})A_0A_3^T)e_1 = x_1-x_0+\sin\sigma_2X_1'A_3^Te_1 = x_3-x_0 = x_2-x_0+\sin\sigma_1X_2'A_3^Te_1 = \sin\sigma_2(X_0'A_2^T+\sin\sigma_1(X_0'A_2^TD_2 - X_0''\frac{J+I}{2})A_0A_3^T)e_1$, we get $X_0'A_1^T(D_2+D_1X^T-C^T(D_1+D_2X^T))e_1 = 0$, or $e_1^T(X(D_1-D_2C)-(D_1C-D_2)) = 0$. If we denote $a := X(D_1-D_2C)-(D_1C-D_2)$, then we

have $e_1^T a = a\frac{J+I}{2} = 0,\ a = a\frac{I-J}{2}$. But $0 = (X(D_2CD_1 + \frac{J+I}{2}) - (D_1CD_2 + \frac{J+I}{2})X)(D_1 - D_2C) = (D_1C - D_2)D_2a + aD_2(D_1C - D_2)$, or $a(\frac{I-J}{2}D_2(D_1C - D_2)\frac{J+I}{2}) = 0$, so $a = 0$ and

$$X(D_1 - D_2C) = D_1C - D_2. \tag{6.7}$$

Therefore we only need to prove that A_3 given by (6.7) satisfies (6.5) for $(A_0,\ \sigma)$ replaced by $(A_1,\ \sigma_2),\ (A_2, \sigma_1)$; by symmetry it is enough to prove only one relation. Since $dA_1 = A_1\omega_0' + A_1\delta A_0^T D_1 A_1 - D_1 A_0\delta,\ dA_2 = A_2\omega_0' + A_2\delta A_0^T D_2 A_2 - D_2 A_0\delta$, we get $d(A_2A_1^T) = (A_2\omega_0' + A_2\delta A_0^T D_2 A_2 - D_2 A_0\delta)A_1^T + A_2(-\omega_0' A_1^T + A_1^T D_1 A_0\delta A_1^T - \delta A_0^T D_1) = A_2(\delta A_0^T(D_2A_2 - D_1A_1) - (A_2^T D_2 - A_1^T D_1)A_0\delta)A_1^T$. Thus if we prove the similar relation $d(A_3A_0^T) = A_3(\delta A_1^T(D_2A_3 + D_1A_0) - (A_3^T D_2 + A_0^T D_1)A_1\delta)A_0^T$, then since $dA_0 = A_0\omega_1' - A_0\delta A_1^T D_1A_0 + D_1A_1\delta$ we obtain what we want: $dA_3 = A_3\omega_1' + A_3\delta A_1^T D_2A_3 - D_2A_1\delta$.

But the formulae for $d(A_2A_1^T),\ d(A_3A_0^T)$ can be written as: $dC = CA_1\delta A_0^T(D_2C - D_1) - (D_2 - CD_1)A_0\delta A_1^T,\ dX = XA_0\delta A_1^T(D_2X + D_1) - (D_2 + XD_1)A_1\delta A_0^T$ and differentiating (6.7) we get $dX(D_1 - D_2C) = (XD_2 + D_1)dC$; thus we need to prove $(XA_0\delta A_1^T(D_2X + D_1) - (D_2 + XD_1)A_1\delta A_0^T)(D_1 - D_2C) = (XD_2 + D_1)(CA_1\delta A_0^T(D_2C - D_1) - (D_2 - CD_1)A_0\delta A_1^T)$. The terms containing $A_1\delta A_0^T$ become: $(-(D_2+XD_1)+(XD_2+D_1)C)A_1\delta A_0^T(D_1-D_2C)$, which is 0 because of (6.7). Using (6.7) we have $(D_2X+D_1)(D_1-D_2C) = D_1^2-D_2^2$, $(XD_2+D_1)(D_2-CD_1) = X(D_2^2-D_1^2)$ and the terms containing $A_0\delta A_1^T$ cancel because $D_2^2 - D_1^2 = (\cot^2\sigma_2 - \cot^2\sigma_1)I$. Finally we need $I = X^TX = ((D_1-D_2C)^{-1})^T(D_1C-D_2)^T(D_1C-D_2)(D_1-D_2C)^{-1}$, or $0 = (D_1C-D_2)^T(D_1C-D_2) - (D_1 - D_2C)^T(D_1 - D_2C) = C^T(D_1^2 - D_2^2)C + D_2^2 - D_1^2 = (\cot^2\sigma_2 - \cot^2\sigma_1)(I - C^TC)$.

Again to prove the existence of the $\mathcal{M}_p$ configuration we need only prove the existence of the $\mathcal{M}_3$ configuration (discrete deformations of $\mathbf{H}^n(\mathbb{R})$ will be obtained by considering $\mathbb{Z}^n$ lattices of B transformations which are subsets of infinite Möbius configurations).

Consider $(D_1D_2)^{-1}((D_1^2 - D_2^2)D_1A_1(D_1A_1 - D_2A_2)^{-1} - D_1^2) = A_3A_0^{-1} = (D_1D_2)^{-1}((D_1^2 - D_2^2)D_2A_2(D_1A_1 - D_2A_2)^{-1} - D_2^2),\ A_5A_0^{-1} = (D_1D_3)^{-1}((D_3^2 - D_1^2)D_1A_1(D_3A_4 - D_1A_1)^{-1} - D_1^2),\ A_6A_0^{-1} = (D_2D_3)^{-1}((D_2^2 - D_3^2)D_2A_2(D_2A_2 - D_3A_4)^{-1} - D_2^2)$; thus with $\square := (D_2^2 - D_3^2)D_1A_1 + (D_3^2 - D_1^2)D_2A_2 + (D_1^2 - D_2^2)D_3A_4$ we have $(D_2A_3A_0^{-1} - D_3A_5A_0^{-1})^{-1}A_1 = ((D_1^2 - D_2^2)(D_1A_1 - D_2A_2)^{-1} - (D_3^2 - D_1^2)(D_3A_4 - D_1A_1)^{-1})^{-1} = (D_1A_1 - D_2A_2)\square^{-1}(D_3A_4 - D_1A_1)$ and similarly $(D_3A_6A_0^{-1} - D_1A_3A_0^{-1})^{-1}A_2 = (D_1A_1 - D_2A_2)\square^{-1}(D_2A_2 - D_3A_4)$. Now $D_1((D_2^2 - D_3^2)D_2A_3A_0^{-1}(D_2A_3A_0^{-1} - D_3A_5A_0^{-1})^{-1}A_1 - D_2^2A_1) = (D_1^2D_2A_2 - D_2^2D_1A_1)\square^{-1}(D_2^2 - D_3^2)(D_3A_4 - D_1A_1) - D_2^2D_1A_1 = (D_1^2D_2A_2 - D_2^2D_1A_1)\square^{-1}(D_3^2 - D_1^2)(D_2A_2 - D_3A_4) - D_1^2D_2A_2 = D_2((D_3^2 - D_1^2)D_1A_3A_0^{-1}(D_3A_6A_0^{-1} - D_1A_3A_0^{-1})^{-1}A_2 - D_1^2A_2)$, so the very lhs and rhs provide the good definition of and afford themselves the name $D_1D_2D_3A_7$.

6.3. **The Bianchi-Lie approach.**

Following the Bianchi-Lie approach in generalizing the B transformation of the 2-dimensional pseudo-sphere to B transformation of an arbitrary 2-dimensional quadric, it is natural to ask what are the leaves of the considered distribution of isoclinic subspaces when the CGC -1 seed x^0 is in the particular position of being the actual n-dimensional pseudo-sphere. Once the correct metric-projective formulation is found, all that is left to do is the analytic confirmation of its generalization to arbitrary quadrics. If the CGC -1 seed is in the particular position of actually being the pseudo-sphere, then the facets of the considered distribution are situated on confocal pseudo-spheres; this constitutes a strong indication if favor of the Bianchi-Lie approach. However, the Tenenblat-Terng approach has to be slightly modified, as for the n-dimensional pseudo-sphere $|x_0|^2 = -1$ in the Lorentz space $\mathbb{R}^n \times i\mathbb{R}$ the shape is highly degenerated and one cannot prescribe the orthogonal sub-manifold A.

Let $x_0 := -ie_0 + \rho(ie_0 + u),\ \rho := \frac{2}{1-|u|^2},\ u := \sum_{j=1}^n u^j e_j \in \mathbb{C}^n$ be a parametrization of the pseudo-sphere $x_0 \subset \mathbb{C}^{n+1},\ |x_0|^2 = -1$ with asymptotic isotropic cone $|u|^2 = 1$. Note that u may not vary in all of $\mathbb{C}^n$; it may depend on k complex and $n-k$ real parameters such that du is n-dimensional; $\mathbf{H}^n(\mathbb{R})$ is obtained for $u \in \mathbb{R}^n,\ |u|^2 < 1$. We have $d\rho = \rho^2 u^T du$; the unit normal is ix_0, $dx_0 = (ie_0 + u)d\rho + \rho du,\ |dx_0|^2 = \rho^2|du|^2 = -ix_0^T d^2x_0$.

Again it is convenient to introduce a 0 column vector as the first one of an orthonormal frame in the normal bundle; all normal frames will be of the form $X'' \in \mathbf{M}_{2n-1,n}(\mathbb{C})$, $X''e_1 = 0$, $X''^T X'' = I_{2,n}$ and all computations with orthonormal frames remain valid as long as one uses the transpose instead of inverse.

We have $|\partial_{u^j} x_0| = \rho$, $\partial_{u^j} x_0 = \rho(e_j + \rho u^j(ie_0 + u))$, $X_0' := \rho^{-1}[\partial_{u^1} x_0 ... \partial_{u^n} x_0] \in \mathbf{M}_{2n-1,n}(\mathbb{C})$ moving tangent orthonormal frame along x_0 and $X_0'' := [0\ ix_0\ e_{n+1}...e_{2n-2}] \in \mathbf{M}_{2n-1,n}(\mathbb{C})$ moving orthonormal frame in the normal bundle of x_0, $X_0 := [X_0'\ X_0'']$. Further $\omega_0 := X_0'^T dx_0 = \rho du$, $\omega_0' := X_0'^T dX_0' = \rho[u, du]$, $\omega_0'' := X_0'^T dX_0'' = -(X_0''^T dX_0')^T = \rho[0\ idu\ 0\ ...\ 0]$, $\tilde{\omega}_0 := X_0''^T dX_0'' = 0$, $\Omega_0' := d\wedge\omega_0' + \omega_0'\wedge\omega_0' = -\rho^2 du\wedge du^T = -\omega_0\wedge\omega_0^T$, $d\wedge\omega_0 = -\omega_0'\wedge\omega_0$, $\omega_0''^T\wedge\omega_0 = 0$, $\omega_0'^T = -\omega_0'$, $\omega_0'' e_1 = 0$ and we have the GCMR equations

$$\Omega_0' - \omega_0''\wedge\omega_0''^T = 0,\ d\wedge\omega_0'' + \omega_0'\wedge\omega_0'' = 0,\ \omega_0''^T\wedge\omega_0'' = 0.$$

Consider the rolling $(R,t)(x_0, dx_0) = (x, dx)$, $x \subset \mathbb{C}^{2n-1}$ with flat normal bundle; R is chosen so that the normal connection of x is not only flat, but further trivial: $X_0''^T R^T dR X_0'' = 0$.

The corresponding omegas for x have same definition and properties as those of x_0, but with the index 0 removed; the frames of x are those of x_0 multiplied on the left by R. We have $\omega = \omega_0$; from (4.5) we have $\omega' = (RX_0')^T d(RX_0') = \omega_0'$, so $(RX_0)^T dRX_0 = (RX_0)^T d(RX_0) - X_0^T dX_0 = \begin{bmatrix} 0 & \omega'' - \omega_0'' \\ -(\omega'' - \omega_0'')^T & 0 \end{bmatrix}$.

The distribution of n-dimensional facets centered at $x_0 + \sin\sigma X_0' A^T e_1$, $A \in \mathbf{O}_n(\mathbb{C})$, spanned by frames $\sin\sigma(X_0' A^T DJ - X_0'')$ and with normal frames $\sin\sigma(X_0' A^T I_{2,n} + X_0'' D)$ should remain integrable if we roll it as we roll x_0 on x and for any position of x.

. Should such leaves exist, we would have $0 = (X_0' A^T I_{2,n} + X_0'' D)^T R^T d(x + \sin\sigma R X_0' A^T e_1) = 0$, or $-D\omega''^T A^T e_1 + I_{2,n} A(\csc\sigma\omega_0 + \omega_0' A^T e_1 - A^T dA A^T e_1) = 0$. With $K := \frac{1}{2} A\omega'' D$, $L := \frac{1}{2} I_{2,n} A\omega_0 e_1^T = L e_1 e_1^T$ we have $Ke_1 = L^T e_1 = 0$, $LI_{2,n} = 0$ and the last relation becomes

$$\begin{aligned} dAA^T &= A\omega_0' A^T + K - K^T + \csc\sigma(L - L^T) + N, \\ N^T &= -N,\ Ne_1 = (-K^T + \csc\sigma L)e_1. \end{aligned} \tag{6.8}$$

From (6.8) and the GW, GCMR equations of x we get

$$\begin{aligned} &K^T\wedge K = L\wedge L = L^T\wedge L = L\wedge K^T = 0,\ K^T\wedge A\omega_0 = 0,\ K\wedge K^T = -\frac{\cot^2\sigma}{4}(A\omega_0)\wedge(A\omega_0)^T, \\ &d\wedge K = (K + \csc\sigma(L - L^T) + N)\wedge K,\ d\wedge L = \csc\sigma L\wedge A\omega_0 e_1^T + (K + K^T - e_1 e_1^T K + N)\wedge L. \end{aligned} \tag{6.9}$$

Imposing the $d\wedge$ condition on (6.8)A, using (6.8) itself, (6.9) and the intrinsic part of the Gauß equation we get

$$\begin{aligned} &d\wedge N = -(K + \csc\sigma L + N)\wedge(K + \csc\sigma L + N)^T + (A\omega_0)\wedge(A\omega_0)^T + \\ &\csc\sigma((e_1 e_1^T K)\wedge L + L^T\wedge(e_1 e_1^T K)^T) - \csc^2\sigma(L\wedge(A\omega_0 e_1^T) + (A\omega_0 e_1^T)^T\wedge L^T). \end{aligned} \tag{6.10}$$

Further imposing the $d\wedge$ condition on (6.10) and using the equation itself, (6.8) and (6.9) we obtain only further (possibly vacuous) algebraic restrictions on N. Now those conditions may make (6.8) into a totally integrable Ricatti equation in A, in which case the leaves in the particular position of the Lie ansatz reveal themselves by letting $\omega'' := \omega_0''$ and integrating that equation (hopefully such a venue is possible).

6.4. **Particular deformations in $\mathbb{C}^{2n-1}$ of quadrics in $\mathbb{C}^{n+1}$.**

7. Appendix 1: A gist of deformations of quadrics

7.1. **Introduction.**

Quadratic laws and conserved quantities (like the energy or the linear element of a surface) are metric-projective notions (differentials involve tangency, a projective notion). It is thus no accident that confocal quadrics (having a metric-projective definition) behave well with respect to quadratic laws and conserved quantities.

We shall present a partial outline of the theory of deformations (through bending) of quadrics, namely the existence of the Bäcklund transformation for deformations of quadrics and the applicability (isometric) correspondence provided by the Ivory affinity. The analysis naturally splits into *static* (algebraic) computations on quadrics confocal to the given one and *moving* (differential) computations in its tangent bundle; the flat connection form is provided by rolling. All necessary identities of the moving part boil down to valid relevant algebraic identities of the static part; in turn these algebraic identities naturally appear within the static picture if one applies the the method of Archimedes-Bianchi-Lie in the moving picture.

7.2. **Confocal quadrics.**

Consider $\mathbb{R}^3$ with the standard basis e_1, e_2, e_3 and the usual Euclidean scalar product $< x, y >:= x^T y,\ |x|^2 := x^T x$. The family of quadrics $\{x_z\}_z$ confocal ('with same foci') to a given one $x_0 \subset \mathbb{R}^3$ is by definition the adjugate of the pencil of quadrics generated by the Cayley's absolute $\mathbb{CP}^2 \supset C(\infty) \ni [y\ 0]$: $\begin{bmatrix} y \\ 0 \end{bmatrix}^T \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y \\ 0 \end{bmatrix} = 0$ and the adjugate of the initial quadric x_0: $\begin{bmatrix} x_z \\ 1 \end{bmatrix}^T (\begin{bmatrix} A & 0 \\ 0 & -1 \end{bmatrix}^{-1} - z \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix})^{-1} \begin{bmatrix} x_z \\ 1 \end{bmatrix} = 0$ for $A := \mathrm{diag}[a_1^{-1}\ a_2^{-1}\ a_3^{-1}]$ (ellipsoids and hyperboloids with one or two sheets) or $\begin{bmatrix} x_z \\ 1 \end{bmatrix}^T (\begin{bmatrix} A & -e_3 \\ -e_3^T & 0 \end{bmatrix}^{-1} - z \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix})^{-1} \begin{bmatrix} x_z \\ 1 \end{bmatrix} = 0$ for $A := \mathrm{diag}[a_1^{-1}\ a_2^{-1}\ 0]$ (elliptic or hyperbolic paraboloids).

Since rigid motions and homotheties are the only projective transformations (homographies) which preserve $C(\infty)$, it encodes the metric structure of $\mathbb{R}^3$ and the mixed metric-projective character of the definition of confocal quadrics becomes clear.

Through any point $x \in \mathbb{R}^3$ pass 3 quadrics of the confocal family: $x \in x_{z_j},\ j = 1, 2, 3$; replacing the cartesian coordinates of x with $(z_j)_{j=1,2,3}$ gives elliptic coordinates on $\mathbb{R}^3$ (for quadrics of revolution we have less than three z's; we replace in this case one z with the coordinate induced by rotation or two z's with the spherical coordinates for confocal spheres).

The classical theorems about confocal quadrics are as follows:

(Lamé) The normal to x_z is proportional to $\partial_z x_z$ and the vectors $(\partial_z|_{z=z_j} x_{z_j})_{j=1,2,3}$ are orthogonal; thus the family of quadrics confocal to a given general quadric forms an orthogonal system.

(Dupin) Confocal quadrics cut each other along lines of curvature (they are given by the elliptic coordinates).

(Ivory) The orthogonal trajectory of a point on x_z as z varies is a conic, and the correspondence $x_0 \to x_z$ thus established is affine.

This affine transformation (henceforth called the Ivory affinity) preserves the lengths of segments between confocal quadrics: with $V_0^1 := x_z^1 - x_0^0,\ V_1^0 := x_z^0 - x_0^1$ we have $|V_0^1|^2 = |V_1^0|^2$ for pairs of points $(x_0^0, x_z^0),\ (x_0^1, x_z^1)$ corresponding on (x_0, x_z) under the Ivory affinity.



(Bianchi) If we have the rulings $w_0^0,\ w_0^1$ at the points $x_0^0,\ x_0^1 \in x_0$ and by use of the Ivory affinity we get the rulings $w_z^0,\ w_z^1$ at the points $x_z^0,\ x_z^1 \in x_z$, then $[V_0^1\ \ w_0^0\ \ w_z^1]^T[V_0^1\ \ w_0^0\ \ w_z^1] = [-V_1^0\ \ w_z^0\ \ w_0^1]^T[-V_1^0\ \ w_z^0\ \ w_0^1]$, so there exists a rigid motion $(R_0^1, t_0^1) \in \mathbf{O}_3(\mathbb{R}) \ltimes \mathbb{R}^3$ with

$$(R_0^1, t_0^1)(x_0^0, x_z^1, w_0^0, w_z^1) = (x_z^0, x_0^1, w_z^0, w_0^1). \tag{7.1}$$

Moreover $(V_0^1)^T\partial_z|_{z=0}x_z^0 = (V_1^0)^T\partial_z|_{z=0}x_z^1$, so the Ivory affinity has a nice projective property: the symmetry of the tangency configuration

$$x_z^1 \in T_{x_0^0}x_0 \Leftrightarrow x_z^0 \in T_{x_0^1}x_0. \tag{7.2}$$

(Jacobi) The tangent lines to a geodesic on x_0 remain tangent to another confocal quadric.

(Chasles) The common tangents to two confocal quadrics form a normal congruence and envelope geodesics on the two confocal quadrics.

There are many transformations of $\mathbb{R}^3$ which take a family of confocal quadrics to another one, but a transformation which preserves the previous properties must take lines to lines, so it must be a homography; conversely:

(Bianchi) Most homographies take a family of confocal quadrics to another one.

All quadrics are doubly ruled because they are equivalent from a complex projective point of view, but the only doubly ruled real quadrics are the hyperboloid with 1 sheet

$$x_z(u,v) := \sqrt{a_1 - z}\frac{1-uv}{u-v}e_1 + \sqrt{z-a_2}\frac{1+uv}{u-v}e_2 + \sqrt{a_3-z}\frac{u+v}{u-v}e_3,\ a_2 < 0, z < a_1, a_3, \tag{7.3}$$

when the Ivory affinity is given by $x_z(u,v) = \sqrt{I_3 - zA}\ x_0(u,v)$ and the hyperbolic paraboloid

$$x_z(u,v) := \sqrt{a_1 - z}(u+v)e_1 + \sqrt{z-a_2}(u-v)e_2 + (2uv + \frac{z}{2})e_3,\ a_2 < 0, z < a_1, \tag{7.4}$$

when the Ivory affinity is given by $x_z(u,v) = \sqrt{I_3 - zA}\ x_0(u,v) + \frac{z}{2}e_3$.

7.3. **The Bäcklund transformation.**

Bäcklund constructed in 1883 a transformation for constant Gauß curvature -1 surfaces (this point of view is due to Lie): the tangent spaces to the unit *pseudo-sphere* x_0 (the space-like unit sphere in the Lorentz space $\mathbb{R}^2 \times i\mathbb{R}$; it is seen as time-like from the origin) cut a confocal pseudo-sphere x_z along circles, thus highlighting a circle in each tangent space of x_0. Each point of the circle, the segment joining it with the origin of the tangent space and one of the (imaginary) rulings on x_z passing through that point determine a *facet* (pair of a point and a plane passing through that point). We have thus highlighted a 3 dimensional integrable distribution of facets: its leaves are the ruling families on x_z. Consider a surface x (called *seed*) non-rigidly isometric (applicable) to the pseudo-sphere x_0. One can roll x_0 on x such that for each position of the rolled x_0 the rolled x_0 and x will meet tangentially and with the same differential at the tangency point.

If we roll the distribution while rolling x_0 on x (the rolled distribution is obtained as follows: each facet of the original distribution corresponds to a point on x_0; we act on that facet with the rigid motion of the rolling corresponding to the highlighted point of x_0 in order to obtain the corresponding facet of the rolled distribution), then it turns out that the integrability condition of the rolled distribution is always satisfied (we have complete integrability), so the integrability of the rolled distribution does not depend on the shape of the seed. Thus the rolled distribution is integrable and its leaves (called the Bäcklund transforms of x, denoted $B_z(x)$ and whose determination requires the integration of a Ricatti equation) will be applicable to the pseudo-sphere. Moreover the seed and any leaf are the focal surfaces of a Weingarten congruence (their second fundamental forms are proportional).

A-priori one would think that the Bäcklund transformation for the pseudo-sphere exists because the pseudo-sphere has a big group of symmetries and thus a general quadric might not admit such a transformation. But the fact that the Bäcklund transformation is of a general nature

(independent of the shape of the seed) means that its true nature lies not in the differential picture with the seed and the leaf, but in the algebraic picture with confocal pseudo-spheres where the Bäcklund transformation being independent of the shape of the seed is still valid (although the leaves degenerate from surfaces to rulings of x_z). Thus the vanishing of the shape of the seed x in the total integrability condition occurs not because of the symmetries of the pseudo-sphere, but because of general equations (the Gauß-Codazzi-Mainardi equations of the seed and of the pseudo-sphere) coupled with algebraic consequences of the tangency configuration and apparent at the level of confocal pseudo-spheres: these are valid for a general quadrics.

Thus Bianchi generalized in 1906 the picture of the Bäcklund transformation of the pseudo-sphere to arbitrary quadrics: the same statement remains true if in Lie's interpretation 'pseudo-sphere' is replaced with 'quadric' and 'circle' with 'conic'. However, the applicability correspondence on constant Gauß curvature surfaces (which can be found in ∞^3 ways) does not have an easy generalization to general quadrics. According to Bianchi's own account ([5],(122),IV) should the applicability correspondence exist in general, it can be presumed to be of a general nature and thus independent of the shape of the seed: therefore the answer lies not in the differential picture with the seed and the leaf, but in the algebraic picture with confocal quadrics so we have to roll back. Thus to find the applicability correspondence one must look for a natural correspondence between confocal quadrics. The Ivory affinity provides such a correspondence and proving the validity of the applicability correspondence via the Ivory affinity is very easy once the complete integrability of the rolled distribution is checked: if we roll the seed x^0 on x_0^0, then the tangent space of the leaf x^1 corresponding to the point of tangency of the rolled x_0^0 and the seed x^0 will be applied to the facet centered at x_z^1 and spanned by $V_0^1,\ x_{zu_1}^1$; further applying the rigid motion (R_0^1, t_0^1) provided by the Ivory affinity it will be applied to $T_{x_0^1}x_0$. In this process $x_{u_1}^1$ is taken to $x_{zu_1}^1$ and further to $x_{0u_1}^1$, so actually (x^1, dx^1) is taken to (x_0^1, dx_0^1); moreover because of the symmetry of the tangency configuration the seed becomes leaf and the leaf becomes seed.

Although the applicability correspondence is restricted, the structure becomes richer (a quadric with less symmetries has less degeneracies; for example a segment of Bäcklund transformations degenerates to the complementary transformation for a quadric of revolution, or the geodesic flow on an ellipsoid degenerates to the one on a sphere).

7.4. **Two algebraic consequences of the tangency configuration.**
Consider now the tangency configuration $(V_0^1)^T N_0^0 = 0$; let $\mathcal{B} := (u-v)^2$ for (7.3) and $\mathcal{B} := 1$ for (7.4), $m_0^1 := \mathcal{B}_1 x_{zu_1}^1 \times V_0^1$ a normal field of the distribution $\mathcal{D}^1$ of facets $\mathcal{F}^1$ passing through x_z^1 and spanned by $V_0^1,\ x_{zu_1}^1$ (and similarly $m_0'^1 := \mathcal{B}_1 x_{zv_1}^1 \times V_0^1$ by considering the other ruling family on x_z^1); note that m_0^1 depends only on u_0, v_0, v_1 and quadratically in v_1; this will make the integrability condition a Ricatti equation in v_1.

Consider the rulings $w_0^0 := x_{0u_0}^0,\ w_0^1 := x_{0u_1}^1$ at $x_0^0 := x_0(u_0, v_0),\ x_0^1 := x_0(u_1, v_1)$; we thus get a rigid motion (R_0^1, t_0^1) provided by the Ivory affinity. If we change the ruling family on x_0^1 the action of the new rigid motion on the facet $T_{x_0^0}x_0$ does not change, so its new rotation must be the old rotation composed with a reflection in $T_{x_0^0}x_0$, because of which the facets $\mathcal{F}^1,\ \mathcal{F}'^1$ reflect in $T_{x_0^0}x_0$ (thus the distributions $\mathcal{D}^1,\ \mathcal{D}'^1$ reflect in Tx_0):

$$(x_{zv_1}^1)^T (I_3 - 2N_0^0 (N_0^0)^T) m_0^1 = 0, \tag{7.5}$$

and $x_{0v_1}^1 = R_0^1 (I_3 - 2N_0^0 (N_0^0)^T) x_{zv_1}^1$; multiplying this on the left by $(x_{0u_1}^1)^T$ and using the preservation of lengths of rulings under the Ivory affinity we get

$$4(x_{zu_1}^1)^T N_0^0 (N_0^0)^T x_{zv_1}^1 du_1 dv_1 = |dx_z^1|^2 - |dx_0^1|^2 = -\frac{4z}{\mathcal{B}_1} du_1 dv_1. \tag{7.6}$$

Thus we have the next result, essentially due to Bianchi (he uses equivalent computations):

If $x_z^1 \in T_{x_0^0}x_0$, then:

I The change in the linear element from x_z^1 to x_0^1 is four times the product of the orthogonal projections of the differentials of the rulings of x_z^1 on the normal of x_0 at x_0^0.

II The facets at x^1_z spanned by V^1_0 and one of the rulings of x^1_z reflect in $T_{x^0_0}x_0$; therefore the distributions $\mathcal{D}^1$, $\mathcal{D}'^1$ reflect in Tx^0_0.

The algebraic relation

$$(N^0_0)^T(2zm^1_0 + m^1_0 \times m^1_{0v_1}) = 0 \tag{7.7}$$

will appear as the total integrability condition. Using (7.5), (7.6) this becomes: $0 = \frac{z(m^1_0)^T x^1_{zv_1}}{(N^0_0)^T x^1_{zv_1}} - \mathcal{B}_1(x^1_{zu_1})^T N^0_0 (V^1_0)^T m^1_{0v_1} = \frac{z(V^1_0)^T(\mathcal{B}_1 x^1_{zv_1} \times x^1_{zu_1} + (\mathcal{B}_1 x^1_{zu_1} \times V^1_0)_{v_1})}{(N^0_0)^T x^1_{zv_1}}$, which is straightforward. Replacing (m^1_0, v_1) with (m'^1_0, u_1) we get a similar relation.

7.5. **Rolling surfaces and distributions.**
Let $(R_0, t_0)(x^0_0, dx^0_0) = (x^0, dx^0)$ be the rolling of the quadric $x^0_0 = x_0(u_0, v_0)$ on the applicable (rollable) surface $x^0 \subset \mathbb{R}^3$ (in general $(R_0, t_0) \subset \mathbf{O}_3(\mathbb{R}) \ltimes \mathbb{R}^3$ is a surface, but it degenerates to a curve if x^0 is ruled); the facets of the rolled distribution $(R_0, t_0)\mathcal{D}^1$ will become tangent spaces to surfaces (leaves) $x^1 := (R_0, t_0)x^1_z = (R_0, x^0)V^1_0$ iff the integrability condition $0 = (R_0 m^1_0)^T dx^1$ holds. Under the identification $(\mathbf{o}_3(\mathbb{R}), [,]) \simeq (\mathbb{R}^3, \times)$ we have $R_0^{-1}dR_0 \simeq N^0_0 \times R_0^{-1}dR_0 N^0_0 =: \omega_0$; further imposing the compatibility condition $d\wedge$ to $R_0 dx^0_0 = dx^0$ (we shall use the notation $d\wedge$ for exterior derivative and d for tensorial derivative) we get

$$d\wedge\omega_0 + \frac{1}{2}\omega_0 \times \wedge\omega_0 = 0,\ \omega_0 \times \wedge dx^0_0 = 0 \tag{7.8}$$

and thus ω_0 is a flat connection form in Tx^0_0 (it encodes the difference between the Gauß - Codazzi-Mainardi equations for x^0, x^0_0).

We have $R_0^{-1}dx_1 = d(x^0_0 + V^1_0) + R_0^{-1}dR_0 V^1_0 = dx^1_z + \omega_0 \times V^1_0$. But $(\omega_0)^\perp = 0$ and $dx^1_z = x^1_{zv_1}dv_1 + x^1_{zu_1}du_1$, so the integrability condition becomes $-(V^1_0)^T\omega_0 \times N^0_0(m^1_0)^T N^0_0 + (m^1_0)^T x^1_{zv_1}dv_1 = 0$; using (7.5) this becomes $-(V^1_0)^T\omega_0 \times N^0_0 + 2(N^0_0)^T x^1_{zv_1}dv_1 = 0$; multiplying it by $\mathcal{B}_1(N^0_0)^T x^1_{zu_1}$, using (7.6) and $-\mathcal{B}_1(N^0_0)^T x^1_{zu_1}V^1_0 = \mathcal{B}_1(V^1_0 \times x^1_{zu_1}) \times N^0_0 = -m^1_0 \times N^1_0$ we finally get the Ricatti equation:

$$(m^1_0)^T\omega_0 + 2zdv_1 = 0. \tag{7.9}$$

We have $dm^1_0 = m^1_{0v_1}dv_1 + \mathcal{B}_1 dx^0_0 \times x^1_{zu_1}$, so $(dm^1_0)^T \wedge \omega_0 = dv_1 \wedge (m_{0v_1})^T\omega_0$; imposing the total integrability condition $d\wedge$ on (7.9) and using the equation itself we need $-(m^1_0)^T\omega_0 \wedge (m^1_{0v_1})^T\omega_0 + 2z(m^1_0)^T d\wedge\omega_0 = 0$, or, using (7.8): $(N^0_0)^T(2zm^1_0 + m^1_0 \times m^1_{0v_1})(N^0_0)^T\omega_0 \times \wedge\omega_0 = 0$; therefore the total integrability is equivalent to (7.7)*.

7.6. **The method of Archimedes-Bianchi-Lie.**
If we roll x^0_0 on different sides of the seed x^0, then we get the Bäcklund transformation for the other ruling family, so the rolled distributions reflect in the tangent bundle of the seed x^0. Thus (7.5) is obtained if one makes the ansatz $x^0_0 = x^0$; the same ansatz for the applicability correspondence provided by the Ivory affinity and the inversion of the Bäcklund transformation (two focal surfaces of a line congruence are in a symmetric relationship) implies Bianchi's result about the existence of (R^1_0, t^1_0) and the symmetry of the tangency configuration; now (7.6) is obtained as previously described.

In the famous letter *The Method*, sent to Eratosthenes of the University of Alexandria and lost from the XIII$^{\text{th}}$ century until 1906 (the year of Bianchi's discovery, so it was unknown to Bianchi and Lie), Archimedes states: '*... certain things first became clear to me by a mechanical method,*

* Most of § 7.1-§ 7.5 was a 10 minutes presentation at a general MAA session in San Antonio, TX January 12 2006. Now in August 2005 when I registered for this general session I did not see a differential geometry AMS session; only algebraic geometry sessions (it turned out I was wrong; on the flyer I have seen several such differential geometry sessions). But the simplification from late December 2005-early January 2006 fortunately made it appropriate for such a MAA session (in fact while trying to present a simpler point of view for the MAA session I managed to implement the simplest point of view at that date); on top of that the link to Archimedes' method makes it even more so, since Archimedes' ideas are currently subject to further research in *The American Mathematical Monthly*. Thus the apparently wrong denomination stuck and fulfilled its purpose.

although they had to be proved by geometry afterwards because their investigation by the said method did not furnish an actual proof. But it is of course easier, when we have previously acquired, by the method, some knowledge of the questions, to supply the proof than it is to find it without any previous knowledge'.

Note that the method at the level of points of facets was known to Bianchi and Lie; however, they had never used the full method, at the level of the planes of the facets too (for this reason the *'of course easier'* ingredient is missing from Bianchi's proofs). Thus if one assumes Theorem I of Bianchi's theory of deformations of quadrics a-priori to be true and to be the metric-projective generalization of Lie's approach, then one naturally *geometrically* gets the necessary algebraic identities needed to prove Theorem I.

For surfaces this method is just a fancy way of reformulating already known identities and which appear naturally enough at the analytic level. But keep in mind that (7.2), (7.5) and (7.6) are equivalent from an analytic point of view and this is not the case in higher dimensions: thus it is very difficult to find the necessary algebraic identities of the static picture from an analytic point of view. Therefore the method of Archimedes-Bianchi-Lie, due to its geometric naturalness is useful in the study of higher dimensional problems. Note that although Archimedes and Bianchi-Lie have dealt with different problems, their approach was the same: $P \Rightarrow Q$ with P being either *'The area of a segment of a parabola is an infinite sum of areas of lines'* or *'A line is a deformation of a quadric'*. Such sentences P were not accepted as true according to the standard of proof of the times, but they had valid relevant consequences Q which elegantly solved problems not solvable with other methods of those times; if only for this fact the standard of proof should change so as to accept these statements as true. Note that *The Method* of Archimedes went a little closer to the B transformation: in an a-priori intuitive elementary non-rigorous geometric argument he transferred and stuck all lines (slices of the segment of the parabola) with their centers at the left end of the balance (thus with ∞ multiplicity).

Note that the facets of the 3-dimensional rolled distribution are differently re-distributed into 2-dimensional families of facets as tangent planes to leaves when the shape of the seed changes, but principles and properties independent of the shape of the seed remain valid for facets even in the singular picture: this is Archimedes' contribution to the Bianchi-Lie ansatz.

Thus I can honestly say that the *B transformation (for quadrics)* is *one and the same with* the *Archimedes balance* as *principles of a general nature*: they are valid at the infinitesimal level and survive integration and conversely, being principles of a general nature both induce by differentiation and by particular singular configurations the infinitesimal picture where the simplest explanation of these principles reveals itself.

The natural question thus appears wether Google concurs with this finding or not and indeed Google points to the site where the truth (*'The truth is out there!'*) is spelled out: *'In the same way that Graham was a Man of Ego who looked like a champion, Backlund was a Man of Skill who looked like something completely different. Bob Backlund didn't even look like a wrestler, he looked like your shop teacher from school, or Opie Cunningham grown up. Backlund's motto was that 'any move can be countered, any hold can be broken.' Like Archimedes, Backlund declared, 'Give me where to stand, and I will move the earth,' making of his own body, his lever, and of the wrestling ring, his fulcrum.'*. All Bäcklund needed was a seed and he produced lots of leafs by means of his method. Note also that on this site Albert Victor Bäcklund clearly appears as *an immediate follower* of Archimedes; note also that Bäcklund got his undergraduate education at Lund, then travelled (including in Europe, where he met with his conational Lie's ideas about the contact transformation) and then he went *back to Lund* for the remainder of his life. While Opie Cunningham was a friend of Fonzie's, it seems to me that apparently Bob ain't no friend of Bäcklund's I ever heard of; only that Robert Miura is aka as Bob (not necessarily English Bob) and he declared in ([23], pg 429) in 1976 as to regards to virtually the B transformation of the KdV equation: *'.... However, several simplifications have been achieved, not least of which is that all the equations to be solved are linear!... It is instructive to point out two simplifying features*

that make the method work. First, we were able to show that the discrete eigenvalues are constant. Second, we could compute the time dependence of c_n and b before knowing the time-dependence of the solution u, requiring only that we knew u at $t = 0$.... (We liken this situation to the spirit of Archimedes' statement 'Give me but one firm spot on which to stand and I will move the earth.')'. I can honestly say that Bob is actually pointing out Archimedes' method: first some useful Euclidean identities are derived (corresponding to constant eigenvalues), then we assume the method to work before we know the method actually works and requiring in the process nothing more than some partial initial knowledge of the method working (simple consequences of the method) and *that is all there is to it*; the checking of the fact that the method works brings nothing more besides tedious computations. Post-priori the partial knowledge needed to prove the general configuration clearly appears from the singular configuration by the application of the same method. Note however that Bob (incarnated both as the Wrestling Bob (*'His name is Robert Paulson! His name is Robert Paulsen!'*) and as the Mathematician Bob) was right about the balance: *'That hit the spot!'*. I can also honestly say that if the Mathematician Bob were familiar with Archimedes' method (for example it being taught in high-school), then he would have probably made the connection in 1976 or earlier. Note also that Bianchi in (122,§ 10) was on spot, as he uses angles and moments of pairs of lines to explain the RMPIA; thus his moments provides the physical link to Archimedes' moments of the balance.

Jules's *last minute* interpretation of *Ezekiel 12:57* (*'What is significant is that I felt the touch of God. God got involved. ... First, I'm going to deliver this case to Marsellus. Then, basically I'm just going to walk The Earth. ... Like Caine in 'Kung Fu'. Going from place to place, meeting new people, getting in adventures. ... replaying the incident in my head when I had what alcoholics refer to as a moment of clarity. ... Blessed is he who in the name of charity and good will shepherds the weak through the Valley of Darkness, for he is truly his brother's keeper and the finder of lost children. ... Or maybe it is the world who is evil and selfish. ... I am the tyranny of an evil man, but I am trying Ringo, I am trying really hard to be the shepherd'*) reveals itself: nothing is more powerful than a simple stick drawing a picture on the sand of a forgotten beach only to be found by the survivors of a nearby unfortunate shipwreck: *'Bene speremus, Hominum enim vestigia video!'*.

'Qui la Geometria vive!' Rock'n Roll The Rock & Peace out!

References

[1] Archimedes *The Method.*
[2] V. I. Arnold *On Teaching Mathematics.*
[3] Berger, E., Bryant, R. L., Griffiths, P. A. *The Gauß equations and rigidity of isometric embeddings,* Duke Math. J., **50** (1983), 803-892.
[4] L. Bianchi *Lezioni Di Geometria Differenziale,* Vol **1-4**, Nicola Zanichelli Editore, Bologna (1922-27).
[5] L. Bianchi *Opere,* Vol **1-11**, a cura dell'Unione Matematica Italiana e col contributo del Consiglio Nazionale Delle Richerche, Edizioni Cremonese (1952-59).
[6] A. I. Bobenko, U. Pinkall *Discrete isothermic surfaces,* J. reine angew. Math., **475** (1996), 209-220.
[7] A. I. Bobenko, U. Pinkall *Discrete surfaces with constant negative Gaussian curvature and the Hirota equation,* J. Diff. Geom., **43** (1996), no. 3, 527-611.
[8] C. B. Boyer *A History of Mathematics,* John Wiley & Sons, Inc. (1968).
[9] F. E. Burstall *Isothermic Surfaces: Conformal Geometry, Clifford Algebras and Integrable Systems,* preprint arXiv:math.DG/0003096 v2 11 Oct 2000.
[10] F. E. Burstall, D. Ferus, F. Pedit and U. Pinkall *Harmonic tori in symmetric spaces and commuting Hamiltonian systems on loop algebras,* Ann. of Math., **138** (1993), 173-212.
[11] P. Calapso *Intorno alle superficie applicabili sulle quadriche ed alle loro transformazioni,* Annali di Matematica Pura ed Applicada, **3**, XIX, (1912), 61-82, 102-178.
[12] E. Cartan *Sur les variétées de courboure constante d'un espace euclidien ou non-euclidien,* Bull. Soc. Math. France, **47** (1919), 125-160; and **48** (1920), 132-208.
[13] G. Darboux *Leçons Sur La Théorie Générale Des Surfaces,* Vol **1-4**, Gauthier-Villars, Paris (1894-1917).
[14] G. Darboux *Leçons Sur Les Systèmes Orthogonaux Et Les Coordonées Curvilignes,* Gauthier-Villars, Paris (1910).
[15] G. Darboux *Sur la déformation des surfaces du second degré,* Comptes rendus de l'Académie, Vol CXXVIII, **13**, (1899), 760-766, 854-859.
[16] G. Darboux *Sur la déformation des surfaces générales du second degré,* Comptes rendus de l'Académie, Vol CXXVIII, **21**, (1899), 1264-1270.
[17] G. Darboux *Sur les surfaces isothermiques,* Comptes rendus de l'Académie, Vol CXXVIII, **22**, (1899), 1299-1305.
[18] G. Darboux *Sur une classe de surfaces isothermiques liées à la déformation des surfaces du second degré,* Comptes rendus de l'Académie, Vol CXXVIII, **25**, (1899), 1483-1487.
[19] L.P. Eisenhart *A Treatise on the Differential Geometry of Curves and Surfaces,* Dover Publications, Inc., New York, New York (1909, republished 1960).
[20] L.P. Eisenhart *Transformations of Surfaces,* Princeton University Press, Princeton, (1922).
[21] T. L. Heath *APOLLONIUS OF PERGA Treatise on conic sections,* New York Barnes & Noble Inc, (1961) (reprinted from publisher Cambridge University Press, (1896)).
[22] T. Levi-Civita *The Absolute Differential Calculus (Calculus of Tensors),* Dover Publications, Inc., New York, New York (1977).
[23] R. M. Miura *The Korteweg-de Vries Equation: A Survey of Results,* SIAM Review, Vol. 18, (July 1976), pp. 412-459.
[24] J. Moser *Geometry of Quadrics and Spectral Theory,* The Chern Symposium 1979 Proc. Internat. Sympos., Berkeley, Calif., 1979, pp. 147-188, Springer, New York-Berlin, 1980.
[25] U. Pinkall, I. Sterling, *On the classification of constant mean curvature tori,* Ann. Math., **130** (1989), 407-451.
[26] C. Rogers, W. K. Schieff *Bäcklund and Darboux Transformations; Geometry and Modern Applications in Soliton Theory,* Cambridge Texts in Applied Mathematics, Cambridge University Press, 2002.
[27] I. Kh. Sabitov *Local Theory of Bendings of Surfaces,* Encyclopedia of Mathematical Sciences, **48**, Geometry III, Yu. D. Burago, V. A. Zalgaller Eds., Springer-Verlag (1991), 179-250.
[28] M. Servant *Sur la déformation des surfaces,* Comptes rendus de l'Académie, Vol CXXXVI, **21**, (1903), 1239-1241.
[29] M. Spivak *A Comprehensive Introduction to Differential Geometry,* Vol **1-5**, Publish or Perish, Berkeley, CA, (1975-80).
[30] A. Sym *Soliton surfaces and their applications,* Lecture Notes in Physics **239**, Springer-Verlag, Berlin (1985), 154-231.
[31] Q. Tarantino *Pulp Fiction,* Miramax Films & A Band Apart & Jersey Films Productions (1994).
[32] K. Tenenblat and C.-L. Terng *Bäcklund's theorem for n-dimensional sub-manifolds of* $\mathbb{R}^{2n-1}$, Ann. of Math., **111** (1980), 477-490.
[33] C.-L. Terng *A higher dimension generalization of the Sine-Gordon equation and its soliton theory,* Ann. of Math., **111** (1980), 491-510.

[34] C.-L. Terng, G. Thorbergsson *Submanifold Geometry in Symmetric Spaces,* J. Diff. Geom. **42** (1995), no. 3, 665-717.
[35] C.-L. Terng, K. Uhlenbeck *Poisson Actions and Scattering Theory for Integrable Systems,* arXiv:dg-ga/9707004 v1 7 Jul 1997.
[36] C.-L. Terng, K. Uhlenbeck *Bäcklund Transformations and Loop Group Actions,* Comm. Pure. & Appl. Math., **LIII** (2000), 1-75.
[37] G. Ţiţeica *Sur la déformation de certaines surfaces tétraédrales, les surfaces S et les réseaux R,* Geometria Proiettiva Differenziale Vol **2**, G. Fubini, E. Čech, Nicola Zanichelli Editore, Bologna 1927, 663-668.
[38] J. Weingarten *Sur la déformation des surfaces,* Acta Mathematica **20** 1896, 159-200.
[39] *The MacTutor History of Mathematics archive.*

Slobozia 2, Ţăndărei, Ialomiţa, 8454, România
E-mail address: ionidinca@yahoo.com